



HAL
open science

Liver fibrosis and all-cause mortality in chronic HCV-infected diabetic patients: A paradoxical association? (ANRS CO22 HEPATHER)

Tangui Barré, Clémence Ramier, Vincent Di Beo, Fabrice Carrat, Hélène Fontaine, Fabienne Marcellin, Patrizia Carrieri, Stanislas Pol, Camelia Protopopescu

► To cite this version:

Tangui Barré, Clémence Ramier, Vincent Di Beo, Fabrice Carrat, Hélène Fontaine, et al.. Liver fibrosis and all-cause mortality in chronic HCV-infected diabetic patients: A paradoxical association? (ANRS CO22 HEPATHER). *Liver International*, 2021, 41 (7), pp.1694 - 1698. 10.1111/liv.14949 . inserm-03332288v2

HAL Id: inserm-03332288

<https://inserm.hal.science/inserm-03332288v2>

Submitted on 30 Aug 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1
2
3
4 **Liver fibrosis and all-cause mortality in chronic HCV-infected diabetic patients:**
5 **a paradoxical association? (ANRS CO22 HEPATHER)**
6

7 Tangui Barré¹, Clémence Ramier¹, Vincent Di Beo¹, Fabrice Carrat^{2,3}, Hélène
8 Fontaine^{4,5}, Fabienne Marcellin¹, Patrizia Carrieri^{1#}, Stanislas Pol², Camelia Protopopescu¹
9 on behalf of the ANRS/AFEF Hepather study group
10
11

12 ¹Aix Marseille Univ, INSERM, IRD, SESSTIM, Sciences Economiques & Sociales de la
13 Santé & Traitement de l'Information Médicale, Marseille, France
14

15 ²Sorbonne Université, Institut National de la Santé et de la Recherche Médicale (INSERM),
16 Institut Pierre Louis d'Epidémiologie et de Santé Publique, Paris, France
17
18

19 ³Assistance Publique-Hôpitaux de Paris (AP-HP), Hôpital Saint-Antoine, Unité de Santé
20 Publique, Paris, France
21
22

23 ⁴AP-HP, Hôpital Cochin, Département d'Hépatologie, Paris, France
24
25

26 ⁵INSERM U1223, Institut Pasteur; Université Paris Descartes, Paris, France
27
28

29 ⁶Université de Paris ; AP-HP, Hôpital Cochin, Département d'Hépatologie; INSERM U1223,
30 Institut Pasteur, Paris, France
31
32

33
34
35
36 #Corresponding author
37
38

39 **Funding statement**
40
41
42
43

44 The cohort received financial support from the INSERM-ANRS (France Recherche
45 Nord & Sud Sida-HIV Hépatites), the French ANR (Agence Nationale de la Recherche), the
46 French DGS (Direction Générale de la Santé), Merck Sharp and Dohme, Janssen-Cilag,
47 Gilead, Abbvie, Bristol-Myers Squibb, and Roche. Those funding sources had no role in the
48 writing of the manuscript or the decision to submit it for publication.
49
50
51
52
53

54
55 FC reports grants from INSERM-ANRS, during the conduct of the study; and personal
56 fees from Imaxio, outside the submitted work. MB reports grants and personal fees from
57 AbbVie and Gilead, outside the submitted work; and personal fees from MSD, Janssen,
58
59
60

1
2
3 Boehringer Ingelheim, Intercept, and Bristol-Myers Squibb, outside the submitted work. SP
4
5 has received consulting and lecturing fees from Bristol-Myers Squibb, Janssen, Gilead, Roche,
6
7 MSD, MYR-Pharma, Shionogi, Biotest and Abbvie, as well as grants from Bristol-Myers
8
9 Squibb, Gilead, Roche and MSD. HF reports personal fees and invitations for medical
10
11 meetings from Gilead, AbbVie, Bristol-Myers Squibb, MSD, and Janssen, outside the
12
13 submitted work. Other authors declare no financial support.
14
15

16 17 Conflict of interest disclosure

18
19
20
21 The authors have no conflict of interest to disclose.
22
23

24 25 Ethics approval statement

26
27
28
29 Data were derived from a trial registered with ClinicalTrials.gov, number NCT01953458.
30
31 Written informed consent was obtained from each cohort participant before enrolment. The
32
33 protocol was designed in accordance with the Declaration of Helsinki and French law for
34
35 biomedical research, and was approved by the 'Comité de Protection des Personnes (CPP)
36
37 Ile de France 3' Ethics Committee (Paris, France) and the French Regulatory Authority
38
39 (ANSM).
40

41 42 Patient consent statement

43
44
45 Written informed consent was obtained from each cohort participant before enrolment.
46
47

48 49 Permission to reproduce material from other sources

50
51
52
53 There is no material from other sources.
54
55
56
57
58
59
60

Acknowledgments

We thank the study participants and the participating clinicians at each site. We also thank the INSERM-ANRS for sponsoring, funding and conducting the ANRS CO22 Hepather cohort in collaboration with the French Association for the Study of the liver (Association Française pour l'Etude du Foie: AFEF). Finally, our thanks to Jude Sweeney for the English revision and editing of our manuscript.

ANRS-AFEF Hepather study group

Delphine Bonnet, Virginie Sicart (CHU Purpan, Toulouse, France), François Bailly, Marjolaine Beaudoin, Dominique Giboz, Kerstin Hartig-Lavie, Marianne Maynard (Hospices Civils de Lyon, Lyon, France), Morane Cavellec, Marjorie Cheraud-Carpentier, François Raffi, Florian Vivrel (Hôpital Hôtel-Dieu, Nantes, France), Jaouad Benhida, Jérôme Boursier, Paul Calès, Françoise Lunel, Frédéric Oberti (CHU Angers, Angers, France), Nathalie Boyer, Audrey Gilibert, Nathalie Giuily (Hôpital Beaujon, Clichy, France), Giovanna Scoazec (Hôpital Beaujon, Clichy, France and Hôpital Henri Mondor, Créteil, France), Sandrine Fernandes, Sylvie Keser, Philippe Sultanik, Anaïs Vallet-Pichard (Hôpital Cochin, Paris, France), Juliette Foucher, Jean-Baptiste Hiriart, Aurore Mathias, Julien Vergniol (Hôpital Haut-Lévêque, Pessac, Bordeaux, France), Chrystelle Ansaldi, Laëtitia Chouquet, Emilie De Luca, Valérie Oules (Hôpital Saint Joseph, Marseille, France), Rodolphe Anty, Eve Gelsi, Régine Truchi (CHU de Nice, Nice, France), Elena Luckina, Nadia Messaoudi, Joseph Moussali, (Groupe Hospitalier Pitié-Salpêtrière, Paris, France), Barbara De Dieuleveult, Damien Labarriere, Pascal Poter, Si Nafa Si Ahmed (CHR La Source, Orléans, France), Nathalie Ganne-Carrié, Véronique Grando-Lemaire, Pierre Nahon, Alan Peltier, Judith Ung (Hôpital Jean Verdier, Bondy, France), Mariette Gougeon, Anne Guillygomarch, Caroline Jezequel (CHU Rennes, Rennes, France), Romain Moirand, Thomas F. Baumert, Michel

1
2
3 Dofföel, Catherine Mutter, Pauline Simo-Noumbissie (Hôpitaux Universitaires de Strasbourg,
4 Strasbourg, France), Hélène Barraud, Mouni Bensenane, Abdelbasset Nani, Sarah Hassani-
5 Nani (CHU de Nancy, Vandoeuvre-lès-Nancy, France), Marie-Albertine Bernard (CHU de
6 Nancy, Vandoeuvre-lès-Nancy, France and Centre Hospitalier Régional, Metz, France),
7
8 Michael Bismuth, Ludovic Caillo, Stéphanie Faure, Georges-Philippe Pageaux, Marie Pierre
9
10 Ripault (Hôpital Saint Eloi, Montpellier, France), Karl Barange, Christophe Bureau, Jean
11
12 Marie Peron, Marie Angèle Robic, Léa Tarallo (CHU Purpan, Toulouse, France), Marine
13
14 Faure, Bruno Froissart, Marie-Noelle Hilleret, Vincent Leroy (CHU de Grenoble, Grenoble,
15
16 France), Odile Gorla, Victorien Grard, Hélène Montialoux (CHU Charles Nicolle, Rouen,
17
18 France), Muriel François, Christian Ouedraogo, Christelle Pauleau, Anne Varault (Hôpital
19
20 Henri Mondor, Créteil, France), Tony Andreani, Bénédicte Angoulevant, Azeline Chevance,
21
22 Lawrence Serfaty (Hôpital Saint-Antoine, Paris, France), Teresa Antonini, Audrey Coilly,
23
24 Jean-Charles Duclos Vallée, Mariagrazia Tateo (Hôpital Paul Brousse, Villejuif, France),
25
26 Armand Abergel, Corinne Bonny, Chanteranne Brigitte, Géraldine Lamblin, Léon Muti
27
28 (Hôpital Estaing, Clermont-Ferrand, France), Abdenour Babouri, Virginie Filipe (Centre
29
30 Hospitalier Régional, Metz, France), Camille Barrault, Laurent Costes, Hervé Hagège,
31
32 Soraya Merbah (Centre Hospitalier Intercommunal, Créteil, France), Paul Carrier, Maryline
33
34 Debette-Gratien, Jérémie Jacques (CHU Limoges, Limoges, France), Florent Artu, Valérie
35
36 Canva, Sébastien Dharancy, Alexandre Louvet (CHRU Claude Huriez, Lille, France), Marc
37
38 Bardou, Donya Da Costa Souhiel, Patrick Hillon, Marianne Latournerie (Dijon University
39
40 Hospital, Dijon, France), Yannick Bacq, Didier Barbereau, Charlotte Nicolas (CHU
41
42 Trousseau, 37044 Tours, France), Nisserine Ben Amara, Danièle Botta-Fridlund, Isabelle
43
44 Portal (CHU Timone, Marseille, France), Moana Gelu-Simeon, Marie-Josée Lafrance (CHU
45
46 de Pointe-à-Pitre, Pointe-à-Pitre, Guadeloupe).
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 We read with great interest the article published by Ciancio *et al.*, which reported
4 long-term improvement of glycometabolic control in diabetic patients with chronic hepatitis C
5 following sustained virological response with direct-acting antivirals (DAA) (1). They also
6 reported a detrimental effect of baseline diabetes on hepatocellular carcinoma (HCC)
7 incidence in cirrhotic hepatitis C patients after multiple adjustment, and a higher mortality rate
8 in hepatitis C patients with diabetes than in their non-diabetic counterparts. The authors'
9 valuable data regarding the long-term benefits of hepatitis C virus clearance on glucose
10 homeostasis in diabetic patients, highlight the need to confirm: (i) that diabetic hepatitis C
11 patients present a higher risk of all-cause mortality, and (ii) whether specific mortality risk
12 factors should be given particular scrutiny in this sub-population. We might expect that the
13 higher rate of HCC incidence *Ciancio et al.* reported in diabetic HCV patients contributes to a
14 higher all-cause mortality rate, as was previously found before the DAA era (2,3). Diabetes is
15 also a well-known risk factor for cardiovascular disease, although DAA treatment appears to
16 diminish the risk (4).

17
18
19 We aimed to confirm whether diabetes is associated with higher all-cause mortality in
20 chronic HCV-infected patients in the DAA era, and whether specific risk factors affect all-
21 cause mortality differently in patients with and without diabetes. Our analyses were based on
22 longitudinal data from ANRS CO22 HEPATHER, a national, multicenter, prospective cohort
23 study conducted in France, which collected annual clinical and socio-behavioral data over
24 five years of follow-up in chronic HCV-infected patients who were not on treatment at
25 inclusion (5). Unlike *Ciancio et al.*'s study, no fibrosis stage inclusion criterion was set. We
26 used a Cox proportional hazards model to investigate the association between diabetes and
27 all-cause mortality, as well as the potential role of other covariates on all-cause mortality,
28 using analyses stratified by diabetes status (i.e., diabetic/not diabetic) at baseline. All
29 covariates were measured at baseline, except for HCV cure, which was considered a time-
30 varying variable.

31
32
33 Of the 10 364 participants included, 13.7% (n=1419) had diabetes at baseline, and
34 82.6% (n=8557) were HCV-cured by the end of follow-up. Median time since HCV diagnosis
35 was between 14 and 15 years. Six hundred and ninety-seven deaths occurred in the study
36 population over 37 633.7 person-years, including 160 diabetic patients. For the whole
37 sample, the all-cause mortality rate was 1.85/100 person-years (95% confidence interval [CI],
38 1.72-1.99), specifically 3.14/100 person-years (95% CI, 2.69-3.67) in diabetic patients, and
39 1.65/100 person-years (95% CI, 1.52-1.80) in non-diabetic patients, corresponding to a
40 hazard ratio (HR) of 1.91 [95% CI, 1.60 – 2.28] for diabetes status ($p < 10^{-3}$).

41
42
43 In multivariable stratified analyses, male gender, older age, being underweight, not
44 HCV cured, and advanced liver fibrosis at baseline (assessed by a FIB-4 score > 3.25 (6))

1
2
3 were all associated with increased mortality in both sub-populations (**Table 1**). Current
4 tobacco smoking, daily coffee intake, an educational level less than high-school diploma, and
5 elevated blood pressure were all independently associated with increased mortality in non-
6 diabetic patients only. The risk of all all-cause mortality associated with having advanced
7 liver fibrosis was two times higher in non-diabetic patients than in diabetic patients (adjusted
8 HR: 5.54 [95% CI, 4.51 – 6.81] *versus* 2.16 [95% CI, 1.51 – 3.09]).
9
10
11
12

13 Our results confirm that diabetes is a significant risk factor for all-cause mortality in
14 chronic HCV patients. While Ciancio *et al.* found that a higher risk of HCC incidence was
15 associated with diabetes in cirrhotic HCV patients, we found that the risk of all-cause
16 mortality associated with having advanced liver fibrosis was two times higher in non-diabetic
17 patients than in diabetic patients. These findings may appear paradoxical.
18
19
20

21 The relatively lower risk of death associated with advanced liver fibrosis in diabetic
22 patients suggests that non-liver-related deaths may also heavily weigh on all-cause mortality
23 in this sub-population, and that the benefits of DAA treatments on glycometabolic control
24 may not be sufficient to counteract the cardiovascular consequences of diabetes as a major
25 cardiovascular risk factor. While a lower hemoglobin A1c level has been associated with
26 lower cardiovascular-related mortality in the general population (7), the fact that more than
27 80% of our study population was over 50, together with the relatively recent arrival of DAA
28 treatments, may mean that the long duration of diabetes in our study sample counteracted
29 the effects which recent improvements in glucose homeostasis have had on cardiovascular
30 events (8). As diabetes is considered a cofactor for liver disease, the lower adjusted hazard
31 ratio for advanced fibrosis in diabetic HCV patients which we found may also be partly due to
32 closer clinical follow-up which focuses on the prevention of liver disease in this sub-
33 population (9). Additionally, this result must be interpreted in the context of the higher all-
34 cause mortality of diabetic patients than their non-diabetic counterparts.
35
36
37
38
39
40
41
42
43

44 Further studies investigating cause-specific mortality in diabetic and non-diabetic
45 HCV patients in the DAA era need to be conducted, in order to shed light on the respective
46 major risk factors for mortality in these two sub-populations, and adapt their clinical follow-up
47 accordingly.
48
49
50
51
52
53
54
55
56
57
58
59
60

References

1. Ciancio A, Ribaldone DG, Dotta A, Giordanino C, Sacco M, Fagoonee S, et al. Long-term follow-up of diabetic and non-diabetic patients with chronic hepatitis C successfully treated with direct-acting antiviral agents. *Liver Int.* 2021;41:276–287.
2. Huang T-S, Lin C-L, Lu M-J, Yeh C-T, Liang K-H, Sun C-C, et al. Diabetes, hepatocellular carcinoma, and mortality in hepatitis C-infected patients: A population-based cohort study. *J. Gastroenterol. Hepatol.* 2017;32:1355–1362.
3. Elkrief L, Chouinard P, Bendersky N, Hajage D, Larroque B, Babany G, et al. Diabetes mellitus is an independent prognostic factor for major liver-related outcomes in patients with cirrhosis and chronic hepatitis C. *Hepatology.* 2014;60:823–831.
4. Butt AA, Yan P, Shuaib A, Abou-Samra A-B, Shaikh OS, Freiberg MS. Direct-Acting Antiviral Therapy for HCV Infection Is Associated With a Reduced Risk of Cardiovascular Disease Events. *Gastroenterology.* 2019;156:987-996.e8.
5. Pol S, Bourliere M, Lucier S, Hezode C, Dorival C, Larrey D, et al. Safety and efficacy of daclatasvir-sofosbuvir in HCV genotype 1-mono-infected patients. *J. Hepatol.* 2017;66:39–47.
6. Sterling Richard K., Lissen Eduardo, Clumeck Nathan, Sola Ricard, Correa Mendes Cassia, Montaner Julio, et al. Development of a simple noninvasive index to predict significant fibrosis in patients with HIV/HCV coinfection. *Hepatology.* 2006;43:1317–1325.
7. Raghavan S, Vassy JL, Ho Y, Song RJ, Gagnon DR, Cho K, et al. Diabetes Mellitus–Related All-Cause and Cardiovascular Mortality in a National Cohort of Adults. *J Am Heart Assoc [Internet].* 2019 [cited 2021 Apr 13];8. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6405678/>
8. Fox CS, Sullivan L, D’Agostino RB, Wilson PWF, Framingham Heart Study. The significant effect of diabetes duration on coronary heart disease mortality: the Framingham Heart Study. *Diabetes Care.* 2004;27:704–708.
9. European Association for the Study of the Liver. Electronic address: easloffice@easloffice.eu. EASL Recommendations on Treatment of Hepatitis C 2016. *J Hepatol.* 2017;66:153–194.

Table 1: Factors associated with all-cause mortality (Cox proportional hazards models), multivariable analyses stratified on diabetes status at baseline: the ANRS CO22 HEPATHER cohort (N = 10 364 patients)

	Whole population (N = 10 364)		Multivariable analyses			
	Non-diabetic patients (N = 8945)	Diabetic patients (N = 1419)	Non-diabetic patients (N = 8129)		Diabetic patients (N = 1309)	
	N (%)	N (%)	aHR [95% CI]	P-value	aHR [95% CI]	P-value
Gender						
Female	4 063 (45.4%)	515 (36.3%)	0.48 [0.39 – 0.59]	< 10 ⁻³	0.59 [0.40 – 0.86]	0.007
Male	4 882 (54.6%)	904 (63.7%)	1		1	
Age (years)						
< 50	2 330 (26.1%)	173 (12.2%)	1		1	
50 – 69	5 515 (61.7%)	946 (66.7%)	2.16 [1.61 – 2.89]	< 10 ⁻³	2.61 [1.04 – 6.58]	0.041
≥ 70	1 100 (12.3%)	300 (21.1%)	3.87 [2.70 – 5.55]	< 10 ⁻³	4.28 [1.58 – 11.57]	0.004
HCV genotype						
1	5 758 (64.4%)	897 (63.2%)				
2/4	1 66 (18.6%)	345 (24.3%)				
3/5/6/7	1 401 (15.7%)	159 (11.2%)				
Missing	120 (1.3%)	18 (1.3%)				
Body mass index (kg.m⁻²)						
Underweight (<18.5)	329 (3.7%)	20 (1.4%)	1.60 [1.03 – 2.49]	0.036	2.80 [1.04 – 7.54]	0.042
Normal (≥ 18.5 and < 25)	4 772 (53.3%)	472 (33.3%)	1		1	
Overweight (≥ 25 and < 30)	2 756 (30.8%)	536 (37.8%)	0.92 [0.75 – 1.13]	0.442	0.78 [0.52 – 1.18]	0.239
Obese (> 30)	1 014 (11.3%)	379 (26.7%)	1.02 [0.76 – 1.38]	0.877	1.00 [0.64 – 1.57]	0.984
Missing data	74 (0.8%)	12 (0.9%)				
Years since HCV diagnosis						
Median [IQR]	14.3 [6.8 – 20.0]	14.77 [8.2 – 20.2]				
Alcohol use^a						
Abstinent with past unhealthy use	1 309 (14.6%)	248 (17.5%)				
Abstinent without past unhealthy use	3 587 (40.1%)	692 (48.8%)				
Moderate use	3 574 (40.0%)	422 (29.7%)				

Unhealthy use	422 (4.7%)	44 (3.1%)				
<i>Missing data</i>	53 (0.6%)	13 (0.9%)				
Tobacco smoking						
Never	3 128 (35.0%)	583 (41.1%)	1		1	
Past	2 342 (26.2%)	444 (31.3%)	0.98 [0.76 – 1.27]	0.899	1.07 [0.71 – 1.62]	0.740
Current	3 460 (38.7%)	392 (27.6%)	1.79 [1.39 – 2.31]	< 10 ⁻³	1.31 [0.84 – 2.06]	0.238
<i>Missing data</i>	15 (0.2%)	0				
HCV transmission category						
IDU	2 488 (27.8%)	244 (17.2%)				
Transfusion	2 588 (28.9%)	467 (32.9%)				
Other	1 554 (17.4%)	277 (19.5%)				
<i>Missing</i>	2 315 (25.9%)	431 (30.4%)				
Educational level						
< high-school diploma	4 687 (52.4%)	859 (60.5%)	1		1	
≥ high-school diploma	4 122 (46.1%)	537 (37.8%)	0.71 [0.59 – 0.86]	0.001	1.01 [0.70 – 1.45]	0.961
<i>Missing data</i>	136 (1.5%)	23 (1.6%)				
HCV treatment status^b						
Not cured	1 571 (17.6%)	236 (16.6%)	1		1	
Cured	7 374 (82.4%)	1 183 (83.4%)	0.26 [0.21 – 0.31]	< 10 ⁻³	0.31 [0.22 – 0.44]	< 10 ⁻³
Advanced liver fibrosis						
No (FIB-4 ≤ 3.25)	6 355 (71.1%)	800 (56.4%)	1		1	
Yes (FIB-4 > 3.25)	2 028 (22.7%)	560 (39.5%)	5.54 [4.51 – 6.81]	< 10 ⁻³	2.16 [1.51 – 3.09]	< 10 ⁻³
<i>Missing data</i>	562 (6.3%)	59 (4.2%)				
Coffee consumption						
0 cup/day	2 524 (28.2%)	419 (29.5%)	1		1	
≥ 1 cup/day	6 266 (70.1%)	965 (68.0%)	0.77 [0.63 – 0.93]	0.007	1.23 [0.84 – 1.81]	0.286
<i>Missing data</i>	155 (1.7%)	35 (2.5%)				
Elevated blood pressure or related treatments						
No	6 674 (74.6%)	615 (43.3%)	1		1	

Yes	2 263 (25.3%)	802 (56.5%)	1.53 [1.25 – 1.87]	< 10 ⁻³	1.36 [0.94 – 1.98]	0.106
<i>Missing data</i>	8 (0.1%)	2 (0.1%)				

Variables with a P-value < 0.20 (Wald test) in the univariable analyses were tested in the multivariable model. The final multivariable model was built using a backward selection of variables). All variables with a P-value < 0.05 in either sub-population was kept in the final model for both sub-populations. Analyses were performed using Stata/SE 14.2 software for Windows (StataCorp LP, College Station, TX).

^a Unhealthy alcohol use was defined as >2 and >3 drinks a day for women and men, respectively.

^b HCV cure was the only time-varying variable, N (%) are given at the end of the follow-up aHR, adjusted hazard ratio; CI, confidence interval; IDU, injecting drug use; IQR, interquartile range

For Peer Review