



Efficacy of Antiviral Treatment in Hepatitis C Virus (HCV)-Driven Monoclonal Gammopathies Including Myeloma

Alba Rodríguez-García, María Linares, María Luz Morales, Sophie Allain-Maillet, Nicolas Mennesson, Ricardo Sanchez, Rafael Alonso, Alejandra Leivas, Alfredo Pérez-Rivilla, Edith Bigot-Corbel, et al.

► To cite this version:

Alba Rodríguez-García, María Linares, María Luz Morales, Sophie Allain-Maillet, Nicolas Mennesson, et al.. Efficacy of Antiviral Treatment in Hepatitis C Virus (HCV)-Driven Monoclonal Gammopathies Including Myeloma. *Frontiers in Immunology*, 2022, 12, pp.797209. 10.3389/fimmu.2021.797209 . inserm-03531984

HAL Id: inserm-03531984

<https://inserm.hal.science/inserm-03531984>

Submitted on 18 Jan 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Efficacy of Antiviral Treatment in Hepatitis C Virus (HCV)-Driven Monoclonal Gammopathies Including Myeloma

OPEN ACCESS

Edited by:

Keqiang Chen,
National Cancer Institute at Frederick,
United States

Reviewed by:

Helga Ogmundsdottir,
University of Iceland, Iceland
Shuo Geng,
Virginia Tech, United States

*Correspondence:

María Linares
mlinares@ucm.es

[†]These authors have contributed
equally to this work

[‡]These authors have contributed
equally to this work

Specialty section:

This article was submitted to
Cancer Immunity
and Immunotherapy,
a section of the journal
Frontiers in Immunology

Received: 18 October 2021

Accepted: 14 December 2021

Published: 11 January 2022

Citation:

Rodríguez-García A, Linares M,
Morales ML, Allain-Maillet S,
Mennesson N, Sanchez R, Alonso R,
Leivas A, Pérez-Rivilla A,
Bigot-Corbel E, Hermouet S and
Martínez-López J (2022) Efficacy of
Antiviral Treatment in Hepatitis C
Virus (HCV)-Driven Monoclonal
Gammopathies Including Myeloma.
Front. Immunol. 12:797209.
doi: 10.3389/fimmu.2021.797209

**Alba Rodríguez-García^{1†}, María Linares^{1,2*†}, María Luz Morales¹, Sophie Allain-Maillet³,
Nicolas Mennesson³, Ricardo Sanchez¹, Rafael Alonso¹, Alejandra Leivas¹,
Alfredo Pérez-Rivilla⁴, Edith Bigot-Corbel^{3,5}, Sylvie Hermouet^{3,6†}
and Joaquín Martínez-López^{1,7‡}**

¹ Department of Translational Hematology, Instituto de Investigación Hospital 12 de Octubre (imas12), Hematological Malignancies Clinical Research Unit H120-CNIO, CIBERONC, Madrid, Spain, ² Department of Biochemistry and Molecular Biology, Pharmacy School, Universidad Complutense de Madrid, Madrid, Spain, ³ Université de Nantes, Institut National de la Santé et de la Recherche Médicale (Inserm), Centre de Recherche en Cancérologie et Immunologie Nantes-Angers (CRCINA), Nantes, France, ⁴ Department of Microbiology, Hospital Universitario 12 de Octubre, Madrid, Spain, ⁵ Laboratoire de Biochimie, Centre Hospitalier Universitaire (CHU) de Nantes, Nantes, France, ⁶ Laboratoire d'Hématologie, Centre Hospitalier Universitaire (CHU) de Nantes, Nantes, France, ⁷ Department of Medicine, Medicine School, Universidad Complutense de Madrid, Madrid, Spain

Multiple myeloma (MM) remains an incurable plasma cell malignancy. While its origin is enigmatic, an association with infectious pathogens including hepatitis C virus (HCV) has been suggested. Here we report nine patients with monoclonal gammopathy of undetermined significance (MGUS) or MM with previous HCV infection, six of whom received antiviral treatment. We studied the evolution of the gammopathy disease, according to anti-HCV treatment and antigen specificity of purified monoclonal immunoglobulin, determined using the INNO-LIA™ HCV Score assay, dot-blot assays, and a multiplex infectious antigen microarray. The monoclonal immunoglobulin from 6/9 patients reacted against HCV. Four of these patients received antiviral treatment and had a better evolution than untreated patients. Following antiviral treatment, one patient with MM in third relapse achieved complete remission with minimal residual disease negativity. For two patients who did not receive antiviral treatment, disease progressed. For the two patients whose monoclonal immunoglobulin did not react against HCV, antiviral treatment was not effective for MGUS or MM disease. Our results suggest a causal relationship between HCV infection and MGUS and MM progression. When HCV was eliminated, chronic antigen-stimulation disappeared, allowing control of clonal plasma cells. This opens new possibilities of treatment for MGUS and myeloma.

Keywords: infection, hepatitis (C) virus, antiviral, multiple myeloma, monoclonal gammopathies

INTRODUCTION

Multiple myeloma (MM) is a common hematologic malignancy (1.2% of all tumors) characterized by the clonal expansion and transformed plasma cells in the bone marrow. MM is always preceded by monoclonal gammopathy of undetermined significance (MGUS), an asymptomatic stage that does not always evolve to MM (1–3). Despite great advances in the understanding and treatment of MM, its origin is unknown, and it remains an incurable disease.

The primary function of plasma cells is to produce and secrete large amounts of immunoglobulins (Ig) that mediate humoral immunity against infection. Healthy plasma cells differentiate from immature B cells when they recognize an antigen foreign to the organism. This process occurs in the germinal centers of the secondary lymphoid organs, where B cells proliferate and select somatic hypermutations that have high affinity with the external antigen. In MM, monoclonal plasma cells secrete large quantities of a single Ig, monoclonal Ig, which serves as a marker of the disease and triggers much of the symptomatology (4).

Latent infection and chronic antigen stimulation are now recognized as initial pathogenic events leading to cancer. This association has been shown in several hematologic malignancies, such as chronic lymphocytic leukemia (CLL) and different types of lymphoma (5, 6). B-cell receptor (BCR) signaling is central for the specific recognition of Igs, suggesting that specific antigens could be involved in the development of different types of CLL. Interestingly, Hooijboom and colleagues recently described a new subset of CLL that expresses stereotypic BCRs specific for β -(1, 6)-glucan, a major component of yeasts and fungi of the microbiota (7). The stimulation of BCR directed from these antigens seems to trigger signaling pathways through different mediators such as p53 and c-Myc, which result in proliferation, suppressed apoptosis, survival and alterations of cell migration (8).

In support of chronic antigenic stimulation as a pathogenic mechanism in MGUS and MM, several studies suggest an association between MM and viral infection, particularly hepatitis C virus (HCV), human immunodeficiency virus or Epstein Barr virus (EBV) (9–14). In addition, Nair et al. identified glucosylsphingosine (GlcSph) as the target of monoclonal Igs both in the context of Gaucher's disease and in sporadic gammopathies (15, 16). Antigen-mediated stimulation led to an increase in the amount of monoclonal Ig and plasma cells in a murine model, confirming the role of chronic antigenic stimulation in the pathogenesis of MM. Independently, we recently reported that one-quarter of all MM cases might be initiated by infectious pathogens, including EBV and HCV (17, 18). In this line, a recent meta-analysis demonstrated a higher risk (2.67-fold) of developing MM in HCV-infected patients than in controls (11). These findings point to a role for HCV in the pathogenic development of MGUS and MM.

This concept opens new possibilities for treatment of MGUS and MM: target antigen reduction. If the target of the monoclonal Ig is eliminated, chronic antigen-stimulation disappears, leading to the control of clonal plasma cells. The

efficacy of this therapeutic approach has been proven for GlcSph-associated MGUS and SMM (19).

In the present study, we explored the efficacy of anti-HCV treatment in a series of MGUS and MM patients linked to HCV. We report on a series of nine MGUS and MM patients with HCV infection, for whom the reactivity of the monoclonal Ig against HCV proteins was analyzed. We demonstrate for the first time that in cases where the monoclonal Ig reacted against HCV, treating the HCV infection improved MGUS and MM disease. Importantly, in a patient with refractory MM whose monoclonal IgG reacted specifically to HCV core protein, treatment of the HCV infection resulted in complete remission (CR) of MM, and the patient has been in clinically stable remission for four years.

MATERIALS AND METHODS

Patients

Nine patients who developed MGUS (n=5) or MM (n=4) after HCV infection was detected in these patients were identified and classified into two groups: those who received antiviral treatment, and those who did not. The characteristics of patients at diagnosis are summarized in **Table 1**. Relevant information about the dates of diagnosis and treatments of the HCV infection and of the gammopathy are summarized in **Supplementary Tables S1, S2**. The study was approved by our Institutional Review Board and the patients provided written informed consent in accordance with the Declaration of Helsinki.

Immunofixation of Igs

Disease status was monitored by the quantification of monoclonal Ig or total Igs or free kappa/lambda light chain levels in serum, depending on the patient. Protein levels were routinely visualized by serum protein electrophoresis and/or immunofixation electrophoresis (20).

Determination of Viral Load

Quantitative determination of RNA from HCV in human plasma containing K2EDTA was performed using the VERIS MDx system (Beckman Coulter). The RNA-HCV assay has been validated to provide quantitative results of samples containing HCV genotypes 1–6 (21). The main characteristics of the HCV infection in patients treated with antivirals are summarized in **Supplementary Tables S1, S2**.

Purification of Monoclonal IgG and IgA

Agarose gel electrophoresis and purification of patients' monoclonal Ig from other Igs present in serum samples was performed as described (17, 18, 22, 23) (**Figure S1** in the **Supplementary Appendix**). Protein concentrations were determined on a Nanodrop ND-1000 spectrophotometer. As an exception, in one patient diagnosed with Bence-Jones MM, kappa light chains were purified using PureProteome™ Protein G and Kappa magnetic beads (Merck Millipore) and purity was

TABLE 1 | Main patient characteristics at diagnosis of the gammopathy disease.

| Patients | P1 | P2 | P3 | P4 | P5 | P6 | P7 | P8 | P9 |
|---------------------------------|------------------|-------------------|-------------------|--------------------|-------------------------|--------------------|-------------------|------------------|-----------------|
| Sex (M/F) | F | F | F | M | M | M | F | F | F |
| Age (years) | 66 | 67 | 53 | 78 | 56 | 53 | 76 | 80 | 79 |
| Diagnosis | MM IgG λ | MGUS IgG κ | MGUS IgG κ | MGUS IgA λ | MM Bence-Jones κ | MGUS IgA λ | MGUS IgG κ | MM IgA λ | MM IgG κ |
| Date of diagnosis | August 2011 | January 2003 | November 2017 | June 2016 | October 2014 | September 2018 | November 2015 | January 2006 | September 2016 |
| Monoclonal Ig (g/dL) | 5.67 | 1.72 | 0.74 | 1.47 | 0.0914 | 1.23 | 1.17 | 4.37 | 1.57 |
| ISS | IIA | NA | NA | NA | III | NA | NA | IIA | III |
| Hemoglobin (g/dL) | 12.8 | 15 | 15.1 | 12.6 | 10.3 | 15.9 | 10.5 | 11.6 | 10 |
| Creatinine (mg/dL) | 0.72 | 0.73 | 0.63 | 1.03 | 7.99 | 0.92 | 1.07 | 0.96 | 1.99 |
| Bone Marrow Plasma Cells (%) | 16 | – | – | 9 | 98 | – | – | 36 | 45 |
| Platelets (10 ⁹ /L) | 163 | 170 | 309 | 173 | 190 | 104 | 206 | 152 | 231 |
| Leukocytes (10 ⁹ /L) | 3.4 | 7.1 | 5 | 4.5 | 4.8 | 5.6 | 6.3 | 3.4 | 6.8 |
| Calcemia (mmol/L) | 2.4 | 2.5 | 2.3 | 2.4 | 2.3 | 2.3 | 2.4 | 2.5 | 2.3 |
| β_2 -Microglobulin (mg/L) | 2.9 | – | 2.1 | – | 13.07 | – | – | 6.44 | 5.9 |
| High risk cytogenetics | No | – | – | – | No | No | Yes ^a | No | No |
| LDH high | No | No | Yes | No | Yes | No | No | No | Yes |

M, male; F, female; Ig, immunoglobulin; MM, multiple myeloma; MGUS, monoclonal gammopathy of undetermined significance; ISS, Multiple Myeloma International Staging System; LDH, lactate dehydrogenase.

^aHigh risk cytogenetics: del17p, t (4;14), t (14;16) or t (14;20).

NA, Not applicable.

evaluated by conventional native 15% polyacrylamide gel electrophoresis.

Analysis of the Specificity of Antigenic Recognition of Purified Monoclonal Igs

The INNO-LIATM HCV Score (Fujirebio) was used to analyze the reactivity of patient monoclonal IgG to HCV proteins. For monoclonal IgA, dot blotting assays with HCV proteins were performed on nitrocellulose membranes (Amersham) spotted with 1 μ g of recombinant HCV core, NS3 and NS4 proteins (Abcam, Advanced Biotechnologies Inc.) (3 spots), which were then incubated with the patient's serum or with the purified monoclonal IgA. The chemiluminescent microparticle immunoassay, Alinity i Anti-HCV (Abbott GmbH & Co. KG), on the Alinity i System was used for the qualitative detection of HCV in the patient diagnosed with Bence-Jones MM.

The multiplex infectious antigen microarray (MIAA) assay was used to analyze the reactivity of serum Igs and of purified monoclonal IgG or IgA against commercially available antigens and/or lysates from EBV, cytomegalovirus (CMV), herpes simplex virus-1 (HSV-1), herpes simplex virus-2 (HSV-2), varicella-zoster virus (VZV), *Helicobacter pylori* (*H. pylori*), *Toxoplasma gondii*, and *Borrelia burgdorferi*, as described (14, 17).

RESULTS

Absence of Disease Progression in Patients With HCV-Specific Monoclonal Ig Who Received HCV Antiviral Treatment

Six patients had a monoclonal Ig that specifically recognized the HCV virus: patients P1–4, P7 and P8. Four of the patients (P1–4) received HCV antiviral treatment (detailed in **Table S1**).

As expected, after antiviral treatment, HCV loads decreased, to undetectable levels for patients P1, P3 and P4 (**Supplementary Table S2**). For patient P1, who suffered from MM in third relapse at the time of anti-HCV treatment, eradication of HCV (**Figure 1A**) was associated with complete remission (CR) of MM in the absence of new anti-MM therapy. The patient's monoclonal IgG drastically decreased (**Figure 1B**) and bone marrow aspirates showed < 5% of plasma cells by cytology and minimal residual disease negativity (**Figure 1C**). The number of plasma cell clones present in the sample and the tumor load was analyzed by next-generation sequencing, as previously reported (20, 24, 25). The monoclonal IgG present in pre-HCV treatment samples disappeared after anti-HCV treatment (**Figure 1D**). Forty-five months later, the patient remains in CR of MM with minimal residual disease negativity as assessed by next generation flow cytometry and undetectable monoclonal IgG. Patient P1's purified monoclonal Ig specifically targeted the core protein of HCV (INNO-LIATM HCV Score assay, **Figure 1E**).

When HCV-positive patients P2–4 (all MGUS patients) received antiviral treatment (and no other treatment), the HCV viral load decreased to minimum level in all cases (**Figure 2A** and **Supplementary Table S2**). Antiviral treatment was followed by a clear and stable decrease in the amount of monoclonal Ig decreased for patients P2 and P4. For patient P3, the concentration of monoclonal Ig decreased slightly and has now been stable for 39 months (**Figure 2B**). The purified monoclonal Ig from the aforementioned 3 MGUS patients (P2–4) specifically targeted antigens of HCV, either C1 (core) or NS3/NS4 proteins (**Figures 2C, D**). It should be noted that when dealing with MGUS, following the recommendations of the clinical guidelines, these patients did not receive any hematological treatment (**Supplementary Table S1**).

To confirm that the monoclonal Igs of patients P1–4 specifically recognized only HCV, we tested their reactivity against other




FIGURE 1 | Stable complete remission and specific recognition of HCV in a patient with MM treated with antiviral drugs. Panel **(A)** shows the HCV load quantified by reverse-transcriptase quantitative polymerase chain reaction in the patient's serum. The black horizontal dotted line represents the threshold of virus detection. Panel **(B)** shows the quantity of monoclonal IgG as determined by serum protein electrophoresis. Panel **(C)** shows the percentage of plasma cells determined by bone marrow examination (cytology) or next generation 8-color multiparametric flow cytometry, and the percentage of pathological plasma cells as determined by immunophenotyping. **(A–D)** The blue vertical dotted line indicates the time of antiviral treatment. Panel **(D)** shows a representative agarose gel electrophoresis of polyclonal IgG in serum before (2014) and after (2017) antiviral treatment – the encircled band corresponds to the patient's monoclonal Ig. Panel **(E)** shows the INNO-LIA™ HCV test and immunoblotting assay used to detect reactivity of the patient's serum IgG and of the purified monoclonal IgG against different HCV proteins. The signal obtained with HCV core for the monoclonal IgG of patient P1 was weak but always reproducible when different preparations of the purified monoclonal IgG were tested. C, positive controls; S, serum; Mc, monoclonal Ig.

microorganisms using the MIAA assay. Serological status (polyclonal Ig + monoclonal Ig) was evaluated in parallel. As expected, serum samples were reactive against several pathogens (EBV, CMV, HSV-1, HSV-2, VZV, *H. pylori*, *B. burgdorferi*, etc.). Confirming their purity and their specificity for HCV only, the monoclonal Ig preparations failed to react against any of these pathogens (Figure S2 in the Supplementary Appendix).

Disease Evolution in Patients for Whom the Monoclonal Ig's Target Was Not Treated

Two patients with Bence-Jones (light chain) MM (P5) or MGUS (P6) were also successfully treated with antivirals (Figure 2A and Supplementary Table S2). However, their monoclonal Ig did not recognize HCV (Figures 2C, D) and these patients did not show an improvement of their MM (P5) or MGUS (P6) disease (Figure 2B). Patient P5 (light chain MM) presented two relapses after HCV therapy, despite different MM treatments (Supplementary Table S1). For patient (P6), light chain MGUS persisted with 7.5% plasma cells and a monoclonal component of 0.86g/dl 9 months after HCV therapy.

Three HCV-infected patients (P7-9) did not receive antiviral treatment, and disease evolution was unfavorable for all three patients. As shown in Table 2, within five months, MGUS patient P7 progressed to smoldering multiple myeloma (SMM) (17% of plasma cells). Despite various MM treatments (Supplementary Table S1), disease also progressed for patients P8 and P9, and both patients died. For patients P7 and P8, their purified monoclonal Ig specifically targeted the HCV NS3 protein (Figure 3), but for patient P9, the monoclonal Ig was shown to target a different virus, HSV-1.

DISCUSSION

Previous reports have suggested an association between MGUS, MM and infection. In particular, we and others showed that at least one-third of MGUS and MM cases present with a monoclonal Ig that targets an infectious pathogen, including HCV (17, 22, 23, 26–28). The recent *in vivo* demonstration that chronic antigen stimulation can lead to emergence of clonal plasma cells supports the role of chronic infection in the




FIGURE 2 | Disease evolution and evaluation of HCV-specificity of the monoclonal Ig from patients who were treated with antiviral drugs. Panel **(A)** shows the HCV load quantified by reverse-transcriptase quantitative polymerase chain reaction in patients' serum. The black horizontal dotted line represents the threshold of virus detection. Panel **(B)** shows the quantity of monoclonal Ig as determined by serum protein electrophoresis. **(A, B)** The blue vertical dotted line indicates the time of antiviral treatment. Panel **(C)** shows the INNO-LIATM HCV test to detect reactivity of patients' serum IgGs and of the purified monoclonal IgG against different HCV proteins. Panel **(D)** shows the dot-blot assay to detect reactivity of patients' serum IgAs and of the purified monoclonal IgA against different HCV proteins (IgAs cannot be studied using the INNO-LIATM HCV test). The monoclonal IgA of patient P4 strongly recognized the HCV NS3 protein, whereas the monoclonal IgA of patient P6 did not recognize any HCV protein of the assay. C, positive controls; S, serum; Mc, monoclonal Ig.

development of subsets of MGUS and MM (15). This new pathogenic model is important since it offers, for the first time, the possibility of treatment for patients with MGUS, as well as novel therapeutic approach for MM: reduction or suppression of


antigenic stimulation by treating the infection that initiated the gammopathy.

In support of this approach, the present study shows that whenever HCV-infected patients later diagnosed with MGUS or

TABLE 2 | Classification of patients, response to anti-HCV treatment and MGUS or MM evolution.

| Patients | Diagnosis | Monoclonal Ig | Purified | Target of the monoclonal Ig | Antiviral treatment | Decrease in monoclonal Ig | Disease progression | Length, progression-free (months) |
|----------|----------------|----------------------|----------|-----------------------------|---------------------|---------------------------|---------------------|-----------------------------------|
| P1 | MM | IgG λ | Yes | HCV core C1 | Yes | Yes | No, stable CR | 45 |
| P2 | MGUS | IgG κ | Yes | HCV NS3 | Yes | Yes | No | 46 |
| P3 | MGUS | IgG κ | Yes | HCV NS3 | Yes | Yes | No | 26 |
| P4 | MGUS | IgA λ | Yes | HCV NS3 | Yes | Yes | No | 39 |
| P5 | MM Bence-Jones | Light chain κ | Yes | Unknown | Yes | No | Yes, 2 relapses | 7 |
| P6 | MGUS | IgA λ | Yes | Unknown | Yes | No | No, stable MGUS | 17 |
| P7 | MGUS | IgG κ | Yes | HCV NS3 | No | No | Yes, to SMM | – |
| P8 | MM | IgA λ | Yes | HCV NS3 | No | No | Yes, death | – |
| P9 | MM | IgG κ | Yes | HSV-1 | No | No | Yes, death | – |

MM, multiple myeloma; HCV, hepatitis C virus; CR, complete remission; MGUS, monoclonal gammopathy of undetermined significance; SMM, smoldering multiple myeloma; HSV-1, herpes simplex virus-1.



MM presented with a monoclonal Ig that targeted HCV, antiviral treatment markedly improved the outcome of their gammopathy. Importantly, we report for the first time that a long and stable CR has been achieved in a patient with refractory MM whose monoclonal IgG targeted HCV, after the sole administration of the antiviral treatment (with sofosbuvir and ledipasvir), and long after the administration of other hematological treatments. After 48 months, the patient remained in CR without MM symptoms. CR of MM is presumably linked to HCV disappearance since the antiviral

drugs received by this patient have never been shown to have any anti-MM effect. For the other five HCV-positive MGUS cases, clonal HCV-specificity was also demonstrated through the recognition of HCV by the purified monoclonal IgG, and lack of recognition of other infectious pathogens, as assessed by the MIAA assay. Logically, for the two HCV-positive patients whose monoclonal Ig did not react against HCV, antiviral treatment had no favorable effect on the patient's MGUS or MM disease (hereby confirming that anti-HCV drugs do not have intrinsic anti-MM effects).

In this study, we selected patients who had had HCV infection prior to the diagnosis of MGUS or MM. We were able to identify the target of the monoclonal Ig for 7/7 patients with a complete monoclonal Ig (heavy + light chains); the targets were HCV in 6 cases, HSV-1 in 1 case. These results are in line with our previous report that when MGUS or MM patients have a history of HCV infection, their monoclonal Ig target HCV in ~85% cases (9). Unfortunately, present assays do not allow to identify the target of clonal light chains, as observed for the two patients with light chain MGUS (P6) or MM (P5).

Thus, together with our previous studies, the present work demonstrates that it is feasible to identify the target of the monoclonal Ig of patients, then link the gammopathy to a previous chronic infection. This new approach should be useful for a significant fraction of MGUS and MM patients: for instance, a recent study revealed that until 7.4% of MM patients were positive for HCV RNA by RT-PCR assay. In this study, the authors emphasize that serologic tests at the time of diagnosis of MM are necessary to identify infected patients, and they propose that confirmation of positive cases by molecular techniques should be mandatory (29). Whether the gammopathy is linked to the infectious agent can then be addressed by showing that the patient's monoclonal Ig reacts against the pathogen. This information allows to propose antiviral or antibiotic treatments to MGUS patients (presently not treated), with the aim of curing the MGUS, and also to SMM and MM patients, as adjuvant treatments, with the aim to improve the patient's response to chemotherapy. This "target antigen reduction" approach should improve the prognosis in patients with gammopathies, especially those at the MGUS or SMM stages. In fact, previous studies reported the need for a complete follow-up of patients with chronic infection due to the possibility that they develop MM or non-Hodgkin lymphoma (NHL) (30). Panfilio et al. (31) previously reported a MM regression after antiviral treatment of one patient with HCV infection; unfortunately, the antigen targeted by the patient's monoclonal Ig was not investigated by these authors (31). Similarly, the beneficial effect of interferon treatment against MM in HCV-infected patients was reported recently by Ioannou et al. (32). Regrettably, the antigen specificity of the monoclonal Igs of patients was not studied by these authors (32). In contrast, our study establishes that anti-HCV treatment improves and even suppresses MGUS or MM disease when the gammopathy is driven by HCV, i.e. when the patient's monoclonal Ig targets HCV. Inversely, when the monoclonal Ig did not target HCV, anti-HCV treatment had no effect on the gammopathy, an indirect confirmation that the beneficial effect of antiviral therapy on MGUS and MM acts *via* the clearance of the HCV infection. Similar observations have been made for HIV-infected patients diagnosed with MM, for whom HIV treatment resulted in a significant reduction in serum monoclonal Ig, a superior response to MM therapy and improved overall and progression-free survival (33). Although the specificity of the monoclonal Ig of these patients was not studied, these observations were consistent with a beneficial role of antiretroviral therapy in terms of control of the plasmacytic clone and Ig production (33).

Despite the fact that HCV has long been shown to be a risk factor for the development of certain proliferative diseases, including MM, the underlying mechanisms remained unclear. Although not completely understood, the mechanisms of HCV cell entry involve the binding of the E2 protein of the virus envelope and CD81, a molecule which is very abundant on the surface of hepatocytes and also of B lymphocytes (34–36). Hence, HCV can infect both hepatocytes and B-cells, and thus directly induce genetic alterations in infected cells and the subsequent development of hepatic or/and B-cell malignancies. However, in the context of MGUS and MM, the most frequent mechanism of cell transformation by HCV is likely to be indirect, since HCV antigens are detected in peripheral mononuclear cells in chronically infected patients (37). Studies based on the importance of sustained stimulation over time by HCV antigens suggest a mechanism of action analogous to the one at play during infection by *H. pylori*, which increases the risk of indirect carcinogenesis and lymphomagenesis in patients infected with a virulent strain. *H. pylori* infection induces gastric inflammation and chronic antigen stimulation of B-cell immune responses, hereby facilitating the acquisition of genetic alterations and transformation of infected gastric tissues as well as cells of the B lineage. In infected patients, it is not rare to observe the presence of oligoclonal Igs then of a monoclonal Ig (thus of a plasmacytic clone), which normally disappear rapidly. The same process is likely at play in infection-initiated MGUS, except that the plasmacytic clone and monoclonal Ig persist for more than 6 months, often for several years, eventually progressing toward MM. Our data show that in the context of HCV infection, such clones remain antigen-dependent, since successful antiviral treatment results in the reduction or suppression of the plasmacytic clone and monoclonal Ig.

In summary, our study highlights the urgency of treating HCV in infected patients, especially in MGUS, SMM or MM cases who present a monoclonal Ig that reacts against the virus, prior to chemotherapy schemes. Importantly, antiviral treatment of all HCV-positive patients should prevent the development of HCV-driven MGUS and MM. Overall, our findings suggest that chronic stimulation by HCV may promote the development of MGUS and MM in chronically infected patients. This observation has evident clinical consequences, since the identification of a patient with a monoclonal Ig specific for a treatable pathogen, such as HCV, would possibly allow curative antiviral treatment in case of MGUS, and improved response to chemotherapy schemes in case of MM.

DATA AVAILABILITY STATEMENT

The original contributions presented in the study are included in the article/**Supplementary Material**. Further inquiries can be directed to the corresponding author.

ETHICS STATEMENT

The studies involving human participants were reviewed and approved by Comité de Ética del Instituto de Investigación

Hospital 12 de Octubre. The patients/participants provided their written informed consent to participate in this study.

AUTHOR CONTRIBUTIONS

AR-G and ML performed experiments, analyzed data, and wrote the initial manuscript draft. MLM contributed to analyzed and discuss the data. SA-M and NM performed experiments and analyzed data. RS and EB-C contributed to experiments and analyzed the data. RA, AL and AP-R contributed with patient samples and data. SH and JM-L designed the research, provided resources and financial support and edited the manuscript. All authors contributed to the article and approved the submitted version.

FUNDING

This work was supported by grants to SH from the Ligue Nationale contre le Cancer (Comités Départementaux 44, 56,

29, 85, 35) and International Myeloma Foundation (IMF) (Brian D. Novis senior grant). We acknowledge the Instituto de Investigación Sanitaria Hospital 12 de Octubre (imas12), CIBERONC, AECC (Accelerator Award and Ideas Semilla), and the CRIS foundation for their help. MLM has a grant from the Spanish Society of Hematology.

ACKNOWLEDGMENTS

We are particularly indebted to all the patients who participated in the study.

SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fimmu.2021.797209/full#supplementary-material>

REFERENCES

- Palumbo A, Anderson K. Multiple Myeloma. *N Engl J Med* (2011) 364:1046–60. doi: 10.1056/NEJMra1011442
- Rajkumar SV, Dimopoulos MA, Palumbo A, Blade J, Merlini G, Mateos M-V, et al. International Myeloma Working Group Updated Criteria for the Diagnosis of Multiple Myeloma. *Lancet Oncol* (2014) 15:e538–48. doi: 10.1016/S1470-2045(14)70442-5
- Ferlay J, Colombet M, Soerjomataram I, Dyba T, Randi G, Bettio M, et al. Cancer Incidence and Mortality Patterns in Europe: Estimates for 40 Countries and 25 Major Cancers in 2018. *Eur J Cancer Oxf Engl* (2018) 1990:103:356–387. doi: 10.1016/j.ejca.2018.07.005
- Guillerey C, Nakamura K, Vuckovic S, Hill GR, Smyth MJ. Immune Responses in Multiple Myeloma: Role of the Natural Immune Surveillance and Potential of Immunotherapies. *Cell Mol Life Sci* (2016) 73:1569–89. doi: 10.1007/s00018-016-2135-z
- Morrison VA. Infections in Patients With Leukemia and Lymphoma. In: V Stosor, TR Zembower, editors. *Infectious Complications in Cancer Patients*. Cham: Springer International Publishing. (2014) p. 319–49.
- Seifert M, Scholtysik R, Küppers R. Origin and Pathogenesis of B Cell Lymphomas. In: R Küppers, editor. *Lymphoma*. New York, NY: Springer New York. (2019) p. 1–33.
- Hoogeboom R, van Kessel KPM, Hochstenbach F, Wormhoudt TA, Reinten RJA, Wagner K, et al. A Mutated B Cell Chronic Lymphocytic Leukemia Subset That Recognizes and Responds to Fungi. *J Exp Med* (2013) 210:59–70. doi: 10.1084/jem.20121801
- Stevens WBC, Netea MG, Kater AP, van der Velden WJFM. Trained Immunity: Consequences for Lymphoid Malignancies. *Haematologica* (2016) 101:1460–8. doi: 10.3324/haematol.2016.149252
- Bigot-Corbel E, Gassin M, Corre I, Le Carrer D, Delaroché O, Hermouet S. Hepatitis C Virus (HCV) Infection, Monoclonal Immunoglobulin Specific for HCV Core Protein, and Plasma-Cell Malignancy. *Blood* (2008) 112:4357–8. doi: 10.1182/blood-2008-07-167569
- McShane CM, Murray LJ, Engels EA, Landgren O, Anderson LA. Common Community-Acquired Infections and Subsequent Risk of Multiple Myeloma: A Population-Based Study: Infections and Multiple Myeloma. *Int J Cancer* (2014) 134:1734–40. doi: 10.1002/ijc.28479
- Li Y, Li Y, Zhang L, Li W. Hepatitis C Virus Infection and Risk of Multiple Myeloma: Evidence From a Meta-Analysis Based on 17 Case-Control Studies. *J Viral Hepat* (2017) 24:1151–9. doi: 10.1111/jvh.12742
- Yan J, Wang J, Zhang W, Chen M, Chen J, Liu W. Solitary Plasmacytoma Associated With Epstein-Barr Virus: A Clinicopathologic, Cytogenetic Study and Literature Review. *Ann Diagn Pathol* (2017) 27:1–6. doi: 10.1016/j.anndiagpath.2016.09.002
- Hermouet S, Corre I, Gassin M, Bigot-Corbel E, Sutton CA, Casey JW. Hepatitis C Virus, Human Herpesvirus 8, and the Development of Plasma-Cell Leukemia. *N Engl J Med* (2003) 348:178–9. doi: 10.1056/NEJM200301093480219
- Feron D, Charlier C, Gourain V, Garderet L, Coste-Burel M, Le Pape P, et al. Multiplexed Infectious Protein Microarray Immunoassay Suitable for the Study of the Specificity of Monoclonal Immunoglobulins. *Anal Biochem* (2013) 433:202–9. doi: 10.1016/j.ab.2012.10.012
- Nair S, Sng J, Boddupalli CS, Seckinger A, Chesi M, Fulciniti M, et al. Antigen-Mediated Regulation in Monoclonal Gammopathies and Myeloma. *JCI Insight* (2018) 3(8):e98259. doi: 10.1172/jci.insight.98259
- Nair S, Branagan AR, Liu J, Boddupalli CS, Mistry PK, Dhodapkar MV. Clonal Immunoglobulin Against Lysolipids in the Origin of Myeloma. *N Engl J Med* (2016) 374:555–61. doi: 10.1056/NEJMoa1508808
- Bosseboeuf A, Feron D, Tallet A, Rossi C, Charlier C, Garderet L, et al. Monoclonal IgG in MGUS and Multiple Myeloma Targets Infectious Pathogens. *JCI Insight* (2017) 2(19):e95367. doi: 10.1172/jci.insight.95367
- Bosseboeuf A, Allain-Maillet S, Mennesson N, Tallet A, Rossi C, Garderet L, et al. Pro-Inflammatory State in Monoclonal Gammopathy of Undetermined Significance and in Multiple Myeloma Is Characterized by Low Sialylation of Pathogen-Specific and Other Monoclonal Immunoglobulins. *Front Immunol* (2017) 8:1347. doi: 10.3389/fimmu.2017.01347
- Nair S, Bar N, Xu ML, Dhodapkar M, Mistry PK. Glucosylsphingosine But Not Saposin C, is the Target Antigen in Gaucher Disease-Associated Gammopathy. *Mol Genet Metab* (2020) 129:286–91. doi: 10.1016/j.ymgme.2020.01.009
- Kumar S, Paiva B, Anderson KC, Durie B, Landgren O, Moreau P, et al. International Myeloma Working Group Consensus Criteria for Response and Minimal Residual Disease Assessment in Multiple Myeloma. *Lancet Oncol* (2016) 17:e328–46. doi: 10.1016/S1470-2045(16)30206-6
- Braun P, Delgado R, Drago M, Fanti D, Fleury H, Gismondo MR, et al. European Multicenter Study on Analytical Performance of DxN Veris System HCV Assay. *J Clin Microbiol* (2017) 55:1186–92. doi: 10.1128/JCM.02163-16
- Bosseboeuf A, Seillier C, Mennesson N, Allain-Maillet S, Fourny M, Tallet A, et al. Analysis of the Targets and Glycosylation of Monoclonal IgAs From MGUS and Myeloma Patients. *Front Immunol* (2020) 11:854. doi: 10.3389/fimmu.2020.00854
- Bosseboeuf A, Mennesson N, Allain-Maillet S, Tallet A, Piver E, Decaux O, et al. Characteristics of MGUS and Multiple Myeloma According to the Target of Monoclonal Immunoglobulins, Glucosylsphingosine, or Epstein-Barr Virus EBNA-1. *Cancers* (2020) 12:1254. doi: 10.3390/cancers12051254

24. Martinez-Lopez J, Sanchez-Vega B, Barrio S, Cuenca I, Ruiz-Heredia Y, Alonso R, et al. Analytical and Clinical Validation of a Novel in-House Deep-Sequencing Method for Minimal Residual Disease Monitoring in a Phase II Trial for Multiple Myeloma. *Leukemia* (2017) 31:1446–9. doi: 10.1038/leu.2017.58
25. Flores-Montero J, Sanoja-Flores L, Paiva B, Puig N, García-Sánchez O, Böttcher S, et al. Next Generation Flow for Highly Sensitive and Standardized Detection of Minimal Residual Disease in Multiple Myeloma. *Leukemia* (2017) 31:2094–103. doi: 10.1038/leu.2017.29
26. Babel N, Schwarzmann F, Pruss A, Volk H-D, Reinke P. Monoclonal Gammopathy of Undetermined Significance (MGUS) Is Associated With an Increased Frequency of Epstein-Barr Virus (EBV) Latently Infected B Lymphocytes in Long-Term Renal Transplant Patients. *Transplant Proc* (2004) 36:2679–82. doi: 10.1016/j.transproceed.2004.09.046
27. Mameli G, Fozza C, Niegowska M, Corda G, Ruda MF, Barraqueddu F, et al. Epstein-Barr Virus Infection Is Associated to Patients With Multiple Myeloma and Monoclonal Gammopathy of Undetermined Significance. *Leuk Lymphoma* (2017) 58:466–9. doi: 10.1080/10428194.2016.1190976
28. Harb J, Mennesson N, Lepetit C, Fourny M, Louvois M, Bosseboeuf A, et al. Comparison of Monoclonal Gammopathies Linked to Poliovirus or Coxsackievirus vs. Other Infectious Pathogens. *Cells* (2021) 10(2):438. doi: 10.3390/cells10020438
29. Kassem NM, Kassem HA, Ibrahim M, Zawam H, Hamada E. The Clinical Impact of Hepatitis C Virus Infection in Egyptian Multiple Myeloma Patients. *J Egypt Natl Cancer Inst* (2020) 32:43. doi: 10.1186/s43046-020-00054-0
30. Lakatos PL. Development of Multiple Myeloma in a Patient With Chronic Hepatitis C: A Case Report and Review of the Literature. *World J Gastroenterol* (2006) 12:2297. doi: 10.3748/wjg.v12.i14.2297
31. Panfilio S, D'Urso P, Annechini G, D'Elia GM, De Angelis F, Stefanizzi C, et al. Regression of a Case of Multiple Myeloma With Antiviral Treatment in a Patient With Chronic HCV Infection. *Leuk Res Rep* (2013) 2:39–40. doi: 10.1016/j.lrr.2013.01.002
32. Ioannou GN, Green PK, Berry K, Graf SA. Eradication of Hepatitis C Virus Is Associated With Reduction in Hematologic Malignancies: Major Differences Between Interferon and Direct-Acting Antivirals. *Hepatol Commun* (2019) 3:1124–36. doi: 10.1002/hep4.1389
33. Li G, Lewis RD, Mishra N, Axiotis CA. A Retrospective Analysis of Ten Symptomatic Multiple Myeloma Patients With HIV Infection: A Potential Therapeutic Effect of HAART in Multiple Myeloma. *Leuk Res* (2014) 38:1079–84. doi: 10.1016/j.leukres.2014.07.001
34. Pileri P, Uematsu Y, Campagnoli S, Galli G, Falugi F, Petracca R, et al. Binding of Hepatitis C Virus to CD81. *Science* (1998) 282(5390):938–41. doi: 10.1126/science.282.5390.938
35. Marcucci F, Mele A. Hepatitis Viruses and Non-Hodgkin Lymphoma: Epidemiology, Mechanisms of Tumorigenesis, and Therapeutic Opportunities. *Blood* (2011) 117:1792–8. doi: 10.1182/blood-2010-06-275818
36. Douam F, Lavillette D, Cosset F-L. The Mechanism of HCV Entry Into Host Cells. *Prog Mol Biol Transl Sci* (2015) 129:63–107. doi: 10.1016/bs.pmbts.2014.10.003
37. Zignego AL, Macchia D, Monti M, Thiers V, Mazzetti M, Foschi M, et al. Infection of Peripheral Mononuclear Blood Cells by Hepatitis C Virus. *J Hepatol* (1992) 15:382–6. doi: 10.1016/0168-8278(92)90073-X

Conflict of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Publisher's Note: All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Copyright © 2022 Rodríguez-García, Linares, Morales, Allain-Maillet, Mennesson, Sanchez, Alonso, Leivas, Pérez-Rivilla, Bigot-Corbel, Hermouet and Martínez-López. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.