



**HAL**  
open science

## **TV, computer, tablet and smartphone use and autism spectrum disorder risk in early childhood: a nationally-representative study**

Maria Melchior, Katharine Barry, David Cohen, Sabine Plancoulaine, Jonathan Bernard, Karen Milcent, Malamine Gassama, Ramchandar Gomajee, Marie-Aline Charles

### ► To cite this version:

Maria Melchior, Katharine Barry, David Cohen, Sabine Plancoulaine, Jonathan Bernard, et al.. TV, computer, tablet and smartphone use and autism spectrum disorder risk in early childhood: a nationally-representative study. *BMC Public Health*, 2022, 22 (1), pp.865. 10.1186/s12889-022-13296-5 . inserm-03693169

**HAL Id: inserm-03693169**

**<https://inserm.hal.science/inserm-03693169v1>**

Submitted on 10 Jun 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

RESEARCH

Open Access



# TV, computer, tablet and smartphone use and autism spectrum disorder risk in early childhood: a nationally-representative study

Maria Melchior<sup>1\*</sup>, Katharine Barry<sup>1</sup>, David Cohen<sup>2,3</sup>, Sabine Plancoulaine<sup>4</sup>, Jonathan Y. Bernard<sup>4,5</sup>, Karen Milcent<sup>6</sup>, Malamine Gassama<sup>6</sup>, Ramchandar Gomajee<sup>1</sup> and Marie-Aline Charles<sup>4,6</sup>

## Abstract

**Background:** Screen media use in early childhood has largely increased in recent years, even more so during the COVID-19 epidemic, and there is much discussion regarding its influence on neurodevelopment, including Autism Spectrum Disorder (ASD).

**Methods:** We examined the relationship between use of TV, computer, tablet and smartphone at age 2 years and risk of ASD assessed in telephone-based questionnaires among 12,950 children participating in the nationally representative ELFE ('Etude Longitudinale Française sur les Enfants') birth cohort study in France.

**Results:** In inverse-probability weighted (IPW) multinomial regression analyses, children's weekly or daily screen media use was associated with an increased likelihood of an intermediate risk of ASD (IPW-controlled OR for weekly use: 1.07, 95% CI 1.02—1.12; IPW-controlled OR for daily use: 1.05, 95% CI 1.02—1.08) but inversely associated with a high risk (IPW-controlled OR for weekly use: 0.60, 95% CI 0.50—0.73; IPW-controlled OR for daily use: 0.75, 95% CI 0.62—0.91), as ascertained by the M-CHAT. This was confirmed when studying TV as well as computer/tablet exposure separately.

**Conclusions:** Overall, our nationally-representative study conducted among a large sample of 2-year-old children, indicates a complex relationship between screen exposure and ASD risk.

**Keywords:** Screen media, Autism spectrum disorder, Birth cohort study, ELFE

## Background

While children's use of screen-based media is ubiquitous across high-income countries, and seems to have increased since the beginning of the COVID-19 pandemic [1], growing evidence suggests that exposure in infancy and early childhood is associated with unfavourable developmental outcomes over the short and mid-term [2–5]. In particular, there has been

suggestion that high levels of screen-based media viewing are related to a higher level of risk of autism-spectrum disorder (ASD) [2, 6].

Although genetics play an important role with regard to the risk of ASD and symptoms of neurodevelopmental delay, evidence suggests that certain environmental exposures could also be involved [7]. Media screen exposure is a candidate, as children who early on in life spend a lot of time watching screens – particularly if their content is poor and/or if they are unsupervised—are less likely to play, be physically active, and engage in interactions with other children and adults, all of which could be related to worse neurodevelopmental outcomes [3, 5]. However,

\*Correspondence: maria.melchior@inserm.fr

<sup>1</sup> Sorbonne Université, Inserm, Institut Pierre Louis d'Epidémiologie Et de Santé Publique (IPLESP), Paris, France

Full list of author information is available at the end of the article



while it has been documented that children and adolescents diagnosed with ASD are more exposed to screens [2], research examining whether early screen exposure precedes later ASD diagnosis has been limited. A recent study based on 2,100 US-based children participating in the nationally representative National Children's Study (NCS), found no association between exposure to TV/DVD watching at 12 and 18 months and a high risk of ASD at age 2 years, although the authors did report a linear association with the number of ASD symptoms [8]. To our knowledge, associations between the use of other forms of screen media, including computers/tablets and smartphones and children's risk of ASD, have not yet been evaluated.

Our aim was to evaluate the relationship between use of TV, computer/tablet and smartphone in relation to neurodevelopmental difficulties and particularly ASD risk in early childhood, in a general population setting.

## Methods

Data for this study come from the ELFE ("Étude Longitudinale Française depuis l'Enfance") study, a nationally representative birth cohort of 18,329 children born in France in 2011 in 320 maternity wards selected using random sampling methods, with the aims of studying children's developmental and health trajectories through adulthood [9]. Inclusion criteria were: singleton or twin birth at  $\geq 33$  weeks of gestation, mother of  $\geq 18$  years of age and not planning to move outside of Metropolitan France in the three years following study inclusion. Mothers had to be able to give consent either in French, English, Arabic or Turkish. Briefly, participating mothers were invited to take part in the study and completed a research questionnaire after delivery during their maternity stay, then mothers and fathers were contacted again for a telephone interview when the child was aged 2 months, 1 year and 2 years.

In the 2-year ELFE study telephone interview, parents reported on their child's TV watching, use of a computer/tablet and smartphone. If the child watched television or used a computer/tablet, parents were additionally asked the number of hours during weekdays and weekends, based on which we estimated average exposure. Frequency of use of television and computers/tablets being very distinct, we estimated the level of exposure in hours per day and per week, respectively.

Risk of neurodevelopmental difficulties at 2 years was assessed using the Modified Checklist for Autism in Toddlers (M-CHAT), a validated 23-item screening instrument [10, 11]. We used the recommended cut-offs to define risk of ASD: low (0–2), intermediate (3–6), and high (7–23) risk.

Because children who use screens and those who do not are likely to differ with regard to many different characteristics, covariates were controlled for using propensity scores, which were calculated based on the variables listed below [12]:

1. *Family socio-demographic characteristics* reported by the mother after delivery: maternal age (<30 vs.  $\geq 30$  years – the average age at first childbirth for women), paternal age (<30 vs.  $\geq 30$  years), maternal and paternal birth place (France vs. Other), mother's educational level, number of children living in the household (1, 2–3,  $\geq 4$ ), maternal and paternal residential area (Greater Paris Region; Paris and suburbs, North-West, East, West, South-West, Centre-East, and South-East) and household income per capita assessed when the child was 2 years (divided into 3 categories based on the minimum wage, average wage and above average wage in 2013, <https://www.insee.fr/fr/statistiques/1375188>) (<1,423; 1,423–1,996; >1,996 euros/month).
2. *Maternal mental health* reported by the mother when the child was 2 months old (Edinburgh Postnatal Depression Scale (EPDS) – a score of  $\geq 12$  being considered as indicative of possible depression, yes vs. no [13]).
3. *Pregnancy and child characteristics* collected from the child's medical records: child sex (male vs. female), age at the time of assessment ( $\leq 24$ , 25, 26, or  $> 26$  months), season of birth (winter; spring; summer; autumn), prematurity (<36 vs.  $\geq 37$  weeks of gestation), if the child was a twin (yes vs. no), weight for gestational age (based on a z-score and categorized as small (<10<sup>th</sup> percentile), average (10–90<sup>th</sup> percentile), or large (>90 percentile)), and Apgar score at 5 min (alarming/fair vs. good).
4. *Postnatal characteristics* reported by the mother at age 2 years: childcare arrangements (parents or family member, centre-based childcare, nanny or childcare professional) and the number of early stimulation and development activities (ex. painting/drawing, storytelling, singing, reading, doing puzzles) the child completed with his/her parents in the preceding month ( $\leq 4$ , 5–6, 7–9,  $> 9$  activities).

## Statistical analyses

Using survey-weighted data, which takes into account the initial sampling plan and baseline participation as well as longitudinal attrition rates to produce nationally representative estimates for children born in 2011 (<https://www.elfe-france.fr/fichier/rte/178/Cote%20rec%20herche/Weighting-Elfe-surveys-general-document.pdf>),

we assessed associations between screen media use and children's neurodevelopmental risk (high, intermediate vs. low as reference) at age 2 years. Missing covariate data (maximum 3.1%) were imputed 10 times using fully conditional specification (MI-FCS). To balance differences between screen exposure groups, propensity score weights were estimated with a generalized boosting model (GBM) using Kolmogorov–Smirnov (KS) means. Then, multinomial logistic regression models weighted on the product of Inverse Probability Weights (IPW) and survey weights, yielded Odds Ratios (ORs) with the associated 95% Confidence interval (95% CI) and a  $P < 0.05$  examining statistical significance [14]. Missing covariate data (maximum 3.1%) were imputed 10 times with fully conditional specification (MI-FCS). All data were analysed using R 4.0.3 and SAS V9.4.

## Results

A total of 12,950 children were included in the present analysis (Table 1), with 12,012 children with available information on weekly TV, computer or tablet use, and respectively 11,609 and 12,498 children with available information on the frequency of TV and computer/tablet use. Respectively 29.9 and 1.1% of children were classified as having an intermediate and high risk of neurodevelopmental difficulties. 66.4% of children used screen media daily, mostly TV (63.3%, 9.8% being exposed  $> 2$  h/day); 30.5% used a computer/tablet weekly and 26.7% used a smartphone. As shown in Table 1, all studied covariates were associated with children's screen media use.

In IPW-controlled analyses (Table 2), compared to never or seldom-users, children who used screen media weekly or daily had an increased likelihood of being at intermediate risk of neurodevelopmental difficulties (ORs between 1.05–1.07), but a decreased likelihood of being at high risk (ORs between 0.60–0.75). Compared to children who never watched TV daily, had an increased likelihood of intermediate level risk (OR associated with  $> 2$  h per day 1.11, 95% CI 1.06–1.17), but a decreased likelihood of a high risk of neurodevelopmental difficulties (ORs associated with 30 mn to  $> 2$  h between 0.33–0.60). Compared to never or seldom-users, children who used a computer/tablet  $\leq 1$  h/day had a decreased likelihood of being at intermediate risk or high risk of neurodevelopmental difficulties (ORs: 0.94, 95% CI 0.91–0.96 and 0.24, 95% CI 0.22–0.26, respectively); children who used a Tables 1 and 2 hours per week also had a low likelihood of being at high risk of neurodevelopmental difficulties (OR 0.83, 95% CI 0.70–0.98).

## Discussion

In a nationally-representative study of 2-year old children, a majority were exposed to screen media daily. In IPW-controlled analyses, screen media use, mainly TV, was associated with an increased likelihood of intermediate level risk of neurodevelopmental difficulties, but a reduced likelihood of a high risk. Our study adds to a growing literature showing elevated levels of screen-based media use among young children and possible associations with development. However, several studies showed elevated levels of screen use among children at risk of autism spectrum disorder [6, 8] and our data temper the results of prior investigations. Our data suggest a complex relationship between screen-based activities and risk of neurodevelopmental difficulties, including autism risk.

We need to acknowledge several limitations. First, ELFE is based on parents' voluntary participation and may have left out children vulnerable to neurodevelopmental difficulties [9]. Nevertheless, the sample is heterogeneous-enough to study environmental risk factors of neurodevelopment [12, 15] and our data were weighted to be nationally representative. Second, ELFE is a longitudinal cohort study, but the data we used are cross-sectional and we cannot rule out that children's neurodevelopmental difficulties could influence their screen use. This may in part explain lower levels of exposure among children at high risk of neurodevelopmental difficulties. To address this issue, our statistical analyses controlled for a range of covariates which precede and predict children's screen use as well as ASD symptoms, which is a way of taking into account common sources of variation. Future studies with prospective designs are warranted to assess the role of early screen use on children's risk of neurodevelopmental difficulties, particularly those that are within the range of ASD. Third, we used a single M-CHAT measure, which is probably less specific than repeated assessments and it is unlikely that all children identified as having an intermediate or high risk will actually be diagnosed with ASD [10]. This results in the possibility of misclassification, particularly in the intermediate risk group, and calls for close monitoring of children's later neurodevelopmental outcomes [11]. Fourth, ELFE study participants were born in 2011, and children's levels of screen use have since increased, particularly during the COVID-19 pandemic [1]. This implies that prevalence levels of screen exposure at 2 years are probably currently be higher than we report. However, the associations we observe between children's media screen use and risk of neurodevelopmental difficulties are likely to be valid in current circumstance or may have increased.

**Table 1** Description of family socio-demographic, maternal mental health, pregnancy and child, postnatal and MCHAT score variables by screen exposure on survey-weighted data (%), ELFE cohort, France, 2011–2013

Variables	Tv/computer/tablet/smartphone use (N = 12 950)				P-value
	Rarely/never (1714)	Monthly (N = 521)	Weekly N = 2517)	Daily (N = 8198)	
<b>Family socio-demographic characteristics</b>					
<b>Maternal age: ≥ 30 years</b>	55.7%	59.6%	59.8%	49.8%	<.0001
<b>Paternal age: ≥ 30 years</b>	67.6%	73.4%	72.9%	71.1%	<.0001
<b>Mother born in France</b>	88.3%	89.8%	86.3%	80.8%	<.0001
<b>Father born in France</b>	87.3%	87.4%	86.8%	80.5%	<.0001
<b>Mother's educational level</b>					<.0001
<i>No diploma</i>	6.1%	2.0%	6.8%	10.1%	
<i>&lt; Bachelor's degree</i>	38.9%	29.0%	30.3%	45.1%	
<i>Bachelor's degree</i>	16.6%	18.4%	20.8%	19.9%	
<i>≥ Master's degree</i>	38.4%	50.5%	42.0%	24.9%	
<b>Number of children in the household</b>					<.0001
<i>1 child</i>	42.5%	46.5%	40.0%	44.4%	
<i>2–3 children</i>	51.0%	50.5%	53.5%	48.4%	
<i>≥ 4 children</i>	51.0%	50.5%	53.5%	48.4%	
<b>Maternal residential area</b>					<.0001
<i>Greater Paris Region</i>	19.9%	23.7%	20.6%	19.5%	
<i>Paris and suburbs</i>	19.4%	16.2%	15.2%	18.5%	
<i>North-West</i>	5.1%	3.6%	5.9%	7.9%	
<i>East</i>	7.7%	6.9%	7.8%	9.0%	
<i>West</i>	16.5%	16.9%	16.5%	13.3%	
<i>South-West</i>	8.4%	6.4%	9.7%	8.3%	
<i>Centre-East</i>	12.2%	16.0%	11.5%	10.9%	
<i>South-East</i>	10.8%	10.4%	12.8%	12.7%	
<b>Paternal residential area</b>					<.0001
<i>Greater Paris Region</i>	19.8%	23.5%	20.7%	19.4%	
<i>Paris Basin</i>	19.6%	16.3%	15.1%	18.5%	
<i>North-West</i>	5.1%	3.6%	5.9%	7.9%	
<i>East</i>	7.7%	6.9%	7.8%	9.0%	
<i>West</i>	16.5%	16.9%	16.8%	13.2%	
<i>South-West</i>	8.4%	6.4%	9.6%	8.4%	
<i>Centre-East</i>	12.2%	16.0%	11.5%	10.9%	
<i>South-East</i>	10.8%	10.4%	12.7%	12.6%	
<b>Household monthly income</b>					<.0001
<i>&lt; 1,423 euros</i>	9.1%	5.0%	8.1%	1.5%	
<i>1,423–1,996 euros</i>	7.4%	8.2%	6.7%	8.3%	
<i>&gt; 1,996 euros</i>	83.4%	86.8%	85.1%	76.7%	
<b>Maternal depression</b>	12.1%	10.8%	13.0%	15.3%	<.0001
<i>Pregnancy and child characteristics</i>					
<b>Child's sex (female)</b>	48.2%	54.0%	47.8%	49.8%	<.0001
<b>Child's age at interview</b>					<.0001
<i>≤ 24 months</i>	19.4%	19.3%	18.9%	18.3%	
<i>25 months</i>	29.2%	34.7%	34.0%	33.6%	
<i>26 months</i>	25.4%	27.7%	28.2%	27.5%	
<i>&gt; 26 months</i>	25.9%	18.3%	19.7%	20.6%	
<b>Twin birth</b>	6.0%	3.3%	2.3%	3.1%	<.0001

**Table 1** (continued)

Variables	Tv/computer/tablet/smartphone use (N = 12 950)				P-value
	Rarely/never (1714)	Monthly (N = 521)	Weekly N = 2517)	Daily (N = 8198)	
<b>Family socio-demographic characteristics</b>					
<b>Season of birth</b>					<.0001
Spring	18.7%	25.1%	22.6%	22.0%	
Summer	31.1%	29.4%	27.8%	24.9%	
Autumn	25.0%	18.7%	24.9%	27.8%	
Winter	25.3%	26.8%	24.6%	25.3%	
<b>Premature birth (&lt; 37 weeks of gestation)</b>	5.1%	7.0%	5.0%	5.2%	<.0001
<b>Birth weight for gestational age</b>					<.0001
Small (< 10 percentile)	3.5%	3.7%	1.5%	2.2%	
Average (10–90 percentile)	93.6%	94.0%	95.3%	95.0%	
Large (> 90 percentile)	2.8%	2.2%	3.1%	2.8%	
<b>Apgar score (good)</b>	98.4%	99.9%	99.1%	98.6%	<.0001
Postnatal characteristics					
<b>Childcare arrangements</b>					<.0001
Parents or family member	39.8%	28.1%	32.1%	46.5%	
Centre-based care	22.2%	24.8%	24.8%	18.1%	
Nanny or professional	38.0%	47.1%	43.1%	35.3%	
<b>Activities with the child</b>					<.0001
≤ 4 activities	7.5%	83.7%	69.6%	7.1%	
5–6 activities	16.6%	15.1%	15.5%	16.6%	
7–8 activities	25.4%	25.4%	27.8%	28.0%	
≥ 9 activities	50.4%	51.1%	49.7%	48.3%	
<b>Risk of ASD</b>					<.0001
Low	77.3%	77.9%	78.0%	75.2%	
Intermediate	21.4%	21.3%	21.3%	23.9%	
High	1.2%	0.8%	0.6%	0.9%	

Our study also has strengths which we would like to highlight. First, ours is one of few studies to examine the early risk of neurodevelopmental difficulties in a community-based nationally representative sample of young children. Second, to our knowledge, this is the first study in a general population setting to include a measure of computer/tablet and smartphone use in early childhood and study the relationship with neurodevelopment. One of the hypotheses explaining this association is the effect of electromagnetic field exposure on children's brain development [6]. Third, we used propensity scores and inverse-probability weights to control for selection biases and for a large number of child and family characteristics predating measures of screen media use that could confound the association of interest, ruling out a number of alternative explanations.

Overall, young children exposed to screen media may experience some neurodevelopmental delays, captured by the intermediate risk category on the M-CHAT. Our

data suggest that TV use seems more strongly associated with children's ASD risk than other types of screen media. This may reflect children's greater passivity in front of a TV screen, which should be confirmed by other research, using quantitative as well as qualitative designs. This association may reflect reduced play time and physical activity, as well as more limited interactions with other children and adults [5]. Low levels of screen use among children at high risk of neurodevelopmental difficulties suggest limited interest or specific parental behaviours in some – rather highly educated – families which restrict children's exposure [16]. There is suggestion that early environmental exposures such as screen media exposure could increase the experience of symptoms evocative of ASD risk, however this should be further investigated in studies which follow longitudinally large samples of children potentially at risk of ASD.

**Table 2** Associations between screen use and risk of autism risk (M-CHAT score): Inverse probability weighted (IPW), survey weighted, multinomial logistic regression model: Odds Ratios (OR), 95% Confidence Interval (CI). ELFE cohort, France, 2011–2013

	Intermediate risk of autism OR (95%CI)	P-value	High risk of autism OR (95%CI)	P-value
<b>Overall screen media use</b>				
<b>TV/computer/tablet/smartphone use</b>				
Rarely/never (n = 1714)	Reference		Reference	
Monthly (n = 521)	1.03 (0.93—1.14)	.48	0.89 (0.72—1.10)	.26
Weekly (n = 2,517)	1.07 (1.02—1.12)	.006	0.60 (0.50—0.73)	.002
Daily (n = 8,198)	1.05 (1.02—1.08)	.003	0.75 (0.62—0.91)	.008
<b>Number of hours of screen media use</b>				
<b>TV/Computer/tablet (hrs/week)</b>				
Never (n = 2124)	Reference		Reference	
≤ 1 h (n = 1198)	0.96 (0.90- 1.02)	.15	0.72 (0.61—0.86)	.002
1–2 h (n = 1292)	1.03 (0.99- 1.08)	.13	0.09 (0.08—0.11)	< .001
> 2 h (n = 7884)	1.01 (0.99—1.03)	.31	0.61 (0.54—0.68)	< .001
<b>TV (hrs/day)</b>				
Never (n = 1845)	Reference		Reference	
≤ 30 min (n = 3632)	1.03 (0.99—1.08)	.10	0.79 (0.63—0.99)	.04
30 min-1 h (n = 3029)	1.00 (0.96—1.05)	.80	0.33 (0.27—0.41)	< .001
1–2 h (n = 2257)	1.02 (0.98—1.05)	.31	0.60 (0.49—0.74)	.002
≥ 2 h (n = 846)	1.11 (1.06—1.17)	.006	0.52 (0.37—0.73)	.002
<b>Computer/tablet use (hrs/week)</b>				
Never (n = 8485)	Reference		Reference	
≤ 1 h (n = 2150)	0.94 (0.91—0.96)	.004	0.24 (0.22—0.26)	< .001
1–2 h (n = 781)	1.02 (0.98—1.05)	.30	0.82 (0.70—0.98)	.03
> 2 h (n = 596)	0.94 (0.85—1.04)	.19	0.83 (0.64 – 1.07)	.13

## Conclusion

Overall, screen media, mainly TV, use early on in life is related to children's neurodevelopmental difficulties, but these associations appear to be complex.

## Abbreviations

OR: Odds Ratio; ASD: Autism-Spectrum Disorder; IPW: Inverse-Probability Weight; MI-FCS: Missing data Imputation -Fully Conditional Specification.

## Acknowledgements

The ELFE study is a joint project between the French Institute for Demographic Studies (INED) and the National Institute of Health and Medical Research (INSERM), in partnership with the French blood transfusion service (Etablissement français du sang, EFS), Santé Publique France, the National Institute for Statistics and Economic Studies (INSEE), the Direction Générale de la Santé (DGS, part of the Ministry of Health and Social Affairs), the Direction Générale de la Prévention des Risques (DGPR, Ministry for the Environment), the Direction de la Recherche, des Etudes, de l'Evaluation et des Statistiques (DREES, Ministry of Health and Social Affairs), the Département des Etudes, de la Prospective et des Statistiques (DEPS, Ministry of Culture), and the Caisse Nationale des Allocations Familiales (CNAF), with the support of the Ministry of Higher Education and Research and the Institut National de la Jeunesse et de l'Éducation Populaire (INJEP).

## Authors' contributions

Dr. Charles had the initial idea for the study, is responsible for data acquisition, and suggested initial statistical analyses. Drs. Plancoulaine, Bernard and Milcent as well as M. Gassama contributed to crafting the study design and initiated

statistical analyses. Ms. Barry and M. Gomajee pursued and finalized statistical analyses. Dr. Cohen helped interpret the study findings and contributed to draft the manuscript. Dr. Melchior supervised statistical analyses and drafted the first version of the manuscript. All authors read and approved the final manuscript.

## Funding

The ELFE cohort is supported by state funding from the ANR (Agence Nationale de la Recherche) within the framework of the "Future Investments" programme (reference: ANR-11-EQPX-0038; ANR-19-COHO-001), as part of the RECONAI platform. This research is further supported by ANR funding under the reference ANR-20-CE36-0001 and an ERC-Consolidator grant (RESEDA, ERC-2020-COG-101001420).

## Availability of data and materials

The datasets analysed during the current study are not publicly available due to regulatory restrictions in place in France but are available from the corresponding author on reasonable request.

## Declarations

### Ethics approval and consent to participate

The ELFE study received the approval of France's bodies regulating ethical research conduct (Comité Consultatif sur le Traitement des Informations pour la Recherche en Santé: CCTIRS; Commission de Protection des Personnes: CPP; Commission National Informatique et Libertés: CNIL). Written informed consent was obtained from all parents of ELFE participants. All methods were carried out in accordance with relevant guidelines and regulations.

### Consent for publication

Not applicable.

**Competing interests**

The authors have no competing interests to declare.

**Author details**

<sup>1</sup>Sorbonne Université, Inserm, Institut Pierre Louis d'Epidémiologie Et de Santé Publique (IPLESP), Paris, France. <sup>2</sup>Department of Child and Adolescent Psychiatry, Pitié-Salpêtrière Hospital, Paris, France. <sup>3</sup>CNRS UMR 7222, Institute for Intelligent Systems and Robotics, Sorbonne University, Paris, France. <sup>4</sup>Centre for Research in Epidemiology and Statistics (CRESS), Université de Paris, Inserm, Inrae, 16 Av Paul Vaillant-Couturier, 75004 Paris, France. <sup>5</sup>Agency for Science, Technology and Research (A\*STAR), Singapore Institute for Clinical Sciences (SICS), Singapore, Singapore. <sup>6</sup>UMS Elfe Team, Ined, Inserm, EFS, Aubervilliers, France.

Received: 25 January 2022 Accepted: 25 April 2022

Published online: 30 April 2022

**References**

- McArthur BA, Racine N, Browne D, McDonald S, Tough S, Madigan S. Recreational screen time before and during COVID-19 in school-aged children. *Acta Paediatr.* 2021;110(10):2805–7.
- Slobodin O, Heffler KF, Davidovitch M. Screen media and autism spectrum disorder: a systematic literature review. *J Dev Behav Pediatr.* 2019;40(4):303–11.
- Madigan S, Browne D, Racine N, Mori C, Tough S. Association between screen time and children's performance on a developmental screening test. *JAMA Pediatr.* 2019;173(3):244–50.
- Fullwood I. Increased screen time is associated with poorer developmental outcomes in early childhood. *Arch Dis Child.* 2020;105(4):253–4.
- Madigan S, McArthur BA, Anhorn C, Eirich R, Christakis DA. Associations between screen use and child language skills: a systematic review and meta-analysis. *JAMA Pediatr.* 2020;174(7):665–75.
- Kushima M, Kojima R, Shinohara R, Horiuchi S, Otawa S, Ooka T, et al. Association between screen time exposure in children at 1 year of age and autism spectrum disorder at 3 years of age: the Japan Environment and Children's Study. *JAMA Pediatrics.* 2022 [In Press].
- Schmengler H, Cohen D, Tordjman S, Melchior M. Autism spectrum and other neurodevelopmental disorders in children of immigrants: a brief review of current evidence and implications for clinical practice. *Front Psych.* 2021;12:566368.
- Heffler KF, Sienko DM, Subedi K, McCann KA, Bennett DS. Association of early-life social and digital media experiences with development of autism spectrum disorder-like symptoms. *JAMA Pediatr.* 2020;174(7):690–6.
- Charles MA, Thierry X, Lanoe JL, Bois C, Dufourg MN, Popa R, et al. Cohort profile: the French National cohort of children ELFE: birth to 5 years. *Int J Epidemiol.* 2020;49(2):368–9.
- Chlebowski C, Robins D, Barton M, Fein D. Large-scale use of the modified checklist for autism in low-risk toddlers. *Pediatrics.* 2013;131(4):e1121–7.
- Baduel S, Guillon Q, Afzali MH, Foudon N, Kruck J, Roge B. The French version of the Modified-Checklist for Autism in Toddlers (M-CHAT): a validation study on a French sample of 24 month-old children. *J Autism Dev Disord.* 2017;47(2):297–304.
- Schmengler H, El-Khoury Lesueur F, Yermachenko A, Taine M, Cohen D, Peyre H, et al. Maternal immigrant status and signs of neurodevelopmental problems in early childhood: the French representative ELFE birth cohort. *Autism Res.* 2019;12(12):1845–59.
- Levis B, Negeri Z, Sun Y, Benedetti A, Thombs BD. Accuracy of the Edinburgh Postnatal Depression Scale (EPDS) for screening to detect major depression among pregnant and postpartum women: systematic review and meta-analysis of individual participant data. *BMJ.* 2020;371:m4022.
- Hernan M, Robbins J. IP weighting and marginal structural models. *Causal Inference* 2013. p. 11–21.
- Nakamura A, El-Khoury Lesueur F, Sutter-Dallay AL, Franck J, Thierry X, Melchior M, et al. The role of prenatal social support in social inequalities with regard to maternal postpartum depression according to migrant status. *J Affect Disord.* 2020;272:465–73.
- Dong HY, Wang B, Li HH, Yue XJ, Jia FY. Correlation between screen time and autistic symptoms as well as development quotients in children with autism spectrum disorder. *Front Psych.* 2021;12: 619994.

**Publisher's Note**

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more [biomedcentral.com/submissions](https://biomedcentral.com/submissions)

