



Early-Life Environment Influence on Late-Onset Alzheimer's Disease

Thibaut Gauvrit, Hamza Benderradji, Luc Buée, David Blum, Didier Vieau

► To cite this version:

Thibaut Gauvrit, Hamza Benderradji, Luc Buée, David Blum, Didier Vieau. Early-Life Environment Influence on Late-Onset Alzheimer's Disease. *Frontiers in Cell and Developmental Biology*, 2022, 10, pp.834661. 10.3389/fcell.2022.834661 . inserm-03699318

HAL Id: inserm-03699318

<https://inserm.hal.science/inserm-03699318>

Submitted on 20 Jun 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Early-Life Environment Influence on Late-Onset Alzheimer's Disease

Thibaut Gauvrit^{1,2}, Hamza Benderradji^{1,2}, Luc Buée^{1,2}, David Blum^{1,2} and Didier Vieau^{1,2*}

¹Inserm, CHU Lille, U1172 LiNCog—Lille Neuroscience and Cognition, Université de Lille, Lille, France, ²Alzheimer and Tauopathies, LabEx DISTALZ, Lille, France

OPEN ACCESS

Edited by:

Elena Zambrano,
Instituto Nacional de Ciencias Médicas
y Nutrición Salvador Zubirán
(INCMNSZ), Mexico

Reviewed by:

Sofía Díaz Cintra,
National Autonomous University of
Mexico, Mexico
Andrea Fuso,
Sapienza University of Rome, Italy

*Correspondence:

Didier Vieau
didier.vieau@univ-lille.fr

Specialty section:

This article was submitted to
Epigenomics and Epigenetics,
a section of the journal
Frontiers in Cell and Developmental
Biology

Received: 13 December 2021

Accepted: 27 January 2022

Published: 17 February 2022

Citation:

Gauvrit T, Benderradji H, Buée L,
Blum D and Vieau D (2022) Early-Life
Environment Influence on Late-Onset
Alzheimer's Disease.
Front. Cell Dev. Biol. 10:834661.
doi: 10.3389/fcell.2022.834661

With the expand of the population's average age, the incidence of neurodegenerative disorders has dramatically increased over the last decades. Alzheimer disease (AD) which is the most prevalent neurodegenerative disease is mostly sporadic and primarily characterized by cognitive deficits and neuropathological lesions such as amyloid β ($A\beta$) plaques and neurofibrillary tangles composed of hyper- and/or abnormally phosphorylated Tau protein. AD is considered a complex disease that arises from the interaction between environmental and genetic factors, modulated by epigenetic mechanisms. Besides the well-described cognitive decline, AD patients also exhibit metabolic impairments. Metabolic and cognitive perturbations are indeed frequently observed in the Developmental Origin of Health and Diseases (DOHaD) field of research which proposes that environmental perturbations during the perinatal period determine the susceptibility to pathological conditions later in life. In this review, we explored the potential influence of early environmental exposure to risk factors (maternal stress, malnutrition, xenobiotics, chemical factors . . .) and the involvement of epigenetic mechanisms on the programming of late-onset AD. Animal models indicate that offspring exposed to early-life stress during gestation and/or lactation increase both AD lesions, lead to defects in synaptic plasticity and finally to cognitive impairments. This long-lasting epigenetic programming could be modulated by factors such as nutraceuticals, epigenetic modifiers or psychosocial behaviour, offering thus future therapeutic opportunity to protect from AD development.

Keywords: Alzheimer disease, perinatal environment, epigenetic mechanisms, early-life stress, tau, amyloid- β

INTRODUCTION

Alzheimer disease (AD), for which only symptomatic treatments are currently available, is a chronic neurodegenerative disorder and the most prevalence form of dementia. In 2020, there were over 50 million people worldwide living with dementia and this number is estimated to double every 20 years (World Health Organization, 2021). AD is a major health problem characterized by the progressive and irreversible development of neuronal damages, leading to the decline in cognitive functions, notably memory. AD, whose diagnosis is firmly established post-mortem, is defined by the extracellular accumulation of amyloid β ($A\beta$) peptides into extracellular amyloid plaques (Checler, 1995) and the presence of neuronal neurofibrillary tangles (NFT) made up of intraneuronal fibrillar aggregates of hyper- and/or abnormally phosphorylated Tau proteins (« Tau pathology »), a common feature of other neurodegenerative disorders called tauopathies (Sergeant et al., 2008). In AD, spreading of the neurofibrillary lesions in the cortex (first entorhinal cortex, then hippocampus and lastly neocortex) fits to the progression of the

symptoms, supporting that Tau pathology is indeed instrumental in cognitive alterations (for review Colin et al., 2020). In agreement, Tau pathology development impairs various forms of synaptic plasticity and cognitive behaviors in mouse models (Polydoro et al., 2009; Hoover et al., 2010; Van der Jeugd et al., 2011; Burnouf et al., 2013; Lo et al., 2013; Van der Jeugd et al., 2013). In most cases, AD appears as a multifactorial disease resulting from the complex interaction between genetic predisposition and epigenetic environmental factors (Reitz et al., 2011; Armstrong, 2019). Epidemiological studies identified protective factors such as exercise, intellectual activities and consumption of fish or caffeine (La Rue, 2010; Flaten et al., 2014; Najar et al., 2019). By contrast, life-stresses, preceding clinical manifestation of AD, drive the progression of the disease and can exacerbate symptoms (Reitz and Mayeux, 2009; Reitz et al., 2011). In addition, apart from aging itself, cardiovascular dysfunctions such as obesity, hypertension and diabetes have been suggested as important risk factors for AD (Craft et al., 1998; Tortelli et al., 2017; Flores-Dorantes et al., 2020; Livingston et al., 2020; Tini et al., 2020). It is also notable that AD patients have been reported to exhibit altered glucose metabolism (Bucht et al., 1983; Craft et al., 1992; Craft et al., 1998; Matsuzaki et al., 2010) and even present an increased prevalence to develop type 2 diabetes (Janson et al., 2004). Presence of cognitive and metabolic alterations in AD patients fits with the “Type 3 diabetes” concept, stating that AD brain is insulin-resistant (Talbot et al., 2012; de la Monte, 2014; de la Monte et al., 2018; Nguyen et al., 2020). Although the pathophysiological mechanisms linking AD, insulin resistance and impaired glucose homeostasis remain to be clarified, it has been reported that A β oligomers impair insulin signalling by promoting insulin neuronal receptor internalization (Zhao et al., 2008) and inhibit insulin receptor substrate (IRS) 1 (Bomfim et al., 2012). In addition, using Tau knock-out mice, we and others showed that Tau loss-of-function, which is likely a consequence of Tau aggregation, might contribute to brain insulin sensitivity and metabolic impairments (Marciniak et al., 2017; Wijesekara et al., 2018; Wijesekara et al., 2021); an effect reversed by human Tau re-expression (Wijesekara et al., 2021). In line, the H1 Tau haplotype was associated to the glucose homeostasis in humans (Marciniak et al., 2017).

Interestingly, the pioneering epidemiological studies of David Barker in the early 90s established a relationship between intra-uterine-growth retardation (IUGR) and cardiovascular diseases in adulthood paving the way to a new field of research now known as DOHaD (for Developmental Origin of Health and Diseases) stating that environment during early life may shape the rest of our life (Barker, 2007). In accordance, numerous epidemiological studies in humans and experimental works in animals have clearly reinforced the idea that adverse perinatal environment, such as maternal stress, exposure to toxics and early-life malnutrition, has long-term consequences and may program, in the offspring, chronic adult diseases such as glucose dyshomeostasis and energy metabolism impairments but also cognitive disorders (Moody et al., 2017; Gawlińska et al., 2021b), all being encountered in AD patients. In light of these observations, altered perinatal neurodevelopment might

sensitize to the occurrence of late-onset AD (Moceri et al., 2000; Landrigan et al., 2005; Miller and O’Callaghan, 2008; Modgil et al., 2014; Athanasopoulos et al., 2016) presumably *via* epigenetic mechanisms (Gapp et al., 2014; Lemche, 2018). In this review, we will summarize the most recent data suggesting that AD, that is usually associated with metabolic impairments developing at the adult stage and aging, may also date back to very early life. The identification of early mechanisms linking parental environment with modification of brain’s epigenetic landscape in the offspring may offer new therapeutic strategies to prevent the progression of AD and other neurodegenerative diseases.

TAU AND ABPP DURING EARLY-NEURODEVELOPMENT

Maturation of central nervous system (CNS) and establishment of neuronal connections that vary in a region-specific manner can be divided schematically in two major developmental time-periods. The first period that occurs predominantly in the embryonic developmental stages in rodents and within the first two trimesters in humans, is dedicated to neurogenesis, neuronal migration and polarization whereas the second period covering lactation in rodents and the last trimester of gestation in humans is involved in intense neurite stabilization, synapse formation and establishment of brain circuitry (de Graaf-Peters and Hadders-Algra, 2006; Metzger, 2010; Cisneros-Franco et al., 2020). This indicates that the perinatal period covering both gestation and lactation constitutes a particularly sensitive critical time-window during which environmental perturbations may modify brain circuitry and may thus exert long-lasting effects on brain function, synaptic plasticity, homeostasis regulation and cognitive functions. Although the role of Tau and the metabolism of amyloid- β precursor protein (A β PP), the precursor to A β peptides, have been extensively described in the context of AD development, few studies have reported their role during early-neurodevelopment. For instance, A β PP acts as a cell adhesion molecule during different steps of neurodevelopment (Sosa et al., 2017) and may thus participate to the migration and maturation of newborn neurons as well as the formation of synapses. *Via* its large ectodomain, A β PP has been shown to bind extracellular matrix (ECM) molecules which play fundamental roles in the formation and maintenance of brain architecture during neurodevelopment and in adulthood (Dityatev et al., 2010; Mouw et al., 2014). Moreover A β PP, *via* dimerization and binding to ECM, participates to the migration and adaptation of newborn neurons to the early environment (Marín et al., 2010; Parsons et al., 2010; Cooper, 2013). Although the developmental role of Tau remains poorly understood, Tau is highly expressed in both rodent and human fetal brains with developmental changes in both splicing and phosphorylation. Resulting from an alternative splicing mechanism, six major isoforms of Tau coexist in the human brain with the presence of either 3 or 4 repeated sequences (named below as Tau-3R or Tau-4R) known as the microtubule-binding regions (Sergeant et al., 2008). Tau-3R short-isoform, being most abundant in fetal brain, has less affinity with microtubules, and increases neuronal plasticity

(Wang and Liu, 2008; Hefti et al., 2018; Hefti et al., 2019). In rats, the total level of Tau in the brain remains stable after birth and slightly increased during lactation with the appearance of larger Tau isoforms from post-natal day (PND) 5 to PND15 and the concomitant decrease of fetal Tau (Yu et al., 2009). Despite Tau knock-out mice showed moderate neurologic alterations (Ikegami et al., 2000; Lei et al., 2012), presumably due to compensatory mechanisms involving other microtubules associated proteins, *in vitro* studies in cultured mouse and hamster neurons suggested that Tau is involved in axogenesis, neurite outgrowth and neuronal circuit formation (Dawson et al., 2001; Sennvik et al., 2007; Biswas and Kalil, 2018). Recently, it has been proposed that Tau phosphorylation at microtubule domain region at early stages could protect the hippocampal circuit from overexcitation in 1-month-old 3xTg mice (a model of AD) by directly interacting with the pyramidal circuitry that spontaneously generates theta oscillations (Mondragón-Rodríguez et al., 2018). These results suggest that Tau phosphorylation at the initial stages of disease progression could represent a compensatory mechanism that mediates neuroprotection against hyperexcitability (Mondragón-Rodríguez et al., 2020). Taken together, adverse environmental outcomes during early-life which might affect A β PP metabolism and/or Tau protein might be prone to influence brain development and circuitry with long-lasting consequences.

EPIGENETIC MECHANISMS MEDIATE LONG-TERM INFLUENCE OF ENVIRONMENTAL FACTORS

Epigenetic inheritance has been defined as heritable changes in gene expression that are not encoded in the primary DNA sequence. Epigenetic processes can regulate DNA replication and repair, RNA transcription, and chromatin conformation, which influence in turn transcriptional regulation and protein translation. Epigenetic marks are partly inherited (transmission of epigenetic marks from one generation to another) but are also the signature and reflection of exposure to environments throughout life, particularly the early environment during gestation and lactation, which is a very sensitive period to both internal and external factors. Indeed, epigenetic mechanisms represent a means through which environmental factors can leave long-lasting memory of past experiences (Kundakovic and Jaric, 2017; Cirulli et al., 2020; Bellver-Sanchis et al., 2021; Breton et al., 2021). Although epigenetic modifications are often considered stable over time, epigenetic plasticity exists and may mediate adaptation to drastic environmental changes. Epigenetic mechanisms have broad actions but may mainly regulate 3 biological functions: DNA and RNA methylation, chromatin remodelling and expression of non-coding RNAs.

DNA and RNA Methylation

DNA methylation is one of the most studied and characterized epigenetic marks. It corresponds to the addition, *via* an enzymatic process, of a methyl (CH₃) group covalently linked to a cytosine

residue in position 5 (5-methylcytosine, 5mC). DNA methylation that occurs mainly on palindromic CpG sequences, which can form CpG islands, takes place during early development and does not increase over time (Jang et al., 2017). However, recent studies indicate that DNA methylation can also occur on non-CpG sequences (called CpH, where H is A, C or T), especially in the brain of vertebrates (de Mendoza et al., 2021). Methylation of CpH, unlike CpG, is established *de novo* during the post-natal period, which corresponds to the maturation processes of neurons, and is increased later in life (Guo et al., 2014; Stroud et al., 2017). DNA methylation on CpG usually leads to inhibition of gene expression when it occurs at the promoter region of genes, while it increases transcriptional activity when it occurs in the transcribed regions. Although many studies suggest that CpH methylation is inhibitory, this is still debated (Jang et al., 2017). The addition of the CH₃ group is catalysed by DNA methyltransferases (DNMTs). There are 3 DNMTs, DNMT3A and DNMT3B being involved in *de novo* methylation, while DNMT1 catalyses the maintenance of methylation during DNA replication. DNA methylation, although quite stable, is a reversible mechanism mediated by the action of Ten-eleven translocation (TET) enzymes. This family of 3 isoforms catalyses the oxidation of 5mC to hydroxymethylcytosine (5hmC), 5-formylcytosine (5fC) and 5-carboxylcytosine (5caC), the last two products can be excised from DNA through base excision repair, thus resulting in the reversion to demethylated cytosine (for review Lacal and Ventura, 2018). Interestingly, the activity of DNMTs and TETs is particularly high during neurodevelopment and the balance between these two families of enzymes is thought to play a crucial role in the dynamic regulation of brain genes involved in development and in cell differentiation (Tao et al., 2018; Cisternas et al., 2019). A recent study reported the methylation patterns of the Presenilin-1 gene (PSEN1), which encodes the catalytic peptide of the gamma-secretase complex (a key enzyme that cleaves the A β PP), in the brain of AD animals and humans. Interestingly, a temporal correlation between dynamic modifications in the PSEN1 CpG and non-CpG methylation patterns and mRNA expression during neurodevelopment and AD neurodegeneration has been reported (Monti et al., 2020). Moreover, several studies have shown that a decrease in TET and DNMT activity is associated with AD and cognitive impairment (Christian et al., 2020; Zhang et al., 2020; Antunes et al., 2021). Although it is likely that 5hmC represents the first step of active DNA demethylation, several data suggest that the 5hmC mark, which is enriched in adult CNS, is an active and independent epigenetic mark promoting the recruitment of factors allowing or not gene expression (Mellén et al., 2012; Shi et al., 2017; Stoyanova et al., 2021). In addition to DNA, methylation can also occur on mRNAs and lncRNAs. In this case, the addition of the methyl group takes place on adenine at position 6 (m6A) and is catalysed by a protein complex including METTL3 and METTL4 that carry the catalytic domain. This post-transcriptional modification of RNA promotes binding to other proteins, allowing pre-miR formation, translation or mRNA degradation. As with DNA methylation, this mechanism is reversible via the action of FTO and ALKBH5 (for review Zhou et al., 2020).

Chromatin Remodelling

In mammalian cells, histones interact with DNA to form chromatin. There are four types of histones: H2A, H2B, H3 and H4, which are present in duplicate to form an octamer, around which DNA is bound, forming a nucleosome. In the same way as DNA, histones can undergo several post-translational modifications (acetylation, methylation, phosphorylation, ubiquitination ...), altering the structure of chromatin. Histone methylation, via histone methyltransferase (HMT) activity, on the N-terminal tail changes the degree of chromatin compaction, making the DNA accessible or inaccessible to transcription factors, regulating gene expression. Histone methylation can enable or repress gene expression depending on the histones methylated and the amount of methylation. The mechanism is dynamic as histones can be demethylated by histone demethylases (HDMs). Histones can also be acetylated on the lysine residues of the N-terminal tail. The addition of an acetyl group is catalysed by histone acetyltransferase (HAT), while histone deacetylases (HDACs) remove acetyl groups. Acetylation of lysines opens up chromatin, making the chromatin structure transcriptionally active with a greater access of transcriptional activators. In contrast, deacetylation compacts chromatin, preventing the binding of transcription factors, thus inhibiting gene expression (for review Bannister and Kouzarides, 2011).

Non-coding RNAs

There are two types of non-coding RNAs (ncRNAs) classified according to their size, ncRNAs below 200 nt are defined as “short” (sRNAs) and those above this length as “long” (lncRNAs). ncRNAs are single-stranded and the result of post-transcriptional maturation. Among sRNAs, microRNAs (22–25 nt), the most studied, can assemble to a protein complex and associate by complementarity with RNA segments usually on the 3' untranslated region, leading to translational repression or mRNA degradation depending on the level of complementarity. The lncRNAs, whose functions are less studied, have genes similar to the genes that code for proteins and have different modes of action by linking to DNA, RNA and proteins. Non-exhaustively, they can inhibit transcription by preventing the recruitment of the pre-initiation complex, regulate other epigenetic modifications by guiding enzymes (HDACs, HATs, HMTs, DNMTs ...) to their site of action, activate the expression of neighbouring genes, bind to mRNA leading to mRNA degradation, bind to miRNAs limiting the action of the latter, regulate the splicing (for review Hombach and Kretz, 2016).

Although the emergence of Omics technologies has allowed to better characterize epigenetic modifications in the context of AD, these studies, that have been the subject of recent reviews (Monti et al., 2020; Coppedè, 2021; Nikolac Perkovic et al., 2021), are outside the scope of the present manuscript which is focused on the putative influence of the early environment on AD programming.

LONG-TERM CONSEQUENCES OF EARLY ENVIRONMENT ON TAU AND A β

Exposure to environmental agents has been extensively studied as a risk factor for the development of neurodegenerative diseases, such as AD (Killin et al., 2016; Wajman et al., 2018; Olayinka et al., 2019; Rahman et al., 2020). Recent studies suggest that exposure to cafeteria diets, pesticides, nanoparticles, air pollution and heavy metals during the perinatal period may sensitise to the development of AD in the long term, acting on the two lesions of the disease, Tau pathology and amyloid pathology, but also on neuroinflammation, which is considered as culprit component of the disease. Below, we will discuss the recent literature on the effects of perinatal exposure to different factors on Tau and A β .

Early-Life Exposure to Chemical Agents

Exposure to lead (Pb) during early-life has been particularly studied in the context of AD. Indeed, lead pollution is a major public health issue in developing cities. A meta-analysis using data on Pb exposure in Mexico City residents showed high blood Pb levels associated with a 5-point reduction in Intellectual quotient (IQ), and intellectual disability in children aged 0–4 years-old (Caravanos et al., 2014). Although no epidemiological study reports the effects of long-term Pb exposure, experimental studies in rodents (Bihagi et al., 2014; Gąssowska et al., 2016) and monkeys (Bihagi and Zawia, 2013) indicate that early exposure to Pb augments the phosphorylation of Tau, and is associated with an increase in the level of the cyclin-dependant kinase 5 (Cdk5), a major Tau kinase that is involved in abnormal phosphorylation of Tau in AD brain (Cruz and Tsai, 2004). In addition to these effects on Tau, studies also showed an increase in A β PP expression and plaque formation in the aging primate brain induced by early Pb exposure (from birth until 400 days of age). That was associated with a decrease in DNMT1 and DNMT3a activity and with a reduction in the acetylation and methylation of certain histones, resulting in a reprogramming of gene expression associated to neurotransmitter, growth factors and signal transduction pathways, suggesting an epigenetic effect (Wu et al., 2008; Bihagi et al., 2011).

The action of bisphenol A (BPA), an endocrine disruptor used to manufacture polycarbonate plastics, is also well documented. Interestingly BPA, due to its lipophilic nature, is able to penetrate the blood brain barrier and has been found in the placenta, amniotic fluid, blood breastmilk, suggesting that it could act on fetuses/neonates during the perinatal life (Hines et al., 2015). Indeed, epidemiological studies have associated BPA exposure during early life stages with low birth weight and an increased risk of developing metabolic diseases in adulthood (Chevalier and Fénelichel, 2015). In mice, maternal BPA exposure disrupts brain function and induces cognitive deficits (Tian et al., 2010) and learning-memory impairment in offspring (Xu et al., 2010). It also augments A β PP levels and Tau phosphorylation via increased activities of GSK3 β and CDK5—two major Tau kinases (Sergeant et al., 2008)—but decreases insulin signalling and alters synaptic plasticity in adulthood (Fang et al., 2016; Xue et al., 2020). Prenatal BPA exposure deregulates the offspring hippocampal

transcriptome with genes associated with AD, oxidative stress and inflammation (Sukjamnong et al., 2020). In particular, it induces elevation of NF- κ B protein and its AD-related target BACE1, a key enzyme involved in A β PP processing, suggesting that prenatal BPA exposure triggers neuroinflammation in the hippocampus and increases the susceptibility of AD through NF- κ B. These modifications may explain the cognitive impairments observed in other studies (Negishi et al., 2004; Ryan and Vandenberg, 2006; Wang et al., 2014), even with a single dose of BPA at PND10 (Viberg et al., 2011). Indeed, *in utero* and neonatal exposure to low dose of BPA may cause deregulation of miRNAs expression, DNA methylation levels and histone modifications (Singh and Li, 2012). A genome-wide analysis of the foetal mouse forebrain epigenome shows that exposure to low doses of BPA modifies the methylation in the promoter-associated CpG islands at several loci, which can alter the brain development (Yaoi et al., 2008). In rat, the exposure to the Di-(2-thyhexyl)-phtalate (DEHP), another endocrine disruptor, during gestation and lactation also increases the phosphorylation of Tau, without modification of A β PP level in hippocampus, and is associated with cognitive impairments in adulthood (Sun, 2014). However, so far no studies have investigated the long-term consequences of early-life BPA on epigenetic changes in the Microtubule Associated Protein Tau (MAPT, the gene encoding Tau) and A β PP genes.

Exposure to xenobiotics, man-made chemicals found in the environment but not endogenously produced, is increasingly common, and their presence in the brain during the perinatal period can alter CNS development. Polybrominated diphenyl ethers 209 (PBDE 209), a combustion inhibitor that spreads in the environment, has been found in breast milk and in consumer products. Studies in animals show that a single exposure to PBDE 209 at PND10, at the time of brain growth spurt in rodent, does not modify the level of Tau but affects CaMKII and synaptophysin levels in the hippocampus (Viberg, 2009), two proteins playing an essential role in neurodevelopment, whose alterations may likely explain impaired spontaneous behaviour seen in adulthood (Johansson et al., 2008). Similarly, exposure to another xenobiotic, the non-dioxin-like polychlorinated biphenyls (NDL-PCB), during lactation in Swiss albino mice led to a more pronounced decrease of the synaptic proteins synaptophysin and PSD95 and impairment of long-term memory when the brain is challenged by the injection of A β oligomers (Elnar et al., 2016).

Early-Life Exposure to High-Fat Diet

Maternal obesity has considerably increased among women of reproductive age in the last decades. Current estimates suggest that by 2025 more than 21% of women in the world will be obese, and in 2014, the estimated percentage of overweight and obesity among pregnant women was 33% in the United States (Poston et al., 2016). Cumulative data suggest that maternal obesity leads to metabolic and cognitive disorders in the offspring (Hasebe et al., 2021), which makes maternal obesity a major public health issue. However, although cognitive and metabolic impairments are two components of AD, the consequences of maternal perinatal high-fat diet on Tau and A β have been poorly

investigated. A recent study by Di Meco *et al.* reports that a maternal high-fat diet (mHFD with 42% calories from fat) during gestation decreases total Tau and pathological Tau conformation level and increases the synaptic integrity, leading to improved cognitive performance in adult mice (Di Meco and Praticò, 2019). However, it has also been reported that a mHFD (45% fat) during gestation and lactation increases the level of A β peptides at the vascular level, and is associated with a modification of the neurovascular environment in the brain of adult mice, which leads to the alteration of the perivascular clearance of A β peptides (Hawkes et al., 2015). Although many works have looked at the epigenetic changes induced by mHFD in the programming of peripheral metabolic diseases, very few studies have looked at the consequences in the CNS of the offspring (Gawlińska et al., 2021a). For instance, it has been reported that consumption of a high-fat diet during gestation and lactation in the Wistar rat decreases the global level of histone H4 acetylation in the hippocampus at P50, leading to transcriptional repression (Gonçalves et al., 2017). Interestingly, a diet rich in grape juice has an opposite effect on the level of this epigenetic mark at PND21 and is associated with a beneficial effect on the brain with an antioxidant effect (Gonçalves et al., 2017; Schaffer et al., 2019) which could limit deleterious consequences of mHFD by reducing oxidative stress and acetylcholinesterase activity (Proença et al., 2021). However, to our knowledge, the putative consequences of mHFD on epigenetic changes in MAPT and A β PP genes have never been reported.

Early-Life Exposure to Tobacco

According to the WHO, more than 1.3 billion people worldwide smoke, about 40% of them being women. In low- and middle-income countries, where the proportion of people who smoke is highest, the prevalence of pregnant women who smoke is 2.6%, but can be as high as 30.8% in Russia (Caleyachetty et al., 2014). The deleterious effects of smoking on offspring are widely accepted and can lead to IUGR and neurodevelopmental changes (Perera et al., 2005; El Marroun et al., 2016). An epidemiological study on 471 individuals in Finland showed that male offspring of mothers who were exposed to cigarette smoke prenatally had more cognitive problems, although these effects were minor or absent in females (Ramsay et al., 2016). In mice, it has been shown that exposure to cigarette smoke during lactation increases the phosphorylation of Tau, and the levels of the 3R form of brain Tau, leading to a deregulation of the 3R/4R ratio of Tau during a critical period of brain development in 4-week-old offspring (La Maestra et al., 2011). Indeed, Tau phosphorylation but also the proportion of each isoform plays an important role in brain development, and particularly during lactation, during which synaptogenesis and axonal growth take place. Although not reported to date, these early changes could lead to neuronal dysfunction and exert long-term consequences.

Early-Life Exposure to Ionizing Radiation

Humans are routinely exposed to ionizing radiation (IR) from natural sources, and from man-made sources such as nuclear energy. IR is also increasingly used in the medical field in the diagnosis or treatment of diseases, for example in X-ray imaging. For example, in children, for the treatment of

brain tumours, exposure at a very young age can also influence the development of CNS. An epidemiological study on 3,094 Swedish males indicated that low-dose IR exposure during early human development can have a negative impact on cognitive abilities during childhood (Hall et al., 2004). An experimental study in mice reported that a single low-dose exposure of IR to PND10 animals deregulates the hippocampal and cortical proteome and transcriptome, impairs synaptic plasticity, reduces neurogenesis and leads to neuroinflammation later in life. Interestingly, miRNAs analysis showed an increased expression of miR-132 and miR-134 repressing the translation of proteins of the Rac1-Cofilin signalling pathway involved in synaptic plasticity and neurogenesis, highlighting a potential epigenetic mechanism (Kempf et al., 2014). Another team using a similar protocol demonstrated an increase in cortical Tau levels at PND11 (i.e. 1 day after exposure) and at 6 months. That was associated with a decline of long-term spatial memory when IR is combined to ketamine exposure, an anaesthesia commonly used to facilitate radiotherapy in children. Although it has not been directly assessed, these long-term consequences on Tau protein levels and cognitive deficits suggested the involvement of epigenetic mechanisms (Buratovic et al., 2014; Buratovic et al., 2018).

Early-Life Exposure to Asphyxia

Perinatal asphyxia is a defect in blood flow or gas exchange from the mother to the foetus or vice versa, occurring immediately before, during or after birth. It accounts for up to 20 cases per 1,000 births in developing countries, with mortality during the neonatal period of 15–20%, while 25% of survivors develop short- and long-term neurological disorders, associated with neonatal encephalopathy in most cases (Gillam-Krakauer and Gowen Jr, 2021). The most common diagnosis of perinatal asphyxia is addressed by chest X-ray, which, as mentioned in the previous paragraph, can be deleterious, especially in the young individual. Thus, it is important to uncover biomarkers released into the cerebrospinal fluid (CSF) during perinatal asphyxia to avoid X-ray exposure. With this idea in mind, one team evaluated the levels of Tau, pTau and A β 42 in the CSF of newborn pigs that had undergone a perinatal hypoxia event a few hours after birth. They showed a decrease in the level of A β 42, with no change in the level of Tau and pTau. They also observed an increase in S100B (Benterud et al., 2015), a glial marker of neuroinflammation which is also increased in CSF of infants with perinatal asphyxia (Massaro et al., 2014). In contrast, an epidemiological study reported that delivery by caesarean, known to limit hypoxia in cases of perinatal asphyxia, was associated with a reduction in cord blood Tau protein levels, indicating that Tau is modulated by blood oxygen concentration (Tunc et al., 2010) although it could not be excluded that Tau variation results from other factors associated with caesarean delivery. However, it is clear that perinatal asphyxia modified Tau, pTau and A β 42 in the fetus/neonate, it remains to be determined if these early-life changes influence (or not) the development of late-onset AD.

LONG-TERM CONSEQUENCES OF EARLY ENVIRONMENT IN ANIMAL MODELS OF AD

Although early-life stressors exposure affects Tau and A β peptides levels, very few studies investigated the effects of perinatal perturbations in the pathological context of AD. This section aims at summarizing the literature on the consequences of early environment on the development of AD lesions.

Early-Life Exposure to Maternal High-Fat Diet

As previously mentioned, mHFD can have long-term deleterious effects on the levels of Tau and A β peptides. The consequences of mHFD have also been studied in the 3xTgAD mouse, a model of AD that develops early amyloid pathology and later Tau pathology. A first team reported that the application of mHFD (60% fat) during gestation and lactation impaired short- and long-term spatial memory and increased the number of Tau-positive neurons in the hippocampus without altering A β peptides level, suggesting that mHFD exacerbated Tau pathology and may sensitize to the development of AD (Martin et al., 2014). Surprisingly, using the same animal model it has also been reported that mHFD (42% fat) only applied during gestation protects from synaptic dysfunctions and associated cognitive disorders and decreases Tau and amyloid pathology by reducing the amount A β 40 and A β 42. This putative protective effect of the maternal diet could result from an increased gene expression of the transcriptional repressor FOXP2, which could be responsible for the decreased gene expression of Tau, CDK5 (a Tau kinase) and BACE-1 (involved in the amyloidogenic pathway) possibly via epigenetic mechanisms (Di Meco et al., 2019). It is important to note that an important limitation of this model is that the mice are homozygous for all 3 mutant alleles, and thus the observed phenotype is the result of the diet given to transgenic dams. Therefore, it is not entirely clear if changes result from the diet effect only or from the convergent impact of the diet and the transgenic phenotype of dams. Using the Tg2576 mouse, another model of AD expressing the human Swedish A β PP mutation and developing only amyloid pathology, it has been reported that a mHFD (45% fat) starting before crossing until weaning increases the level of soluble A β and the number of amyloid plaques. It has been suggested that these alterations could modify the extracellular matrix leading to increase A β plaques retention within the parenchyma (Nizari et al., 2016). These seemingly contradictory results may be explained by the different amount or types of fat used in the diets and/or by the timing of exposure of mHFD, as well as by the model used regarding amyloid pathology. Although the biological mechanisms that explain the consequences of mHFD on the development of AD are still poorly understood, recent studies suggest an important role for the gut microbiota. Indeed changes in maternal diet during the perinatal period modify both the mother's and the offspring's microbiota with maternal transmission both at delivery and through milk intake (Mueller et al., 2015). In mice, perinatal mHFD alters the short-term memory and is

associated with a deregulation of the gut microbiota in offspring, which could partly explain the observed phenotype (Sanguinetti et al., 2019). Interestingly, a high-fibre diet in mothers or offspring can restore cognitive deficits in offspring by regulating the composition of bacteria and SCFA metabolites in the long-term (Liu et al., 2021). It has been suggested that early deregulation of the microbiota may participate to cognitive impairment and to the onset of later lesions in the 3xTgAD mouse model (Bello-Medina et al., 2021). Interestingly, addition of bioactive food to the diet restores these effects and modulates the composition of the gut microbiota (Syeda et al., 2018). Together, these studies indicate that the modification of the diets or the addition of probiotics, both of which being able to modulate epigenetic mechanisms, offer the opportunity for identification of new diagnostic biomarkers and treatments (Ji and Shen, 2021).

Early-Life Exposure to Caffeine

Tea and coffee are two commonly substances consumed in the world, whose main component is caffeine, a psychoactive substance. Numerous epidemiological studies in humans and experimental studies in AD mouse models have shown the beneficial and long-term protective effects of moderate caffeine consumption in adulthood in reducing the development of lesions and related to cognitive impairment, decreasing the risk of developing AD by 65% in humans (Eskelinen et al., 2009; Flaten et al., 2014; Polito et al., 2018). Sustaining epidemiological studies, caffeine treatment of adult mice developing amyloid or Tau demonstrate positive outcomes at the pathological and behavioural levels (Arendash et al., 2006; Arendash et al., 2009; Laurent et al., 2014). In sharp contrast perinatal caffeine, by blocking adenosine A2A receptors, not only exert a detrimental impact on neuronal migration but caffeine receptors themselves strongly impact GABAergic synapse stabilization during synaptogenesis (Gomez-Castro et al., 2021). Further, perinatal caffeine strongly impacts hippocampal excitability and memory at the adult stage. Whether perinatal caffeine modulates the later development of AD pathology or has cognitive consequences remains largely unclear. However, recent data showed that perinatal caffeine consumption starting before gestation until PND15 in a transgenic mouse model of tauopathy, even not modifying the pathological load, leads to earlier long-term memory impairments together with excitatory/inhibitory synaptic drive in adulthood. These results indicate that early-life caffeine may influence brain development and exert long-lasting consequences in a Tau background (Zappettini et al., 2019).

Early-Life Exposure to Stress


Evidence indicates that chronic stress can participate to the progression of AD and other neurodegenerative disorders. In 3xTgAD mice, chronic social stress in adulthood, by regularly changing the composition of the mice in the cage, decreases BDNF levels and increases amyloid pathology and anxiety, with no change in control animals indicating a greater sensitivity to stress in mutant mice (Rothman et al., 2012). During the perinatal period, although few studies have been reported, Sierksma et al.

have shown that chronic restraint stress performed during gestation (E1-E7) in APPswe/PS1dE9 mice model of AD induced sexual-dimorphic results in the adult offspring with males exhibiting altered spatial memory, while females have improved spatial memory associated with decreased amyloid load in the dorsal hippocampus (Sierksma et al., 2012). Although these modifications were not associated with changes in DNMT expression nor 5mC and 5hmC levels, other epigenetic mechanisms targeting histones or non-coding RNAs remain to be studied. Using the same AD animal model, it has been shown that repeated maternal separation during lactation, which is also used to induce stress during early-life, increases amyloid deposits in the hippocampus and cortex, and impairs learning and spatial memory in adulthood (Hui et al., 2017). The same experimental protocol increases Tau and amyloid pathology, decreases synaptophysin and PSD95 (two synaptic markers), impairs memory and is associated with hyperactivity of the hypothalamo-pituitary-adrenal (HPA) axis in adult male rats (Martisova et al., 2013). These observations are in agreement with the elevated glucocorticoid levels in adulthood found associated with a higher risk of developing AD, by altering glial function, Tau and amyloid lesions, and synaptic plasticity (Vyas et al., 2016). The first evidence that early-life stress could induce epigenetic programming in the rat brain came from the pioneer studies of Weaver et al. who showed that poor maternal care leads to increased methylation in the glucocorticoid receptor (GR) promoter which was accompanied by aberrant behaviours and altered HPA responses in adult offspring (Weaver et al., 2004). Using the 3xTgAD mice, it has been reported that handling offspring during lactation (PND2-PND9), by separating the dam from her pups for 15 min, increases maternal care of the dam towards her pups upon reunion. Increased maternal care leads to a decrease both in short-term memory decline and in amyloid load in the hippocampus, suggesting that prenatal stress could be mitigated or to have antagonistic effect by postnatal maternal care.

Recently, it has been reported in a subclinical AD-like rat model, that adult animals which were prenatally-stressed (PS) exhibited increased dysfunctions of HPA axis activity and working and long-term spatial memory as well as modified cellular organisation in CA1 hippocampal region. Interestingly, these alterations were alleviated by increased postnatal maternal care during lactation. Unfortunately, the authors did not report the putative consequences of cross-fostering in adult PS rats on A β plaques and NFT nor on phosphorylation and protein levels of Tau (Rostami et al., 2020).

Conclusion and Perspectives

In the present review, we provided strong evidence that early-life stress may constitute another risk factor contributing to the development of late-onset AD possibly *via* long-term influence of epigenetic mechanisms (Figure 1). However, important questions, which are still unanswered in the context of AD, such as the putative contribution of transgenerational epigenetic inheritance, paternal contribution and sexual dimorphism on late-onset AD development, have been overlooked. It should be kept in mind that although epigenetic



marks may be transmitted transgenerationally (Burton and Greer, 2021), the existence of multigenerational epigenetic inheritance in humans is still a matter of debate (Cavalli and Heard, 2019). Epigenetic mechanisms are indeed plastic, modifiable and reversible processes providing thus opportunities to try to reverse deleterious effects of environment *via* early-life interventional strategies (Lesuis et al., 2018). Recently, the consequences of mHFD (60% fat) have been examined in F1 (intergenerational inheritance) and F2 (transgenerational inheritance) generations of SAMP8 mice, a model of accelerated ageing leading to increased cognitive decline associated with the development of neuropathological AD hallmarks (Liu et al., 2020). The HFD applied to female mice of the F0 generation was supplemented or not with resveratrol, a natural phytoalexin known to have beneficial effects in

pathological conditions, notably via its anti-inflammatory effects, and by reducing amyloid pathology and playing a role in epigenetic mechanisms (Rahman et al., 2020). Interestingly, the impairment of both short- and long-term memory and metabolic parameters (increased plasma levels of triglycerides and leptin) induced by the HFD was attenuated by resveratrol supplementation in the F0, F1 and F2 generations supporting a multigenerational effect. In addition, the level of DNA and RNA methylation (on 5mC and m6A, respectively) is increased in F1 offspring and associated with a change in the expression of enzymes involved in these epigenetic mechanisms, strongly suggesting epigenetic inheritance (Izquierdo et al., 2021). Future studies will be clearly required to understand if and how early nutritional interventions in mothers may exert beneficial effects in offspring and mitigate the long-term

deleterious effects of adverse environmental exposure on late-onset AD.

Although most of the studies focused on the consequences of maternal environmental stress in the offspring, a growing body of evidence indicates that paternal preconception stress may also have long-term effects on the offspring. Studies in rodents indicate that paternal obesity induced by HFD alters the development of the 2-cell embryo. These early events have long-lasting consequences and decrease neurogenesis and memory, and are associated with epigenetic marks changes in adulthood, including hypermethylation of BDNF promoter, a neurotrophic factor that plays a key role in brain development and function (McPherson et al., 2015; Zhou et al., 2018). It has also been reported that repeated paternal stress (Harker et al., 2015, 2018), as well as paternal BPA exposure (Luo et al., 2017), lead to behavioural alterations, associated with changes in dendritic spines and brain organisation in the offspring. In addition, paternal stress during spermatogenesis leads to altered behaviour and DNA methylation patterns with increased level in the hippocampus of both male and female PND21 offspring (Mychasiuk et al., 2013). In mice, paternal cognitive spatial training improves cognitive performance in adult offspring and is associated with an increase in the hippocampal expression of synaptotagmin, a pre-synaptic protein. This was associated with elevation of histone acetylation at the promoter of synaptotagmin in the hippocampi of both fathers and offspring as well as in fathers sperm (Zhang et al., 2017). Thus, it would be interesting to delineate the role of paternal stress in animal models of AD and to determine whether both maternal and paternal exposure to stress would have a cumulative effect. Although it would more closely mimic what happens in humans, to our knowledge, this has never been reported in the context of AD.

REFERENCES

- Antunes, C., Da Silva, J. D., Guerra-Gomes, S., Alves, N. D., Ferreira, F., Loureiro-Campos, E., et al. (2021). Tet3 Ablation in Adult Brain Neurons Increases Anxiety-like Behavior and Regulates Cognitive Function in Mice. *Mol. Psychiatry* 26, 1445–1457. doi:10.1038/s41380-020-0695-7
- Arendash, G. W., Mori, T., Cao, C., Mamcarz, M., Runfeldt, M., Dickson, A., et al. (2009). Caffeine Reverses Cognitive Impairment and Decreases Brain Amyloid-Beta Levels in Aged Alzheimer's Disease Mice. *J. Alzheimers Dis.* 17, 661–680. doi:10.3233/JAD-2009-1087
- Arendash, G. W., Schleif, W., Rezai-Zadeh, K., Jackson, E. K., Zacharia, L. C., Cracchiolo, J. R., et al. (2006). Caffeine Protects Alzheimer's Mice Against Cognitive Impairment and Reduces Brain Beta-Amyloid Production. *Neuroscience* 142, 941–952. doi:10.1016/j.neuroscience.2006.07.021
- Armstrong, A. R. (2019). Risk Factors for Alzheimer's Disease. *fn* 57, 87–105. doi:10.5114/fn.2019.85929
- Athanasopoulos, D., Karagiannis, G., and Tsolaki, M. (2016). Recent Findings in Alzheimer Disease and Nutrition Focusing on Epigenetics. *Adv. Nutr.* 7, 917–927. doi:10.3945/an.116.012229
- Bannister, A. J., and Kouzarides, T. (2011). Regulation of Chromatin by Histone Modifications. *Cell Res* 21, 381–395. doi:10.1038/cr.2011.22
- Barker, D. J. P. (2007). The Origins of the Developmental Origins Theory. *J. Intern. Med.* 261, 412–417. doi:10.1111/j.1365-2796.2007.01809.x
- Bello-Medina, P. C., Hernández-Quiroz, F., Pérez-Morales, M., González-Franco, D. A., Cruz-Pauseno, G., García-Mena, J., et al. (2021). Spatial Memory and Gut Microbiota Alterations Are Already Present in Early Adulthood in a Pre-clinical Transgenic Model of Alzheimer's Disease. *Front. Neurosci.* 15, 595583. doi:10.3389/fnins.2021.595583
- Bellver-Sanchis, A., Pallàs, M., and Griñán-Ferré, C. (2021). The Contribution of Epigenetic Inheritance Processes on Age-Related Cognitive Decline and Alzheimer's Disease. *Epigenomes* 5, 15. doi:10.3390/epigenomes5020015
- Benterud, T., Pankratov, L., Solberg, R., Bolstad, N., Skinningsrud, A., Baumbusch, L., et al. (2015). Perinatal Asphyxia May Influence the Level of Beta-Amyloid (1-42) in Cerebrospinal Fluid: An Experimental Study on Newborn Pigs. *PLoS One* 10, e0140966. doi:10.1371/journal.pone.0140966
- Bihagi, S. W., Bahmani, A., Adem, A., and Zawia, N. H. (2014). Infantile Postnatal Exposure to lead (Pb) Enhances Tau Expression in the Cerebral Cortex of Aged Mice: Relevance to AD. *Neurotoxicology* 44, 114–120. doi:10.1016/j.neuro.2014.06.008
- Bihagi, S. W., Huang, H., Wu, J., and Zawia, N. H. (2011). Infant Exposure to Lead (Pb) and Epigenetic Modifications in the Aging Primate Brain: Implications for Alzheimer's Disease. *Jad* 27, 819–833. doi:10.3233/JAD-2011-111013
- Bihagi, S. W., and Zawia, N. H. (2013). Enhanced Tauopathy and AD-like Pathology in Aged Primate Brains Decades after Infantile Exposure to lead (Pb). *Neurotoxicology* 39, 95–101. doi:10.1016/j.neuro.2013.07.010
- Biswas, S., and Kalil, K. (2018). The Microtubule-Associated Protein Tau Mediates the Organization of Microtubules and Their Dynamic Exploration of Actin-Rich Lamellipodia and Filopodia of Cortical Growth Cones. *J. Neurosci.* 38, 291–307. doi:10.1523/JNEUROSCI.2281-17.2017
- Bomfim, T. R., Forny-Germano, L., Sathler, L. B., Brito-Moreira, J., Houzel, J.-C., Decker, H., et al. (2012). An Anti-diabetes Agent Protects the Mouse Brain from Defective Insulin Signaling Caused by Alzheimer's Disease-

As previously mentioned, a large majority of works have examined the consequences of maternal stress on male offspring. To date and to the best of our knowledge, only three studies look at the consequences of early-life environment on females, even though Alzheimer's disease affects women preferentially. In the US, nearly 70% of AD patients are women, which can be explained by a longer life expectancy, but also by intrinsic biological differences, including brain organisation and function, age-associated loss of protection exerted by steroid hormones, higher inflammatory response, but also different susceptibility to genetic variations (for review Fisher et al., 2018). In addition, during early-development brain volume is higher in males, while the hippocampus is larger in females, pointing out to sexual dimorphism during a critical time-window (Kaczurkin et al., 2019). Finally, although few studies have been reported in the context of AD, the use of mouse models led to discordant results in males and females showing the importance of studying the effects in the two sexes independently.

In conclusion, although the way by which perinatal epigenetic mechanisms act to program long-term memory and cognition remains to be clarified, it seems clear that the development of multi-omics strategies will allow molecular dissection of this fascinating new field of research. In particular, it will offer novel opportunities to identify early markers of AD and to develop novel preventive strategies to attenuate age-associated epigenetic alterations.

AUTHOR CONTRIBUTIONS

All authors listed have made a substantial, direct, and intellectual contribution to the work and approved it for publication.

- Associated A β Oligomers. *J. Clin. Invest.* 122, 1339–1353. doi:10.1172/JCI57256
- Breton, C. V., Landon, R., Kahn, L. G., Enlow, M. B., Peterson, A. K., Bastain, T., et al. (2021). Exploring the Evidence for Epigenetic Regulation of Environmental Influences on Child Health across Generations. *Commun. Biol.* 4, 1–15. doi:10.1038/s42003-021-02316-6
- Bucht, G., Adolfsson, R., Lithner, F., and Winblad, B. (1983). Changes in Blood Glucose and Insulin Secretion in Patients with Senile Dementia of Alzheimer Type. *Acta Med. Scand.* 213, 387–392. doi:10.1111/j.0954-6820.1983.tb03756.x
- Buratovic, S., Stenerlöv, B., Fredriksson, A., Sundell-Bergman, S., Viberg, H., and Eriksson, P. (2014). Neonatal Exposure to a Moderate Dose of Ionizing Radiation Causes Behavioural Defects and Altered Levels of Tau Protein in Mice. *Neurotoxicology* 45, 48–55. doi:10.1016/j.neuro.2014.09.002
- Buratovic, S., Stenerlöv, B., Sundell-Bergman, S., Fredriksson, A., Viberg, H., Gordh, T., et al. (2018). Effects on Adult Cognitive Function after Neonatal Exposure to Clinically Relevant Doses of Ionising Radiation and Ketamine in Mice. *Br. J. Anaesth.* 120, 546–554. doi:10.1016/j.bja.2017.11.099
- Burnouf, S., Martire, A., Derisbourg, M., Laurent, C., Belarbi, K., Leboucher, A., et al. (2013). NMDA Receptor Dysfunction Contributes to Impaired Brain-Derived Neurotrophic Factor-Induced Facilitation of Hippocampal Synaptic Transmission in a Tau Transgenic Model. *Aging Cell* 12, 11–23. doi:10.1111/acel.12018
- Burton, N. O., and Greer, E. L. (2021). Multigenerational Epigenetic Inheritance: Transmitting Information across Generations. *Semin. Cell Develop. Biol.* S1084–9521 (21), 00210–X. doi:10.1016/j.semcdb.2021.08.006
- Caleyachetty, R., Tait, C. A., Kengne, A. P., Corvalan, C., Uauy, R., and Echouffo-Tcheugui, J. B. (2014). Tobacco Use in Pregnant Women: Analysis of Data from Demographic and Health Surveys from 54 Low-Income and Middle-Income Countries. *Lancet Glob. Health* 2, e513–e520. doi:10.1016/S2214-109X(14)70283-9
- Caravanas, J., Dowling, R., María Téllez-Rojo Dra, M., Cantoral, A., Kobrosly, R., Estrada, D., et al. (2014). Blood lead Levels in Mexico and Pediatric burden of Disease Implications. *Ann. Glob. Health* 80, 269–277. doi:10.1016/j.aogh.2014.08.002
- Cavalli, G., and Heard, E. (2019). Advances in Epigenetics Link Genetics to the Environment and Disease. *Nature* 571, 489–499. doi:10.1038/s41586-019-1411-0
- Checler, F. (1995). Processing of the β -Amyloid Precursor Protein and its Regulation in Alzheimer's Disease. *J. Neurochem.* 65, 1431–1444. doi:10.1046/j.1471-4159.1995.65041431.x
- Chevalier, N., and Fénichel, P. (2015). Bisphenol A: Targeting Metabolic Tissues. *Rev. Endocr. Metab. Disord.* 16, 299–309. doi:10.1007/s11154-016-9333-8
- Christian, D. L., Wu, D. Y., Martin, J. R., Moore, J. R., Liu, Y. R., Clemens, A. W., et al. (2020). DNMT3A Haploinsufficiency Results in Behavioral Deficits and Global Epigenomic Dysregulation Shared across Neurodevelopmental Disorders. *Cel Rep.* 33, 108416. doi:10.1016/j.celrep.2020.108416
- Cirulli, F., Musillo, C., and Berry, A. (2020). Maternal Obesity as a Risk Factor for Brain Development and Mental Health in the Offspring. *Neuroscience* 447, 122–135. doi:10.1016/j.neuroscience.2020.01.023
- Cisneros-Franco, J. M., Voss, P., Thomas, M. E., and de Villers-Sidani, E. (2020). Critical Periods of Brain Development. *Handb Clin. Neurol.* 173, 75–88. doi:10.1016/B978-0-444-64150-2.00009-5
- Cisternas, C. D., Cortes, L. R., Bruggeman, E. C., Yao, B., and Forger, N. G. (2019). Developmental Changes and Sex Differences in DNA Methylation and Demethylation in Hypothalamic Regions of the Mouse Brain. *Epigenetics* 15, 72–84. doi:10.1080/15592294.2019.1649528
- Colin, M., Dujardin, S., Schraen-Maschke, S., Meno-Tetang, G., Duyckaerts, C., Courade, J.-P., et al. (2020). From the Prion-like Propagation Hypothesis to Therapeutic Strategies of Anti-tau Immunotherapy. *Acta Neuropathol.* 139, 3–25. doi:10.1007/s00401-019-02087-9
- Cooper, J. A. (2013). Mechanisms of Cell Migration in the Nervous System. *J. Cell Biol* 202, 725–734. doi:10.1083/jcb.201305021
- Coppède, F. (2021). Epigenetic Regulation in Alzheimer's Disease: Is it a Potential Therapeutic Target. *Expert Opin. Ther. Targets* 25, 283–298. doi:10.1080/14728222.2021.1916469
- Craft, S., Peskind, E., Schwartz, M. W., Schellenberg, G. D., Raskind, M., and Porte, D. (1998). Cerebrospinal Fluid and Plasma Insulin Levels in Alzheimer's Disease. *Neurology* 50, 164–168. doi:10.1212/wnl.50.1.164
- Craft, S., Zallen, G., and Baker, L. D. (1992). Glucose and Memory in Mild Senile Dementia of the Alzheimer Type. *J. Clin. Exp. Neuropsychol.* 14, 253–267. doi:10.1080/01688639208402827
- Cruz, J. C., and Tsai, L.-H. (2004). Cdk5 Deregulation in the Pathogenesis of Alzheimer's Disease. *Trends Mol. Med.* 10, 452–458. doi:10.1016/j.molmed.2004.07.001
- Dawson, H. N., Ferreira, A., Eyster, M. V., Ghoshal, N., Binder, L. I., and Vitek, M. P. (2001). Inhibition of Neuronal Maturation in Primary Hippocampal Neurons from τ Deficient Mice. *J. Cell Sci* 114, 1179–1187. doi:10.1242/jcs.114.6.1179
- de Graaf-Peters, V. B., and Hadders-Algra, M. (2006). Ontogeny of the Human central Nervous System: what Is Happening when. *Early Hum. Develop.* 82, 257–266. doi:10.1016/j.earlhumdev.2005.10.013
- de la Monte, S. M., Tong, M., and Wands, J. R. (2018). The 20-Year Voyage Aboard the Journal of Alzheimer's Disease: Docking at 'Type 3 Diabetes', Environmental/Exposure Factors, Pathogenic Mechanisms, and Potential Treatments. *Jad* 62, 1381–1390. doi:10.3233/JAD-170829
- de la Monte, S. M. (2014). Type 3 Diabetes Is Sporadic Alzheimer's Disease: Mini-Review. *Eur. Neuropsychopharmacol.* 24, 1954–1960. doi:10.1016/j.euroneuro.2014.06.008
- de Mendoza, A., Poppe, D., Buckberry, S., Pflueger, J., Albertin, C. B., Daish, T., et al. (2021). The Emergence of the Brain Non-CpG Methylation System in Vertebrates. *Nat. Ecol. Evol.* 5, 369–378. doi:10.1038/s41559-020-01371-2
- Di Meco, A., Jelinek, J., Lauretti, E., Curtis, M. E., Issa, J.-P. J., and Praticò, D. (2019). Gestational High Fat Diet Protects 3xTg Offspring from Memory Impairments, Synaptic Dysfunction, and Brain Pathology. *Mol. Psychiatry* 26, 7006–7019. doi:10.1038/s41380-019-0489-y
- Di Meco, A., and Praticò, D. (2019). Early-life Exposure to High-fat Diet Influences Brain Health in Aging Mice. *Aging Cell* 18, e13040. doi:10.1111/acel.13040
- Dityatev, A., Seidenbecher, C. I., and Schachner, M. (2010). Compartmentalization from the outside: the Extracellular Matrix and Functional Microdomains in the Brain. *Trends Neurosciences* 33, 503–512. doi:10.1016/j.tins.2010.08.003
- El Marroun, H., Tiemeier, H., Franken, I. H. A., Jaddoe, V. W. V., van der Lugt, A., Verhulst, F. C., et al. (2016). Prenatal Cannabis and Tobacco Exposure in Relation to Brain Morphology: A Prospective Neuroimaging Study in Young Children. *Biol. Psychiatry* 79, 971–979. doi:10.1016/j.biopsych.2015.08.024
- Elnar, A. A., Allouche, A., Desor, F., Yen, F. T., Soulimani, R., and Oster, T. (2016). Lactational Exposure of Mice to Low Levels of Non-dioxin-like Polychlorinated Biphenyls Increases Susceptibility to Neuronal Stress at a Mature Age. *Neurotoxicology* 53, 314–320. doi:10.1016/j.neuro.2015.10.003
- Eskelinen, M. H., Ngandu, T., Tuomilehto, J., Soininen, H., and Kivipelto, M. (2009). Midlife Coffee and tea Drinking and the Risk of Late-Life Dementia: a Population-Based CAIDE Study. *Jad* 16, 85–91. doi:10.3233/JAD-2009-0920
- Fang, F., Gao, Y., Wang, T., Chen, D., Liu, J., Qian, W., et al. (2016). Insulin Signaling Disruption in Male Mice Due to Perinatal Bisphenol A Exposure: Role of Insulin Signaling in the Brain. *Toxicol. Lett.* 245, 59–67. doi:10.1016/j.toxlet.2016.01.007
- Fisher, D. W., Bennett, D. A., and Dong, H. (2018). Sexual Dimorphism in Predisposition to Alzheimer's Disease. *Neurobiol. Aging* 70, 308–324. doi:10.1016/j.neurobiolaging.2018.04.004
- Flaten, V., Laurent, C., Coelho, J. E., Sandau, U., Batalha, V. L., Burnouf, S., et al. (2014). From Epidemiology to Pathophysiology: what about Caffeine in Alzheimer's Disease. *Biochem. Soc. Trans.* 42, 587–592. doi:10.1042/BST20130229
- Flores-Dorantes, M. T., Díaz-López, Y. E., and Gutiérrez-Aguilar, R. (2020). Environment and Gene Association with Obesity and Their Impact on Neurodegenerative and Neurodevelopmental Diseases. *Front. Neurosci.* 14, 863. doi:10.3389/fnins.2020.00863
- Gapp, K., Woldemichael, B. T., Bohacek, J., and Mansuy, I. M. (2014). Epigenetic Regulation in Neurodevelopment and Neurodegenerative Diseases. *Neuroscience* 264, 99–111. doi:10.1016/j.neuroscience.2012.11.040
- Gąssowska, M., Baranowska-Bosiacka, I., Moczyłowska, J., Tarnowski, M., Pilutin, A., Gutowska, I., et al. (2016). Perinatal Exposure to lead (Pb) Promotes Tau Phosphorylation in the Rat Brain in a GSK-3 β and CDK5 Dependent Manner: Relevance to Neurological Disorders. *Toxicology* 347–349, 17–28. doi:10.1016/j.tox.2016.03.002
- Gawlińska, K., Gawliński, D., Borczyk, M., Korostyński, M., Przeglasiński, E., and Filip, M. (2021a). A Maternal High-Fat Diet during Early Development

- Provokes Molecular Changes Related to Autism Spectrum Disorder in the Rat Offspring Brain. *Nutrients* 13, 3212. doi:10.3390/nu13093212
- Gawlińska, K., Gawliński, D., Filip, M., and Przeglasiński, E. (2021b). Relationship of Maternal High-Fat Diet during Pregnancy and Lactation to Offspring Health. *Nutr. Rev.* 79, 709–725. doi:10.1093/nutrit/nuaa020
- Gillam-Krakauer, M., and Gowen, C. W., Jr (2021). "Birth Asphyxia," in *StatPearls* (Treasure Island (FL): StatPearls Publishing). Available at: <http://www.ncbi.nlm.nih.gov/books/NBK430782/> (Accessed November 28, 2021).
- Gomez-Castro, F., Zappettini, S., Pressey, J. C., Silva, C. G., Russeau, M., Gervasi, N., et al. (2021). Convergence of Adenosine and GABA Signaling for Synapse Stabilization During Development. *Science* 374, eabk2055. doi:10.1126/science.abk2055
- Gonçalves, L. K., da Silva, I. R. V., Cechinel, L. R., Frusciante, M. R., de Mello, A. S., Elsnér, V. R., et al. (2017). Maternal Consumption of High-Fat Diet and Grape Juice Modulates Global Histone H4 Acetylation Levels in Offspring hippocampus: A Preliminary Study. *Neurosci. Lett.* 661, 29–32. doi:10.1016/j.neulet.2017.09.042
- Guo, J. U., Su, Y., Shin, J. H., Shin, J., Li, H., Xie, B., et al. (2014). Distribution, Recognition and Regulation of Non-CpG Methylation in the Adult Mammalian Brain. *Nat. Neurosci.* 17, 215–222. doi:10.1038/nn.3607
- Hall, P., Adami, H.-O., Trichopoulos, D., Pedersen, N. L., Laggiou, P., Ekblom, A., et al. (2004). Effect of Low Doses of Ionising Radiation in Infancy on Cognitive Function in Adulthood: Swedish Population Based Cohort Study. *BMJ* 328, 19. doi:10.1136/bmj.328.7430.19
- Harker, A., Carroll, C., Raza, S., Kolb, B., and Gibb, R. (2018). Preconception Paternal Stress in Rats Alters Brain and Behavior in Offspring. *Neuroscience* 388, 474–485. doi:10.1016/j.neuroscience.2018.06.034
- Harker, A., Raza, S., Williams, K., Kolb, B., and Gibb, R. (2015). Preconception Paternal Stress in Rats Alters Dendritic Morphology and Connectivity in the Brain of Developing Male and Female Offspring. *Neuroscience* 303, 200–210. doi:10.1016/j.neuroscience.2015.06.058
- Hasebe, K., Kendig, M. D., and Morris, M. J. (2021). Mechanisms Underlying the Cognitive and Behavioural Effects of Maternal Obesity. *Nutrients* 13, 240. doi:10.3390/nu13010240
- Hawkes, C. A., Gentleman, S. M., Nicoll, J. A., and Carare, R. O. (2015). Prenatal High-Fat Diet Alters the Cerebrovasculature and Clearance of β -Amyloid in Adult Offspring. *J. Pathol.* 235, 619–631. doi:10.1002/path.4468
- Hefti, M. M., Farrell, K., Kim, S., Bowles, K. R., Fowkes, M. E., Raj, T., et al. (2018). High-resolution Temporal and Regional Mapping of MAPT Expression and Splicing in Human Brain Development. *PLOS ONE* 13, e0195771. doi:10.1371/journal.pone.0195771
- Hefti, M. M., Kim, S., Bell, A. J., Betters, R. K., Fiock, K. L., Iida, M. A., et al. (2019). Tau Phosphorylation and Aggregation in the Developing Human Brain. *J. Neuropathol. Exp. Neurol.* 78, 930–938. doi:10.1093/jnen/nl073
- Hines, E. P., Mendola, P., von Ehrenstein, O. S., Ye, X., Calafat, A. M., and Fenton, S. E. (2015). Concentrations of Environmental Phenols and Parabens in Milk, Urine and Serum of Lactating North Carolina Women. *Reprod. Toxicol.* 54, 120–128. doi:10.1016/j.reprotox.2014.11.006
- Hombach, S., and Kretz, M. (2016). Non-coding RNAs: Classification, Biology and Functioning. *Adv. Exp. Med. Biol.* 937, 3–17. doi:10.1007/978-3-319-42059-2_1
- Hoover, B. R., Reed, M. N., Su, J., Penrod, R. D., Kotilinek, L. A., Grant, M. K., et al. (2010). Tau Mislocalization to Dendritic Spines Mediates Synaptic Dysfunction Independently of Neurodegeneration. *Neuron* 68, 1067–1081. doi:10.1016/j.neuron.2010.11.030
- Hui, J., Feng, G., Zheng, C., Jin, H., and Jia, N. (2017). Maternal Separation Exacerbates Alzheimer's Disease-like Behavioral and Pathological Changes in Adult APP^{swe}/PS1^{dE9} Mice. *Behav. Brain Res.* 318, 18–23. doi:10.1016/j.bbr.2016.10.030
- Ikegami, S., Harada, A., and Hirokawa, N. (2000). Muscle Weakness, Hyperactivity, and Impairment in Fear Conditioning in Tau-Deficient Mice. *Neurosci. Lett.* 279, 129–132. doi:10.1016/S0304-3940(99)00964-7
- Izquierdo, V., Palomera-Ávalos, V., Pallás, M., and Grinán-Ferré, C. (2021). Correction: Izquierdo et al. Resveratrol Supplementation Attenuates Cognitive and Molecular Alterations under Maternal High-Fat Diet Intake: Epigenetic Inheritance over Generations. *Int. J. Mol. Sci.* 2021, 22, 1453. *Int. J. Mol. Sci.* 22, 9155. doi:10.3390/ijms22179155
- Jang, H., Shin, W., Lee, J., and Do, J. (2017). CpG and Non-CpG Methylation in Epigenetic Gene Regulation and Brain Function. *Genes* 8, 148. doi:10.3390/genes8060148
- Janson, J., Laedtke, T., Parisi, J. E., O'Brien, P., Petersen, R. C., and Butler, P. C. (2004). Increased Risk of Type 2 Diabetes in Alzheimer Disease. *Diabetes* 53, 474–481. doi:10.2337/diabetes.53.2.474
- Jeugd, A. V. d., Ahmed, T., Burnouf, S., Belarbi, K., Hamdame, M., Grosjean, M.-E., et al. (2011). Hippocampal Tauopathy in Tau Transgenic Mice Coincides with Impaired Hippocampus-dependent Learning and Memory, and Attenuated Late-phase Long-Term Depression of Synaptic Transmission. *Neurobiol. Learn. Mem.* 95, 296–304. doi:10.1016/j.nlm.2010.12.005
- Ji, H.-F., and Shen, L. (2021). Probiotics as Potential Therapeutic Options for Alzheimer's Disease. *Appl. Microbiol. Biotechnol.* 105, 7721–7730. doi:10.1007/s00253-021-11607-1
- Johansson, N., Viberg, H., Fredriksson, A., and Eriksson, P. (2008). Neonatal Exposure to Deca-Brominated Diphenyl Ether (PBDE 209) Causes Dose-Response Changes in Spontaneous Behaviour and Cholinergic Susceptibility in Adult Mice. *Neurotoxicology* 29, 911–919. doi:10.1016/j.neuro.2008.09.008
- Kaczurkin, A. N., Raznahan, A., and Satterthwaite, T. D. (2019). Sex Differences in the Developing Brain: Insights from Multimodal Neuroimaging. *Neuropsychopharmacol.* 44, 71–85. doi:10.1038/s41386-018-0111-z
- Kempf, S. J., Casciati, A., Buratovic, S., Janik, D., Toerne, C. v., Ueffing, M., et al. (2014). The Cognitive Defects of Neonatally Irradiated Mice Are Accompanied by Changed Synaptic Plasticity, Adult Neurogenesis and Neuroinflammation. *Mol. Neurodegeneration* 9, 57. doi:10.1186/1750-1326-9-57
- Killin, L. O. J., Starr, J. M., Shiue, I. J., and Russ, T. C. (2016). Environmental Risk Factors for Dementia: a Systematic Review. *BMC Geriatr.* 16, 1–28. doi:10.1186/s12877-016-0342-y
- Kundakovic, M., and Jaric, I. (2017). The Epigenetic Link between Prenatal Adverse Environments and Neurodevelopmental Disorders. *Genes* 8, 104. doi:10.3390/genes8030104
- La Maestra, S., Kisby, G. E., Micale, R. T., Johnson, J., Kow, Y. W., Bao, G., et al. (2011). Cigarette Smoke Induces DNA Damage and Alters Base-Excision Repair and Tau Levels in the Brain of Neonatal Mice. *Toxicol. Sci.* 123, 471–479. doi:10.1093/toxsci/kfr187
- La Rue, A. (2010). Healthy Brain Aging: Role of Cognitive reserve, Cognitive Stimulation, and Cognitive Exercises. *Clin. Geriatr. Med.* 26, 99–111. doi:10.1016/j.cger.2009.11.003
- Lacal, L., and Ventura, R. (2018). Epigenetic Inheritance: Concepts, Mechanisms and Perspectives. *Front. Mol. Neurosci.* 11, 292. doi:10.3389/fnmol.2018.00292
- Landrigan, P. J., Sonawane, B., Butler, R. N., Trasande, L., Callan, R., and Droller, D. (2005). Early Environmental Origins of Neurodegenerative Disease in Later Life. *Environ. Health Perspect.* 113, 1230–1233. doi:10.1289/ehp.7571
- Laurent, C., Eddarkaoui, S., Derisbourg, M., Leboucher, A., Demeyer, D., Carrier, S., et al. (2014). Beneficial Effects of Caffeine in a Transgenic Model of Alzheimer's Disease-Like Tau Pathology. *Neurobiol. Aging* 35, 2079–2090. doi:10.1016/j.neurobiolaging.2014.03.027
- Lei, P., Ayton, S., Finkelstein, D. I., Spoerri, L., Ciccostoto, G. D., Wright, D. K., et al. (2012). Tau Deficiency Induces Parkinsonism with Dementia by Impairing APP-Mediated Iron export. *Nat. Med.* 18, 291–295. doi:10.1038/nm.2613
- Lemche, E. (2018). Early Life Stress and Epigenetics in Late-Onset Alzheimer's Dementia: A Systematic Review. *Cg* 19, 522–602. doi:10.2174/1389202919666171229145156
- Lesuis, S. L., Hoeijmakers, L., Korosi, A., de Rooij, S. R., Swaab, D. F., Kessels, H. W., et al. (2018). Vulnerability and Resilience to Alzheimer's Disease: Early Life Conditions Modulate Neuropathology and Determine Cognitive reserve. *Alz Res. Ther.* 10, 95. doi:10.1186/s13195-018-0422-7
- Liu, B., Liu, J., and Shi, J.-S. (2020). SAMP8 Mice as a Model of Age-Related Cognition Decline with Underlying Mechanisms in Alzheimer's Disease. *Jad*, 385–395. doi:10.3233/JAD-200063
- Liu, X., Li, X., Xia, B., Jin, X., Zou, Q., Zeng, Z., et al. (2021). High-Fiber Diet Mitigates Maternal Obesity-Induced Cognitive and Social Dysfunction in the Offspring Via Gut-Brain Axis. *Cell Metab.* 33, 923–938.e6. doi:10.1016/j.cmet.2021.02.002
- Livingston, G., Huntley, J., Sommerlad, A., Ames, D., Ballard, C., Banerjee, S., et al. (2020). Dementia Prevention, Intervention, and Care: 2020 Report of the

- Lancet Commission. *The Lancet* 396, 413–446. doi:10.1016/S0140-6736(20)30367-6
- Lo, A. C., Iscru, E., Blum, D., Tesseur, I., Callaerts-Vegh, Z., Bué, L., et al. (2013). Amyloid and Tau Neuropathology Differentially Affect Prefrontal Synaptic Plasticity and Cognitive Performance in Mouse Models of Alzheimer's Disease. *Jad* 37, 109–125. doi:10.3233/JAD-122296
- Luo, G., Wei, R., Wang, S., and Wang, J. (2017). Paternal Bisphenol A Diet Changes Prefrontal Cortex Proteome and Provokes Behavioral Dysfunction in Male Offspring. *Chemosphere* 184, 720–729. doi:10.1016/j.chemosphere.2017.06.050
- Marciniak, E., Leboucher, A., Caron, E., Ahmed, T., Tailleur, A., Dumont, J., et al. (2017). Tau Deletion Promotes Brain Insulin Resistance. *J. Exp. Med.* 214, 2257–2269. doi:10.1084/jem.20161731
- Marin, O., Valiente, M., Ge, X., and Tsai, L.-H. (2010). Guiding Neuronal Cell Migrations. *Cold Spring Harbor Perspect. Biol.* 2, a001834. doi:10.1101/cshperspect.a001834
- Martin, S. A. L., Jameson, C. H., Allan, S. M., and Lawrence, C. B. (2014). Maternal High-Fat Diet Worsens Memory Deficits in the Triple-Transgenic (3xTgAD) Mouse Model of Alzheimer's Disease. *PLoS One* 9, e99226. doi:10.1371/journal.pone.0099226
- Martisova, E., Aisa, B., Gueren, G., and Javier Ramirez, M. (2013). Effects of Early Maternal Separation on Biobehavioral and Neuropathological Markers of Alzheimer's Disease in Adult Male Rats. *Car* 10, 420–432. doi:10.2174/1567205011310040007
- Massaro, A. N., Chang, T., Baumgart, S., McCarter, R., Nelson, K. B., and Glass, P. (2014). Biomarkers S100B and Neuron-specific Enolase Predict Outcome in Hypothermia-Treated Encephalopathic Newborns*. *Pediatr. Crit. Care Med.* 15, 615–622. doi:10.1097/PCC.0000000000000155
- Matsuzaki, T., Sasaki, K., Tanizaki, Y., Hata, J., Fujimi, K., Matsui, Y., et al. (2010). Insulin Resistance Is Associated with the Pathology of Alzheimer Disease: the Hisayama Study. *Neurology* 75, 764–770. doi:10.1212/WNL.0b013e3181ee25f
- McPherson, N. O., Bell, V. G., Zander-Fox, D. L., Fullston, T., Wu, L. L., Robker, R. L., et al. (2015). When Two Obese Parents Are Worse Than One! Impacts on Embryo and Fetal Development. *Am. J. Physiology-Endocrinology Metab.* 309, E568–E581. doi:10.1152/ajpendo.00230.2015
- Mellén, M., Ayata, P., Dewell, S., Kriacucionis, S., and Heintz, N. (2012). MeCP2 Binds to ShmC Enriched within Active Genes and Accessible Chromatin in the Nervous System. *Cell* 151, 1417–1430. doi:10.1016/j.cell.2012.11.022
- Metzger, F. (2010). Molecular and Cellular Control of Dendrite Maturation during Brain Development. *Cmp* 3, 1–11. doi:10.2174/1874467211003010001
- Miller, D. B., and O'Callaghan, J. P. (2008). Do early-life insults contribute to the late-life development of Parkinson and Alzheimer diseases. *Metabolism* 57 (Suppl. 2), S44–S49. doi:10.1016/j.metabol.2008.07.011
- Moceri, V. M., Kukull, W. A., Emanuel, I., van Belle, G., and Larson, E. B. (2000). Early-life Risk Factors and the Development of Alzheimer's Disease. *Neurology* 54, 415. doi:10.1212/wnl.54.2.415
- Modgil, S., Lahiri, D. K., Sharma, V. L., and Anand, A. (2014). Role of Early Life Exposure and Environment on Neurodegeneration: Implications on Brain Disorders. *Transl Neurodegener* 3, 9. doi:10.1186/2047-9158-3-9
- Mondragón-Rodríguez, S., Salas-Gallardo, A., González-Pereyra, P., Macías, M., Ordaz, B., Peña-Ortega, F., et al. (2018). Phosphorylation of Tau Protein Correlates with Changes in Hippocampal Theta Oscillations and Reduces Hippocampal Excitability in Alzheimer's Model. *J. Biol. Chem.* 293, 8462–8472. doi:10.1074/jbc.RA117.001187
- Mondragón-Rodríguez, S., Salgado-Burgos, H., and Peña-Ortega, F. (2020). Circuitry and Synaptic Dysfunction in Alzheimer's Disease: A New Tau Hypothesis. *Neural Plasticity* 2020, 1–11. doi:10.1155/2020/2960343
- Monti, N., Cavallaro, R. A., Stoccoro, A., Nicolai, V., Scarpa, S., Kovacs, G. G., et al. (2020). CpG and Non-CpG Presenilin1 Methylation Pattern in Course of Neurodevelopment and Neurodegeneration Is Associated with Gene Expression in Human and Murine Brain. *Epigenetics* 15, 781–799. doi:10.1080/15592294.2020.1722917
- Moody, L., Chen, H., and Pan, Y.-X. (2017). Early-Life Nutritional Programming of Cognition-The Fundamental Role of Epigenetic Mechanisms in Mediating the Relation between Early-Life Environment and Learning and Memory Process. *Adv. Nutr.* 8, 337–350. doi:10.3945/an.116.014209
- Mouw, J. K., Ou, G., and Weaver, V. M. (2014). Extracellular Matrix Assembly: a Multiscale Deconstruction. *Nat. Rev. Mol. Cell Biol* 15, 771–785. doi:10.1038/nrm3902
- Mueller, N. T., Bakacs, E., Combellick, J., Grigoryan, Z., and Dominguez-Bello, M. G. (2015). The Infant Microbiome Development: Mom Matters. *Trends Mol. Med.* 21, 109–117. doi:10.1016/j.molmed.2014.12.002
- Mychasiuk, R., Harker, A., Illynskyy, S., and Gibb, R. (2013). Paternal Stress Prior to conception Alters DNA Methylation and Behaviour of Developing Rat Offspring. *Neuroscience* 241, 100–105. doi:10.1016/j.neuroscience.2013.03.025
- Najar, J., Östling, S., Gudmundsson, P., Sundh, V., Johansson, L., Kern, S., et al. (2019). Cognitive and Physical Activity and Dementia. *Neurology* 92, e1322–e1330. doi:10.1212/WNL.0000000000007021
- Negishi, T., Kawasaki, K., Suzuki, S., Maeda, H., Ishii, Y., Kyuwa, S., et al. (2004). Behavioral Alterations in Response to Fear-Provoking Stimuli and Tranylcypromine Induced by Perinatal Exposure to Bisphenol A and Nonylphenol in Male Rats. *Environ. Health Perspect.* 112, 1159–1164. doi:10.1289/ehp.6961
- Nguyen, T. T., Ta, Q. T. H., Nguyen, T. K. O., Nguyen, T. T. D., and Van Giau, V. (2020). Type 3 Diabetes and its Role Implications in Alzheimer's Disease. *Ijms* 21, 3165. doi:10.3390/ijms21093165
- Nikolac Perkovic, M., Videtic Paska, A., Konjevod, M., Kouter, K., Svob Strac, D., Nedic Erjavec, G., et al. (2021). Epigenetics of Alzheimer's Disease. *Biomolecules* 11, 195. doi:10.3390/biom11020195
- Nizari, S., Carare, R. O., and Hawkes, C. A. (2016). Increased Aβ Pathology in Aged Tg2576 Mice Born to Mothers Fed a High Fat Diet. *Sci. Rep.* 6, 21981. doi:10.1038/srep21981
- Olayinka, O., Olayinka, O. O., Alemu, B. T., Akpinar-Elci, M., and Grossberg, G. T. (2019). Toxic Environmental Risk Factors for Alzheimer's Disease: A Systematic Review. *Aging Med. Healthc.* 10, 4–17. doi:10.33879/amh.2019.1727
- Parsons, J. T., Horwitz, A. R., and Schwartz, M. A. (2010). Cell Adhesion: Integrating Cytoskeletal Dynamics and Cellular Tension. *Nat. Rev. Mol. Cell Biol* 11, 633–643. doi:10.1038/nrm2957
- Perera, F. P., Rauh, V., Whyatt, R. M., Tang, D., Tsai, W. Y., Bernert, J. T., et al. (2005). A Summary of Recent Findings on Birth Outcomes and Developmental Effects of Prenatal ETS, PAH, and Pesticide Exposures. *Neurotoxicology* 26, 573–587. doi:10.1016/j.neuro.2004.07.007
- Polito, C., Cai, Z.-Y., Shi, Y.-L., Li, X.-M., Yang, R., Shi, M., et al. (2018). Association of Tea Consumption with Risk of Alzheimer's Disease and Anti-beta-amyloid Effects of Tea. *Nutrients* 10, 655. doi:10.3390/nu10050655
- Polydoro, M., Acker, C. M., Duff, K., Castillo, P. E., and Davies, P. (2009). Age-dependent Impairment of Cognitive and Synaptic Function in the Htau Mouse Model of Tau Pathology. *J. Neurosci.* 29, 10741–10749. doi:10.1523/JNEUROSCI.1065-09.2009
- Poston, L., Caleyachetty, R., Nattangius, S., Corvalán, C., Uauy, R., Herring, S., et al. (2016). Preconceptional and Maternal Obesity: Epidemiology and Health Consequences. *Lancet Diabetes Endocrinol.* 4, 1025–1036. doi:10.1016/S2213-8587(16)30217-0
- Proença, I. C. T., Gonçalves, L. K., Schmitz, F., Mello, A., Funchal, C. S., Wyse, A., et al. (2021). Purple Grape Juice Consumption during the Gestation Reduces Acetylcholinesterase Activity and Oxidative Stress Levels Provoked by High-Fat Diet in hippocampus from Adult Female Rats Descendants. *Acad. Bras. Ciênc.* 93, e20191002. doi:10.1590/0001-376520210191002
- Rahman, M. H., Akter, R., Bhattacharya, T., Abdel-Daim, M. M., Alkahtani, S., Arafah, M. W., et al. (2020). Resveratrol and Neuroprotection: Impact and its Therapeutic Potential in Alzheimer's Disease. *Front. Pharmacol.* 11, 619024. doi:10.3389/fphar.2020.619024
- Ramsay, H., Barnett, J. H., Murray, G. K., Mäki, P., Hurtig, T., Nordström, T., et al. (2016). Smoking in Pregnancy, Adolescent Mental Health and Cognitive Performance in Young Adult Offspring: Results from a Matched Sample within a Finnish Cohort. *BMC Psychiatry* 16, 430. doi:10.1186/s12888-016-1142-9
- Reitz, C., Brayne, C., and Mayeux, R. (2011). Epidemiology of Alzheimer Disease. *Nat. Rev. Neurol.* 7, 137–152. doi:10.1038/nrneurol.2011.2

- Reitz, C., and Mayeux, R. (2009). Endophenotypes in normal Brain Morphology and Alzheimer's Disease: a Review. *Neuroscience* 164, 174–190. doi:10.1016/j.neuroscience.2009.04.006
- Rostami, F., Javan, M., Moghimi, A., Haddad-Mashadrizheh, A., and Fereidoni, M. (2020). Prenatal Stress Promotes Icv-STZ-Induced Sporadic Alzheimer's Pathology through central Insulin Signaling Change. *Life Sci.* 241, 117154. doi:10.1016/j.lfs.2019.117154
- Rothman, S. M., Herdener, N., Camandola, S., Texel, S. J., Mughal, M. R., Cong, W.-N., et al. (2012). 3xTgAD Mice Exhibit Altered Behavior and Elevated A β after Chronic Mild Social Stress. *Neurobiol. Aging* 33, 830. e1. doi:10.1016/j.neurobiolaging.2011.07.005
- Ryan, B. C., and Vandenbergh, J. G. (2006). Developmental Exposure to Environmental Estrogens Alters Anxiety and Spatial Memory in Female Mice. *Horm. Behav.* 50, 85–93. doi:10.1016/j.yhbeh.2006.01.007
- Sanguinetti, E., Guzzardi, M. A., Tripodi, M., Panetta, D., Selma-Royo, M., Zega, A., et al. (2019). Microbiota Signatures Relating to Reduced Memory and Exploratory Behaviour in the Offspring of Overweight Mothers in a Murine Model. *Sci. Rep.* 9, 12609. doi:10.1038/s41598-019-48090-8
- Schaffer, T. K., Wohlenberg, M. F., de Souza Machado, F., Bortolato, G., Marinho, J. P., da Silva Medeiros, N., et al. (2019). Chronic Consumption of Purple Grape Juice in Gestational-Lactation and post Lactation Promotes Anxiolytic Effect and Antioxidant Defense Improvement in Brain from Wistar Male Offsprings. *J. Nutr. Intermediary Metab.* 15, 46–54. doi:10.1016/j.jnim.2018.12.005
- Sennvik, K., Boekhoorn, K., Lasrado, R., Terwel, D., Verhaeghe, S., Korrr, H., et al. (2007). Tau-4R Suppresses Proliferation and Promotes Neuronal Differentiation in the hippocampus of Tau Knockin/Knockout Mice. *FASEB j.* 21, 2149–2161. doi:10.1096/fj.06-7735com
- Sergeant, N., Bretteville, A., Hamdane, M., Caillet-Boudin, M.-L., Grognet, P., Bombois, S., et al. (2008). Biochemistry of Tau in Alzheimer's Disease and Related Neurological Disorders. *Expert Rev. Proteomics* 5, 207–224. doi:10.1586/14789450.5.2.207
- Shi, D.-Q., Ali, I., Tang, J., and Yang, W.-C. (2017). New Insights into 5hmC DNA Modification: Generation, Distribution and Function. *Front. Genet.* 8, 100. doi:10.3389/fgene.2017.00100
- Sierksma, A. S. R., Vanmierlo, T., De Vry, J., Raijmakers, M. E. A., Steinbusch, H. W.-N., van den Hove, D. L. A., et al. (2012). Effects of Prenatal Stress Exposure on Soluble A β and Brain-Derived Neurotrophic Factor Signaling in Male and Female APPswe/PS1dE9 Mice. *Neurochem. Int.* 61, 697–701. doi:10.1016/j.neuint.2012.06.022
- Singh, S., and Li, S. S.-L. (2012). Epigenetic Effects of Environmental Chemicals Bisphenol A and Phthalates. *Ijms* 13, 10143–10153. doi:10.3390/ijms130810143
- Sosa, L. J., Cáceres, A., Dupraz, S., Oksdath, M., Quiroga, S., and Lorenzo, A. (2017). The Physiological Role of the Amyloid Precursor Protein as an Adhesion Molecule in the Developing Nervous System. *J. Neurochem.* 143, 11–29. doi:10.1111/jnc.14122
- Stoyanova, E., Riad, M., Rao, A., and Heintz, N. (2021). 5-Hydroxymethylcytosine-mediated Active Demethylation Is Required for Mammalian Neuronal Differentiation and Function. *Elife* 10, e66973. doi:10.7554/eLife.66973
- Stroud, H., Su, S. C., Hrvatin, S., Greben, A. W., Renthal, W., Boxer, L. D., et al. (2017). Early-Life Gene Expression in Neurons Modulates Lasting Epigenetic States. *Cell* 171, 1151–1164. e16. doi:10.1016/j.cell.2017.09.047
- Sukjamnong, S., Thongkorn, S., Kanlayaprasit, S., Saeliw, T., Hussem, K., Warayanon, W., et al. (2020). Prenatal Exposure to Bisphenol A Alters the Transcriptome-Interactome Profiles of Genes Associated with Alzheimer's Disease in the Offspring hippocampus. *Sci. Rep.* 10, 9487. doi:10.1038/s41598-020-65229-0
- Sun, W., Ban, J.-B., Zhang, N., Zu, Y.-K., and Sun, W.-X. (2014). Perinatal exposure to Di-(2-ethylhexyl)-Phthalate leads to cognitive dysfunction and phospho-tau level increase in aged rats. *Environ Toxicol* 29, 596–603. doi:10.1002/tox.21785
- Syeda, T., Sanchez-Tapia, M., Pinedo-Vargas, L., Granados, O., Cuervo-Zanatta, D., Rojas-Santiago, E., et al. (2018). Bioactive Food Abates Metabolic and Synaptic Alterations by Modulation of Gut Microbiota in a Mouse Model of Alzheimer's Disease. *Jad* 66, 1657–1682. doi:10.3233/JAD-180556
- Talbot, K., Wang, H.-Y., Kazi, H., Han, L.-Y., Bakshi, K. P., Stucky, A., et al. (2012). Demonstrated Brain Insulin Resistance in Alzheimer's Disease Patients Is Associated with IGF-1 Resistance, IRS-1 Dysregulation, and Cognitive Decline. *J. Clin. Invest.* 122, 1316–1338. doi:10.1172/JCI59903
- Tao, H., Xie, P., Cao, Y., Shu, L., Li, L., Chen, J., et al. (2018). The Dynamic DNA Demethylation during Postnatal Neuronal Development and Neural Stem Cell Differentiation. *Stem Cell Int.* 2018, 1–10. doi:10.1155/2018/2186301
- Tian, Y.-H., Baek, J.-H., Lee, S.-Y., and Jang, C.-G. (2010). Prenatal and Postnatal Exposure to Bisphenol A Induces Anxiolytic Behaviors and Cognitive Deficits in Mice. *Synapse* 64, 432–439. doi:10.1002/syn.20746
- Tini, G., Scagliola, R., Monacelli, F., La Malfa, G., Porto, I., Brunelli, C., et al. (2020). Alzheimer's Disease and Cardiovascular Disease: A Particular Association. *Cardiol. Res. Pract.* 2020, 1–10. doi:10.1155/2020/2617970
- Tortelli, R., Lozupone, M., Guerra, V., Barulli, M. R., Imbimbo, B. P., Capozzo, R., et al. (2017). Midlife Metabolic Profile and the Risk of Late-Life Cognitive Decline. *Jad* 59, 121–130. doi:10.3233/JAD-170153
- Tunc, T., Karaoglu, A., Cayci, T., Demirkaya, E., Kul, M., Yaman, H., et al. (2010). The Relation between Delivery Type and Tau Protein Levels in Cord Blood. *Pediatr. Int.* 52, 872–875. doi:10.1111/j.1442-200X.2010.03213.x
- Van der Jeugd, A., Vermaercke, B., Derisbourg, M., Lo, A. C., Hamdane, M., Blum, D., et al. (2013). Progressive Age-Related Cognitive Decline in Tau Mice. *Jad* 37, 777–788. doi:10.3233/JAD-130110
- Viberg, H. (2009). Exposure to Polybrominated Diphenyl Ethers 203 and 206 during the Neonatal Brain Growth Spurt Affects Proteins Important for normal Neurodevelopment in Mice. *Toxicol. Sci.* 109, 306–311. doi:10.1093/toxsci/kfp074
- Viberg, H., Fredriksson, A., Buratovic, S., and Eriksson, P. (2011). Dose-dependent Behavioral Disturbances after a Single Neonatal Bisphenol A Dose. *Toxicology* 290, 187–194. doi:10.1016/j.tox.2011.09.006
- Vyas, S., Rodrigues, A. J., Silva, J. M., Tronche, F., Almeida, O. F. X., Sousa, N., et al. (2016). Chronic Stress and Glucocorticoids: From Neuronal Plasticity to Neurodegeneration. *Neural Plasticity* 2016, 1–15. doi:10.1155/2016/6391686
- Wajman, J. R., Mansur, L. L., and Yassuda, M. S. (2019). Lifestyle Patterns as a Modifiable Risk Factor for Late-Life Cognitive Decline: A Narrative Review Regarding Dementia Prevention. *Cas* 11, 90–99. doi:10.2174/1874609811666181003160225
- Wang, C., Niu, R., Zhu, Y., Han, H., Luo, G., Zhou, B., et al. (2014). Changes in Memory and Synaptic Plasticity Induced in Male Rats after Maternal Exposure to Bisphenol A. *Toxicology* 322, 51–60. doi:10.1016/j.tox.2014.05.001
- Wang, J.-Z., and Liu, F. (2008). Microtubule-associated Protein Tau in Development, Degeneration and protection of Neurons. *Prog. Neurobiol.* 85, 148–175. doi:10.1016/j.pneurobio.2008.03.002
- Weaver, I. C. G., Cervoni, N., Champagne, F. A., D'Alessio, A. C., Sharma, S., Seckl, J. R., et al. (2004). Epigenetic Programming by Maternal Behavior. *Nat. Neurosci.* 7, 847–854. doi:10.1038/nn1276
- Wijesekara, N., Gonçalves, R. A., Ahrens, R., De Felice, F. G., and Fraser, P. E. (2018). Tau Ablation in Mice Leads to Pancreatic β Cell Dysfunction and Glucose Intolerance. *FASEB j.* 32, 3166–3173. doi:10.1096/fj.201701352
- Wijesekara, N., Gonçalves, R. A., Ahrens, R., Ha, K., De Felice, F. G., and Fraser, P. E. (2021). Combination of Human Tau and Islet Amyloid Polypeptide Exacerbates Metabolic Dysfunction in Transgenic Mice. *J. Pathol.* 254, 244–253. doi:10.1002/path.5674
- World Health Organization (2021). Dementia. Available at: <https://www.who.int/news-room/fact-sheets/detail/dementia> (Accessed November 28, 2021).
- Wu, J., Basha, M. R., Brock, B., Cox, D. P., Cardozo-Pelaez, F., McPherson, C. A., et al. (2008). Alzheimer's Disease (AD)-Like Pathology in Aged Monkeys after Infantile Exposure to Environmental Metal Lead (Pb): Evidence for a Developmental Origin and Environmental Link for AD. *J. Neurosci.* 28, 3–9. doi:10.1523/JNEUROSCI.4405-07.2008
- Xu, X.-h., Zhang, J., Wang, Y.-m., Ye, Y.-p., and Luo, Q.-q. (2010). Perinatal Exposure to Bisphenol-A Impairs Learning-Memory by Concomitant Down-Regulation of N-Methyl-D-Aspartate Receptors of hippocampus in Male Offspring Mice. *Horm. Behav.* 58, 326–333. doi:10.1016/j.yhbeh.2010.02.012
- Xue, J., Zhang, L., Xie, X., Gao, Y., Jiang, L., Wang, J., et al. (2020). Prenatal Bisphenol A Exposure Contributes to Tau Pathology: Potential Roles of CDK5/GSK3 β /PP2A axis in BPA-Induced Neurotoxicity. *Toxicology* 438, 152442. doi:10.1016/j.tox.2020.152442
- Yao, T., Itoh, K., Nakamura, K., Ogi, H., Fujiwara, Y., and Fushiki, S. (2008). Genome-wide Analysis of Epigenomic Alterations in Fetal Mouse Forebrain after Exposure to Low Doses of Bisphenol A.

- Biochem. Biophysical Res. Commun.* 376, 563–567. doi:10.1016/j.bbrc.2008.09.028
- Yu, Y., Run, X., Liang, Z., Li, Y., Liu, F., Liu, Y., et al. (2009). Developmental Regulation of Tau Phosphorylation, Tau Kinases, and Tau Phosphatases. *J. Neurochem.* 108, 1480–1494. doi:10.1111/j.1471-4159.2009.05882.x
- Zappettini, S., Faivre, E., Ghestem, A., Carrier, S., Buée, L., Blum, D., et al. (2019). Caffeine Consumption during Pregnancy Accelerates the Development of Cognitive Deficits in Offspring in a Model of Tauopathy. *Front. Cel. Neurosci.* 13, 438. doi:10.3389/fncel.2019.00438
- Zhang, S., Li, X., Wang, Z., Liu, Y., Gao, Y., Tan, L., et al. (2017). Paternal Spatial Training Enhances Offspring's Cognitive Performance and Synaptic Plasticity in Wild-type but Not Improve Memory Deficit in Alzheimer's Mice. *Sci. Rep.* 7, 1521. doi:10.1038/s41598-017-01811-3
- Zhang, Y., Zhang, Z., Li, L., Xu, K., Ma, Z., Chow, H. M., et al. (2020). Selective Loss of 5hmC Promotes Neurodegeneration in the Mouse Model of Alzheimer's Disease. *FASEB j.* 34, 16364–16382. doi:10.1096/fj.202001271R
- Zhao, W. Q., De Felice, F. G., Fernandez, S., Chen, H., Lambert, M. P., Quon, M. J., et al. (2008). Amyloid Beta Oligomers Induce Impairment of Neuronal Insulin Receptors. *FASEB j.* 22, 246–260. doi:10.1096/fj.06-7703com
- Zhou, Y., Kong, Y., Fan, W., Tao, T., Xiao, Q., Li, N., et al. (2020). Principles of RNA Methylation and Their Implications for Biology and Medicine. *Biomed. Pharmacother.* 131, 110731. doi:10.1016/j.biopha.2020.110731
- Zhou, Y., Zhu, H., Wu, H. Y., Jin, L. Y., Chen, B., Pang, H. Y., et al. (2018). Diet-Induced Paternal Obesity Impairs Cognitive Function in Offspring by Mediating Epigenetic Modifications in Spermatozoa. *Obesity* 26, 1749–1757. doi:10.1002/oby.22322
- Conflict of Interest:** The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.
- Publisher's Note:** All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.
- Copyright © 2022 Gauvrit, Benderradji, Buée, Blum and Vieau. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.