



HAL
open science

Role of αv integrins in mesangial cell adhesion to vitronectin and von Willebrand factor

Zakia Hafdi, Philippe Lesavre, Pierre Louis Tharaux, Gilles Bessou, Dominique Baruch, Lise Halbwachs-Mecarelli

► **To cite this version:**

Zakia Hafdi, Philippe Lesavre, Pierre Louis Tharaux, Gilles Bessou, Dominique Baruch, et al.. Role of αv integrins in mesangial cell adhesion to vitronectin and von Willebrand factor. *Kidney International*, 1997, 51 (6), pp.1900-1907. 10.1038/ki.1997.259 . inserm-03788592

HAL Id: inserm-03788592

<https://inserm.hal.science/inserm-03788592v1>

Submitted on 26 Sep 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Role of α_v integrins in mesangial cell adhesion to vitronectin and von Willebrand factor

ZAKIA HAFDI, PHILIPPE LESAVRE,¹ PIERRE LOUIS THARAUX, GILLES BESSOU, DOMINIQUE BARUCH, and LISE HALBWACHS-MECARELLI

INSERM U 90 and Department of Nephrology, Necker Hospital, Paris, and INSERM U 143, Hôpital de Bicêtre, Bicêtre, France

Role of α_v integrins in mesangial cells adhesion to vitronectin and von Willebrand factor. This study demonstrates (by flow cytometry and immunoprecipitation after cell surface radiolabeling and by using monoclonal antibodies to α_v , β_3 , and $\alpha_v\beta_3$ and $\alpha_v\beta_5$ complexes) that $\alpha_v\beta_3$, the vitronectin receptor, and $\alpha_v\beta_5$ are expressed *in vitro* on cultured human mesangial cells (HMC) of the 5th to 8th passages. Antibodies to α_v , β_3 and $\alpha_v\beta_3$ respectively precipitated an $\alpha\beta$ heterodimer with molecular weights of 140 and 97 kDa. We analyzed the role of the various integrins in HMC interactions with vitronectin, and with fibronectin and von Willebrand factor (vWf), which are synthesized respectively by mesangial and endothelial cells. Cell adhesion increased in a dose dependent manner with the concentration of plastic-coated matrix protein and vWf. Inhibition of cell attachment with monoclonal antibodies to integrins indicated that HMC adhesion to vWf primarily involves $\alpha_v\beta_3$, and that $\alpha_v\beta_5$ may also contribute to cell binding to vWf. Adhesion to vitronectin involves both $\alpha_v\beta_3$ and $\alpha_v\beta_5$ complexes. In contrast, adhesion to fibronectin was not affected by monoclonal antibodies to $\alpha_v\beta_3$ and $\alpha_v\beta_5$ complexes. We propose that integrins $\alpha_v\beta_3$ and $\alpha_v\beta_5$, present on HMC, could mediate an interaction between mesangial and endothelial cells by binding to vWf, released at the basal site of endothelial cells.

Integrins are heterodimeric transmembrane glycoproteins consisting of varying combinations of noncovalently bound α and β chains, and are involved in adhesive interactions with the extracellular matrix. They are critical for normal physiological processes such as embryonic development, tissue structure and wound repair [1].

Integrins that contain the α_v chain mediate attachment of a variety of cells to diverse matrix proteins [2] and are critical for angiogenesis [3, 4], migration [5], cell proliferation and survival [6–8]. The integrin $\alpha_v\beta_3$, the vitronectin receptor, is the most promiscuous member of the integrin family, allowing cells to interact with a wide variety of proteins including vitronectin, fibrinogen, von Willebrand factor, thrombospondin, fibronectin, osteopontin, laminin and denatured collagen types I and IV [2]. $\alpha_v\beta_3$ is expressed on a variety of cells including endothelial cells, fibroblasts, melanoma, osteosarcoma and smooth muscle cells. In addition to β_3 , α_v chain associates with multiple β chains, that is,

β_1 , β_5 , β_6 and β_8 . The nature of cellular interactions with matrix proteins apparently depends on the β subunit. $\alpha_v\beta_1$ and $\alpha_v\beta_5$ recognize fibronectin and vitronectin [9], while $\alpha_v\beta_6$ binds to fibronectin [10, 11] and $\alpha_v\beta_8$ binds to vitronectin [12].

The renal localization of β_3 integrins has been investigated in developing and adult kidneys [13–15]. In the human adult glomerulus, $\alpha_v\beta_3$ is normally strongly expressed by glomerular endothelial cells, capsular epithelial cells and podocytes, and to a lesser extent by mesangial cells [13–18]. In the developing glomeruli, the β_3 chain is expressed in the cells of Bowman's capsule [19]. The role of $\alpha_v\beta_3$ in the metanephric development is suggested by its spatio-temporal distribution that varies with time, is restricted to the epithelia before vascularization, and is concentrated in the glomerular and extraglomerular vasculature in the mature kidney [19].

In the present study, we investigated the presence and the adhesive function of the α_v family of integrins on human mesangial cells in culture. We present evidence that $\alpha_v\beta_3$ and $\alpha_v\beta_5$ and possibly $\alpha_v\beta_1$ integrins are expressed at the membrane of this cell type. We also show that adhesion of cultured HMC to vitronectin and to vWf involves $\alpha_v\beta_3$ and $\alpha_v\beta_5$.

Methods

Antibodies

The antibodies used in this study are presented in Table 1.

Monoclonal antibodies included: anti- α_v (clone AMF/7), anti- α_6 (clone GoH3), anti- α_2 (clone Gi9), anti- β_1 (clone Lia 1/2) and control mouse IgG were from Immunotech (Marseille, France); anti- α_3 (clone P1B5) was from Chemicon (Temecula, CA, USA); anti- α_5 (clone 11, 16) was a generous gift from Dr. Steven K. Akiyama (Laboratory of Developmental Biology, NIH, MD, USA); anti- α_v (clone LM142, ascites) and anti- $\alpha_v\beta_3$ (clone LM 609, ascites) were a generous gift from Dr. David Cheresh (Scripps Research Institute, La Jolla, CA, USA); anti- β_3 (clone AP3, ascites) was a generous gift from Dr. Peter J. Newman (The Blood Center of Southeastern, WI, USA); anti- β_1 (clone K20) was a generous gift from Dr. Alain Bernard (INSERM U 343, Nice, France); anti- $\alpha_v\beta_5$ (clone P1F6) was from Chemicon; FITC-goat anti-mouse F(ab')₂ IgG fragments were from Jackson IRL (West Grove, PA, USA). The concentration for maximal monoclonal antibody binding was determined individually by flow cytometry and excess concentrations were used, as indicated,

¹ Current address: INSERM U 64, Hôpital Tenon, 4 rue de la Chine, 75020 Paris, France.

Received for publication August 16, 1996

and in revised form January 29, 1997

Accepted for publication January 30, 1997

© 1997 by the International Society of Nephrology

Table 1. Antibodies to integrins used in this study

Specificity	Clone	Species	Reference
α_2	Gi9	mouse	[20]
α_3	P1B5 ^a	mouse	[21]
α_5	11	mouse	[22]
	16 ^a	mouse	
α_6	GoH3 ^a	rat	[23]
α_v	AMF7	mouse	[24]
	LM142	mouse	[25]
β_1	Lia1/2 ^a	mouse	[26]
	K20	mouse	[27]
β_3	AP3	mouse	[28]
β_5	polyclonal	rabbit	[29]
$\alpha_v\beta_3$	LM609 ^a	mouse	[25]
$\alpha_v\beta_5$	PIF6 ^a	mouse	[30]

^aAntibodies known to inhibit cell adhesion. The origin of these antibodies is indicated in the text.

during incubation both for flow cytometry analysis and adhesion inhibition experiments. Rabbit polyclonal anti- β_5 antibodies were from Novus Molecular (San Diego, CA, USA). Biotinylated goat anti-mouse IgG and anti-rabbit IgG were respectively from Immunotech and from Novus Molecular.

Matrix components

Human vitronectin was purchased from Becton Dickinson (Bedford, MA, USA), human fibronectin from Sigma (St. Louis, MO, USA). According to the manufacturer, both proteins were purified to homogeneity (> 95%) as assessed by SDS PAGE analysis. Von Willebrand factor (vWf) was purified to homogeneity as described [31]. Purified vWf was homogeneous (> 95%) and vitronectin was not detectable by SDS PAGE analysis.

Human mesangial cells

Human mesangial cells (HMC) were prepared from the normal pole of kidney obtained after nephrectomy as previously described [32]. Briefly, each kidney was decapsulated, glomeruli were isolated from renal cortex by serial sieving, treated with collagenase type IV 300 UI/ml (Sigma) 20 to 30 minutes at 37°C, washed, and cultured in flasks with RPMI 1640 (ATGC, Paris, France) supplemented with 10% heat-inactivated fetal bovine serum FBS (ATGC), 2 mM L-glutamin, 60 UI/ml penicillin, 60 UI/ml streptomycin, and 10 mM HEPES at 37°C under 5% CO₂. HMC grown from glomeruli within two to three weeks were then subcultured and maintained in 10% FBS/RPMI medium. HMC were shown to be free of mycoplasma and characterized by morphology, α smooth muscle actin expression, absence of factor VIII expression, and used at the 5th to 8th passages.

Human endothelial cells

Human umbilical vein endothelial cells (HUVEC) were isolated as previously described by Jaffe et al [33]. HUVEC were cultured in M-199 (ATGC, Paris, France) supplemented with 20% heat-inactivated FBS (ATGC), 2 mM L-glutamine, 100 U/ml penicillin, 100 μ g/ml streptomycin, 0.25 μ g/ml amphotericin B, 10 mM HEPES, 50 μ g/ml heparin, 20 μ g/ml endothelial cell growth factor at 37°C under 5% CO₂, and used after two passages.

Fibroblasts

Fibroblasts were obtained from forearm skin biopsies as described [34] and grown in the same condition as we used for mesangial cell culture (see above) with 20% heat-inactivated FBS.

Flow cytometry

Expression of integrins by cultured cells in suspension was evaluated by flow cytometry. HMC were detached from culture flasks by trypsin (ATGC) treatment for five minutes at 37°C, washed with RPMI, counted and resuspended in PBS containing 1% BSA and 0.1% sodium azide. A total of 10⁵ cells were incubated at 4°C with anti-integrin antibodies for 30 minutes, washed three times with PBS and stained with fluorescein-conjugated goat anti-mouse IgG. Stained cells were fixed with 1% formaldehyde and analyzed on a FACScan flow cytometer (Becton Dickinson).

Adhesion assays

Adhesion assays were performed in 96-well plates (Costar, Cambridge, MA, USA) coated with 50 μ l/well of matrix proteins in PBS: fibronectin (1 μ g/ml), vitronectin (0.5 μ g/ml), vWf (5 μ g/ml). Plates were incubated at 37°C for 60 minutes, washed three times with phosphate-buffered saline (PBS) and unbound sites were blocked one hour at 37°C with 1% BSA (Sigma) previously heat-inactivated for 60 minutes at 56°C. Control wells were coated with 1% heat-inactivated BSA. HMC were detached from culture flasks by trypsin/EDTA treatment, and resuspended in RPMI with 0.1% BSA. A total of 10⁴ cells were allowed to adhere in each well at 37°C for 60 minutes. Non-adherent cells were then removed by washing with PBS, and adherent cells were fixed with 4% paraformaldehyde for five minutes. After rinsing with PBS, cells were stained for five minutes with 0.5% toluidine blue in 4% formaldehyde and rinsed with water. Cells were solubilized with 1% SDS and quantified in a microtiter plate reader at 630 nm. This procedure of quantitation gives values proportional to the number of adherent cells [24]. In adhesion inhibition experiments, cells were first preincubated for 30 minutes at 4°C with anti-integrin antibodies and treated thereafter as described above. All inhibition experiments were repeated independently, the number of which is indicated in each case.

Immunoprecipitation

Immunoprecipitation was performed on HMC. Cells were washed three times with PBS and surface labeled with 2 mCi/ml of Na¹²⁵I in tubes coated with iodogen (Pierce, Rockford, IL, USA), according to the manufacturer's instructions. After 30 minutes at 4°C, cells were washed and solubilized with 2% NP-40 in PBS containing protease inhibitors: 2 mM PMSF, 1 mg/ml SBTI, 1000 U/ml aprotinin, 5 mM iodoacetamide and 1 μ g/ml leupeptin. The cell lysate was cleared by centrifugation for 10 minutes at 4°C and the supernatant, precleared on normal rabbit IgG protein A-Sepharose, was incubated with different antibodies coupled to protein A-Sepharose for three hours at 4°C on a roller. Antigen-antibody complexes bound to the Sepharose beads were pelleted on a microcentrifuge and washed ten times with PBS 0.1% NP40. Samples were analyzed by SDS-PAGE under nonreducing conditions on a 6% polyacrylamide gel and radiolabeled bands visualized




Fig. 1. FACS analysis of $\alpha_v\beta_3$ and $\alpha_v\beta_5$ integrins on mesangial cells. Suspended (10^5) mesangial cells were labeled by indirect immunofluorescence and analyzed by flow cytometry, as described in the **Methods** section, with monoclonal antibodies to integrins: **(A)** anti- α_v (AMF/7 and LM142), anti- β_3 (AP3), anti- $\alpha_v\beta_3$ (LM609) and anti- $\alpha_v\beta_5$ (PIF6) antibodies; and **(B)** anti- α_2 (Gi9), anti- α_5 (16, 11), anti- α_6 (GoH3) and anti- β_1 (K 20, Li1/2) antibodies. Cells were then stained with fluorescein-conjugated goat anti-mouse IgG (1:1000 dilution) and analyzed by flow cytometry. The ordinates depicts the number of cells per channel and the abscissa depicts the relative fluorescence intensity in arbitrary units (log scale). Control mouse monoclonal IgG was used as control. Representative experiment of six independent determinations.

by autoradiography. The molecular mass markers were myosine (200 kDa), β -galactosidase (116 kDa), phosphorylase-b (97 kDa), bovine serum albumin (66 kDa) and ovalbumin (45 kDa) (Biorad Lab., Paris, France).

Western blotting

HMC were solubilized with 2% NP40 in PBS with the same protease inhibitors as above for 30 minutes at 4°C. Proteins were




Fig. 2. Immunoprecipitation of α_v integrins and CD44 by human mesangial and endothelial cells in culture. Subconfluent cultures of mesangial cells (A) and confluent cultures of endothelial cells (B) were surface radiolabeled, lysed with detergent and subjected to immunoprecipitation as described in the **Methods** section, with monoclonal antibodies directed to the α_v chain (clone LM 142), β_3 chain (clone AP3), β_1 chain (K20) or to the complex $\alpha_v\beta_3$ (LM 609) and $\alpha_v\beta_5$ (P1F6) and to CD44 (J-173) used as control [50]. The precipitated material (HMC, 10^6 cells; HUVEC, 4×10^6 cells) was analyzed by SDS-PAGE under non-reducing conditions and exposed to x-ray film as described. Representative gel of three independent immunoprecipitations.

analyzed by electrophoresis on 7.5% polyacrylamide gel under non-reducing conditions, and transferred to a nitrocellulose membrane by electrophoresis. After saturation overnight at room temperature with TBST (10 mM Tris-HCl, pH 8; 1 mM NaN_3 ; 0.05% Tween 20; 0.15 M NaCl) containing 10% skimmed milk, membranes were incubated one hour at room temperature with anti-integrin β_3 or anti- β_5 mAbs, followed by biotinylated anti-mouse or -rabbit IgG respectively, washed three times with TBST and incubated with HRP-streptavidin complex (Amersham, Buckinghamshire, UK). The membranes were finally revealed by chemiluminescence using an Amersham enhancing reagent according to the manufacturer's instruction.

Statistical analysis

Results of cell adhesion inhibition assays were expressed as means \pm SEM of adherent cells (OD at 630 nm). Groups of duplicates were analyzed. Comparisons were made by ANOVA and post hoc comparisons were made by the Dunnett's test.

Results

Membrane expression

Surface expression of integrin $\alpha_v\beta_3$ and $\alpha_v\beta_5$ was examined by flow cytometry on suspended cells using monoclonal antibodies to




Fig. 3. Western blot analysis indicates that β_3 and β_5 chains are present on mesangial cells. HMC, 10^6 cells, (Mes); HUVEC, 4×10^6 cells (End); human cultured fibroblasts, 10^6 cells (Fib) were solubilized and the blots were probed with antibodies to β_3 (AP3), β_5 chain of integrins (or with control monoclonal IgG1 or polyclonal rabbit IgG; data not shown) and performed as described in the **Methods** section. Biotinylated molecular mass markers (*) are analyzed on the two external lanes of the gel. Representative gel of three independent experiments.




Fig. 4. Mesangial cells attachment to various matrix proteins. Mesangial cells were allowed to attach for 60 minutes to 96-well plates, coated with either fibronectin (FN), vitronectin (VN) or vWF, at various concentration (0.1 to 100 μ g/ml). Adhesion assays were performed as described in the **Methods** section. Results are expressed as the percentage of adherent cells compared with the total number of cells offered. Symbols are: (\square) BSA; (\diamond) FN; (\circ) VN; (\triangle) vWF.

the β_3 and α_v chains and to the $\alpha_v\beta_3$ and $\alpha_v\beta_5$ complexes. As shown on Figure 1a, HMC are labeled by these four mAbs, thus showing that mesangial cells express both $\alpha_v\beta_3$ and $\alpha_v\beta_5$ integrins. In addition, β_1 integrins are known to be expressed on mesangial cells [13, 35]. We confirmed these data by showing a positive labeling with antibodies directed against α_2 , α_5 and α_6 chains (Fig. 1b).

Immunoprecipitation and Western blotting

To determine the molecular weight of the $\alpha_v\beta_3$ and $\alpha_v\beta_5$ complexes identified on the cell surface, mesangial cell membranes were radiolabeled with 125 I and immunoprecipitated with specific antibodies to the α_v and β_3 chains and to the $\alpha_v\beta_3$ and $\alpha_v\beta_5$ complexes. The resulting fractions were subjected to PAGE electrophoresis and autoradiography. Endothelial cells were treated in the same conditions, as a control. As shown in Figure 2A, antibodies to the α_v and β_3 chains or to the $\alpha_v\beta_3$ complex immunoprecipitated proteins with molecular weight of 97 kDa and 140 kDa, corresponding to the known molecular weights of β_3 and α_v subunits, respectively. Immunoprecipitation of α_v and β_3 chains from mesangial cells produced bands identical to that observed with endothelial cells (Fig. 2B). In addition to α_v and β_3 chains, the anti- α_v antibody precipitated, on mesangial cells (as well as on endothelial cells), a third band with an apparent molecular weight of 115 kDa, which may correspond to the β_1 chain. The antibody to $\alpha_v\beta_5$ complex, which binds to mesangial cells as assessed by FACS analysis, failed to immunoprecipitate this integrin (Fig. 2A). β_5 chain expression was nevertheless




Fig. 5. Specific inhibition of mesangial cell attachment to fibronectin by anti-integrin α_5 and β_1 antibodies. Ninety-six-well plates were coated with fibronectin at 1 μ g/ml and unbound sites were blocked with BSA. Cells were preincubated for 30 minutes with anti α_5 (clone 16), anti- $\alpha_v\beta_3$ (LM609), anti- α_2 (Gi9), anti- α_3 (P1B5) and anti- α_5 + anti- β_1 (Lia 1/2). Adhesion assay was then performed as described in the **Methods** section. Results are expressed as percentage of adherent cells compared with the control in presence of a murine monoclonal IgG1. Each point represents the mean \pm SEM of the number of experiments (N is indicated for each antibody tested). * $P < 0.05$ versus control.




Fig. 6. Specific inhibition of mesangial cell attachment to vitronectin by anti-integrin $\alpha_v\beta_3$ and $\alpha_v\beta_5$. Vitronectin at 0.5 μ g/ml was placed into 96-well plates and non specific sites were blocked with BSA. Cells were preincubated for 30 minutes with anti- $\alpha_v\beta_3$ (LM609), anti- $\alpha_v\beta_5$ (P1F6), anti- α_5 (16), anti- β_1 (Li1/2). Adhesion assay was performed as described in **Methods**. Results are expressed as percentage of adherent cells compared with the control in presence of murine monoclonal IgG1. Each point represents the mean \pm SEM of the number of experiments (N is indicated for each antibody tested). * $P < 0.05$ versus control, # $P < 0.05$ versus control and versus inhibition by anti- $\alpha_v\beta_3$ alone.




Fig. 7. Specific inhibition of mesangial cell attachment to von willebrand factor by anti-integrin $\alpha_v\beta_3$ and $\alpha_v\beta_5$. Von Willebrand factor at 5 μ g/ml was placed into 96-well plates and nonspecific sites were blocked with BSA. Cells were preincubated for 30 minutes with anti- $\alpha_v\beta_3$ (LM609), anti- $\alpha_v\beta_5$ (PIF6), α_5 (16), β_1 (Li1/2). The adhesion assay was performed as described in the **Methods** section. Results are expressed as percentage of adherent cells compared with the control in presence of murine monoclonal IgG1. Each point represents the mean \pm SEM of the number of experiments (*N* is indicated for each antibody tested). **P* < 0.05 versus control.

demonstrated by Western blot revealed with an polyclonal antibody to the β_5 chain, as shown on Figure 3. The molecular weight of the β_5 chain was analogous to that of β_3 chain, as already known [10]. In this Western blot analysis (Fig. 3), HUVEC and cultured human fibroblasts, used as control cells, expressed respectively $\alpha_v\beta_3$ and both $\alpha_v\beta_3$ and $\alpha_v\beta_5$, as already known [2].

Role of integrins in HMC adhesion to fibronectin, vitronectin and vWf

Mesangial cells were seeded, in the absence of serum, on fibronectin, vitronectin or vWf coated 96-wells plates. Cell adhesion occurred in a dose dependent manner. As shown in Figure 4, 60% of the maximal response was observed at concentrations of about 1 μ g/ml for fibronectin, 0.5 μ g/ml for vitronectin and 5 μ g/ml for vWf. Maximal adhesion was 100% of the offered cells on fibronectin or vitronectin and 80% on vWf. Adhesion experiments performed in the presence of 10% FBS led to similar results to those obtained in the absence of serum. In order to determine which integrins were involved in adhesion to these ligands, we performed inhibition experiments with blocking anti-integrin antibodies. In the following experiments, ascites fluid containing mAbs or purified mAbs were used at a concentration equal or up to five times the concentration, giving the maximal signal in flow cytometry and equal or up to two times the concentration giving the maximal adhesion inhibition in preliminary experiments.

Adhesion to fibronectin. As shown in Figure 5, mesangial cell adhesion to fibronectin was strongly inhibited ($\geq 75\%$ inhibition) by pre-incubation with mAb to the α_5 or to the β_1 chain alone and with a mixture of both mAbs. In contrast, neither mAb to $\alpha_v\beta_3$

complex and $\alpha_v\beta_5$ complex (data not shown) or to α_2 , α_3 and α_6 chains had any effect on cell adhesion to fibronectin. These results demonstrate that mesangial cell adhesion to fibronectin was mainly mediated by the $\alpha_5\beta_1$ integrin.

Adhesion to vitronectin. As shown in Figure 6, mesangial cell adhesion was inhibited by pre-incubation with mAb to $\alpha_v\beta_3$ or to the $\alpha_v\beta_5$ complexes. When used in combination, these two antibody showed a significant additive effect and strongly inhibited cell adhesion by about 75%. In contrast, mAbs to α_5 had no effect on adhesion to vitronectin. Anti- α_2 , α_3 , α_6 chains also had no effect (data not shown). The slight inhibition of cell adhesion by anti- β_1 mAb was not statistically significant and was not additive to that observed with anti- $\alpha_v\beta_3$ and anti- $\alpha_v\beta_5$. These results demonstrate that adhesion of HMC to vitronectin was primarily mediated by $\alpha_v\beta_3$ and $\alpha_v\beta_5$ integrins.

Adhesion to von Willebrand factor. Figure 7 shows that mesangial cell adhesion to von Willebrand factor was reduced by about 46% by pre-incubation with antibody to the $\alpha_v\beta_3$. Antibody to $\alpha_v\beta_5$ complexes resulted in about 40% inhibition of cell attachment. These results demonstrate that mesangial cell adhesion to von Willebrand factor involves $\alpha_v\beta_3$ and $\alpha_v\beta_5$.

Discussion

We have demonstrated by flow cytometry and immunoprecipitation that $\alpha_v\beta_3$, a broadly distributed integrin (particularly on endothelium, smooth muscle cells and fibroblasts [2]), is expressed on HMC and plays a role in mesangial cell adhesion to vitronectin and vWf. The absence of $\alpha_v\beta_3$ -mediated binding to fibronectin, observed in this study with HMC, is analogous to that observed with human umbilical vein endothelial cells [25], but differs from human dermal microvascular endothelial cells, which bind to fibronectin through $\alpha_v\beta_3$ [36]. One may hypothesize that the interaction between fibronectin and $\alpha_v\beta_3$ is not detected on HMC, because its strength is negligible as compared with that of $\alpha_5\beta_1$.

$\alpha_v\beta_5$ is also present on HMC, as shown by flow cytometry. $\alpha_v\beta_5$ was not detected after immunoprecipitation, probably because of the lack of sensitivity of this method and the low amount of $\alpha_v\beta_5$ present on HMC in culture. Conversely, its presence was directly demonstrated by Western blotting after amplification by chemiluminescence. $\alpha_v\beta_5$ on HMC binds mainly to vitronectin, as already shown for other cell types such as fibroblasts and carcinoma cells [2, 10]. We observed that $\alpha_v\beta_5$ on HMC interacts with vWf, since addition of a monoclonal antibody to the $\alpha_v\beta_5$ complex is inhibitory and increases the inhibition of cell attachment observed with a monoclonal antibody to the $\alpha_v\beta_3$ complex. This differs from previous reports, which demonstrated that $\alpha_v\beta_5$ purified from placenta binds to vitronectin, marginally to fibronectin but not to vWf [10], and that a lung adenocarcinoma cell line expressing $\alpha_v\beta_5$ but not $\alpha_v\beta_3$ binds to vitronectin but not to fibronectin or to vWf. This apparent discordance with our results on HMC may be due to the fact that the $\alpha_v\beta_5$ -vWf interaction is weak and not sufficient to promote adhesion to vWf by itself in these models. Our results suggest that $\alpha_v\beta_5$ on HMC interacts with vWf and could play a role by increasing the strength of $\alpha_v\beta_3$ -mediated cell attachment to vWf.

Finally, $\alpha_v\beta_1$ is likely to be present since we observed that the β_1 chain coprecipitated with the α_v chain. $\alpha_v\beta_1$ has a more restricted distribution, limited to few cell types including fibroblasts, than $\alpha_v\beta_3$, and it binds to vitronectin and fibronectin on those cells [2, 9, 37]. By contrast with what was observed on fibroblasts, we

showed that $\alpha_v\beta_1$ present on human cultured mesangial cells is probably marginally important for fibronectin binding, since a monoclonal antibody to the α_5 chain almost completely inhibited the binding to fibronectin, as already shown by others [13]. Furthermore, $\alpha_v\beta_1$ did not play an important role in adhesion to vitronectin since the antibody to the β_1 chain had a limited (and not significant) effect on cell attachment to vitronectin. Thus, this slight and not significant inhibition observed with the anti- β_1 mAb precluded any firm conclusion on the eventual role of β_1 integrins during adhesion to vitronectin.

The presence of integrins of the α_v family on mesangial cells may have a number of physiopathological consequences. Modifications of this integrin expression have been reported in glomerular diseases [38]. During mesangial proliferation in IgA nephropathy, the $\alpha_v\beta_3$ mesangial expression was shown to be increased when using polyclonal Abs [18, 39, 40], or to be decreased when using mAbs (N. Patey, personal communication). In human crescentic GN, we and others have observed that among other integrins, $\alpha_v\beta_3$ was strongly expressed by the epithelial crescentic cells [17, 41]. Mesangial cells are in close contact with the basal site of endothelial cells [42]. vWf is expressed by all renal tissue endothelial cells, is up-regulated during early fibrosis, and colocalized with TSP in vasculitis, diabetic nephropathy, focal sclerosis and membranous GN, but not in minimal change or IgA nephropathies [16]. On the other hand, other ligands of the α_v integrin family are also present within the glomerulus and are modified during glomerulonephritis. Indeed, thrombospondin [16, 43, 44], and osteopontin [24, 45] are synthesized by mesangial cells, and vitronectin colocalizes with the complement membrane attack complex in glomerular diseases [46, 47].

One may hypothesize that, *in vivo*, excretion of the vWf at the basal site of endothelial cells [21] and vitronectin deposition [48] would promote, mainly via an interaction with $\alpha_v\beta_3$, not only cell adhesion but also migration and cells proliferation. Indeed, $\alpha_v\beta_3$ has multiple functions possibly related to this potential mechanism and, in addition to adhesion, migration and cell survival, $\alpha_v\beta_3$ may also play a role in phagocytosis of apoptotic cells in the inflamed glomerulus. This function of $\alpha_v\beta_3$ has been demonstrated for macrophages but remain to be studied in mesangial cells, which are also able to phagocytize apoptotic neutrophils [49].

Acknowledgments

This work was supported by a grant from the "Programme Hospitalier de Recherche Clinique, PHRC". This work was presented, in part, at the ASN meeting, San Diego, Nov 1995 and published in abstract form (*J Am Soc Nephrol* 6:895, 1995). The authors thank Dr. Paul Landais (Biostatistics Department, Necker Hospital, Paris, France), Dr. David Cheresch (Scripps Research Institute, La Jolla, CA), Dr. Peter J. Newman (The Blood Center of Southeastern Wisconsin, USA), Dr. Steven K. Akiyama (NIH, Bethesda, MD, USA) and Dr. Alain Bernard (INSERM U 343, Marseilles, France) for providing monoclonal antibodies; Dr. Nicole Ardaillou (INSERM U64, Paris, France) for her advice on human mesangial cell culture; and Dr. Dominique Droz (Renal Pathology Laboratory, Necker Hospital, Paris, France) for providing kidney specimens.

Reprint requests to Dr. Z. Hafdi, INSERM U 90, Necker Hospital, 161 rue de Sèvres, 75015 Paris, France.

References

1. HYNES RO: Integrins: A family of cell surface receptors. *Cell* 48:549-544, 1987
2. FELDING-HABERMANN B, CHERESH DA: Vitronectin and its receptors. *Curr Opin Cell Biol* 5:864-868, 1993
3. BROOKS PC, CLARK RA, CHERESH DA: Requirement of vascular integrin alpha v beta 3 for angiogenesis. *Science* 264:569-571, 1994
4. FRIEDLANDER M, BROOKS PC, SHAFFER RW, KINCAID CM, VARNER JA, CHERESH DA: Definition of two angiogenic pathways by distinct alpha v integrins. *Science* 270:1500-1502, 1995
5. BROWN SL, LUNDGREN CH, NORDT T, FUJII S: Stimulation of migration of human aortic smooth muscle cells by vitronectin: Implications for atherosclerosis. *Cardiovasc Res* 28:1815-1820, 1994
6. DRAKE CJ, CHERESH DA, LITTLE CD: An antagonist of integrin alpha v beta 3 prevents maturation of blood vessels during embryonic neovascularization. *J Cell Sci* 108:2655-2661, 1995
7. MONTGOMERY AM, REISFELD RA, CHERESH DA: Integrin alpha v beta 3 rescues melanoma cells from apoptosis in three-dimensional dermal collagen. *Proc Natl Acad Sci USA* 91:8856-8860, 1994
8. BROOKS PC, MONTGOMERY AM, ROSENFELD M, REISFELD RA, HU T, KLIER G, CHERESH DA: Integrin alpha v beta 3 antagonists promote tumor regression by inducing apoptosis of angiogenic blood vessels. *Cell* 79:1157-1164, 1994
9. BODARY SC, MCLEAN JW: The integrin β_1 subunit associates with the vitronectin receptor α_v subunit to form a novel vitronectin receptor in a human embryonic kidney cell line. *J Biol Chem* 265:5938-5941, 1990
10. SMITH JW, VESTAL DJ, IRWIN SV, BURKE TA, CHERESH DA: Purification and functional characterization of integrin $\alpha_v\beta_5$. An adhesion receptor for vitronectin. *J Biol Chem* 265:11008-11013, 1990
11. BUSK M, PYTELA R, SHIEPPARD D: Characterization of the integrin $\alpha_v\beta_6$ as a fibronectin-binding protein. *J Biol Chem* 267:569-571, 1992
12. NISHIMURA SL, SHEPPARD D, PYTELA R: Integrin $\alpha_v\beta_8$: Interaction with vitronectin and functional divergence of the β_8 cytoplasmic domain. *J Biol Chem* 269:28708-28715, 1994
13. COSIO FG, SEDMARK DD, NAHMAN NS: Cellular receptors for matrix proteins in normal human kidney and human mesangial cells. *Kidney Int* 38:886-895, 1990
14. KORHONEN M, YLÄNNE J, LAITINEN L, VIRTANAN I: Distribution of β_1 and β_3 integrins in human fetal and adult kidney. *Lab Invest* 62:616-625, 1990
15. PATEY N, HALBWACHS-MECARELLI L, DROZ D, LESAVRE P, NOËL LH: Distribution of integrin subunits in normal human kidney. *Cell Adhes Commun* 2:159-167, 1994
16. MCGREGOR B, COLON S, MUTIN M, CHIGNIER E, ZECH P, MCGREGOR J: Thrombospondin in human glomerulopathies. A marker of inflammation and early fibrosis. *Am J Pathol* 144:1281-7, 1994
17. BARALDI A, ZAMBRUNO G, FURCI L, BALLESTRI M, TOMBESI A, OTTANI D, LUCCHI L, LUSVARGHI E: β_1 and β_3 integrin upregulation in rapidly progressive glomerulonephritis. *Nephrol Dial Transplant* 10:1155-1161, 1995
18. SHIKATA K, MAKINO H, MORIOKA S, KASHITANI T, HIRATA K, OTA Z, WADA J, KANWAR YS: Distribution of extracellular matrix receptors in various forms of glomerulonephritis. *Am J Kidney Dis* 25:680-688, 1995
19. WADA J, KUMAR A, LIU Z, RUOSHLATI E, REICHHARDT L, MARVALDI J, KANWAR YS: Cloning of mouse integrin α_v cDNA and role of the α -related matrix receptors in metanephric development. *J Cell Biol* 132:1161-1176, 1996
20. TUSYNSKI GP, KOWALSKA MA: Thrombospondin-induced adhesion of human platelets. *J Clin Invest* 87:1387-1394, 1991
21. WAGNER D, MARDER VJ: Biosynthesis of von Willebrand protein by human endothelial cells: Processing steps and their intracellular localization. *J Cell Biol* 99:2123-2129, 1984
22. AKIYAMA S, YAMADA S, CHEN W, YAMADA K: Analysis of fibronectin receptor function with monoclonal antibodies: Roles in cell adhesion, migration, matrix assembly and cytoskeletal organization. *J Cell Biol* 109:863-868, 1989
23. SONNENBERG AH, DAAM MA, VAN DER VALK J, HILKEN J, HILGERS J: Development of mouse mammary gland: identification of stages in differentiation of luminal and myoepithelial cells using monoclonal antibodies and polyvalent antiserum against keratin. *J Histochem Cytochem* 34:1037-1046, 1986
24. LIAW L, SKINNER MP, RAINES EW, ROSS R, CHERESH DA, SCHWARTZ

- SM, GIACHELLI CM: The adhesive and migratory effects of osteopontin are mediated via distinct cell surface integrins. Role of alpha v beta 3 in smooth muscle cell migration to osteopontin *in vitro*. *J Clin Invest* 95:713-724, 1995
25. CHERESH DA: Human endothelial cells synthesize and express an Arg-Gly-Asp-directed adhesion receptor involved in attachment to fibrinogen and von Willebrand factor. *Proc Natl Acad Sci USA* 84:6471-6475, 1987
 26. CAMPANERO MR: Functional role of α 2 β 1 integrin in leucocyte intercellular adhesion induced through the common β 1 subunit. *Eur J Immunol* 22:3111-3119, 1992
 27. AMIOT M, BERNARD A, TRAN H, LECA G, KANELLOPOULOS JM, BOUMSELL L: The human cell surface glycoprotein complex (gp 120,200) recognized by monoclonal antibody K20 is a component binding to phytohaemagglutinin on T cells. *J Immunol* 23:109-118, 1986
 28. NEWMAN PJ, ALLEN RW, KAHN RA, KUNICKI TJ: Quantitation of membrane glycoprotein IIIa on intact human platelets using the monoclonal antibody, AP-3. *Blood* 65:227-232, 1985
 29. ADAMS JC, WATT FM: Expression of β 1, β 3, β 4, and β 5 integrins by human epidermal keratinocytes and non-differentiating keratinocytes. *J Cell Biol* 115:829-841, 1991
 30. GAILLIT J, CLARK RAF: Studies *in vitro* on the role of α v and β 1 integrins in the adhesion of human dermal fibroblasts to provisional matrix proteins fibronectin, vitronectin, and fibrinogen. *J Invest Derm* 106:102-108, 1996
 31. THORELL L, BLOMBACK B: Purification of the factor VIII complex. *Thromb Res* 35:431-437, 1984
 32. STRIKER GE, STRIKER LJ: Glomerular cell culture. *Lab Invest* 53:122-131, 1985
 33. JAFFE EA, NACHMAN RL, BECKER CG, MINICK CR: Culture of human endothelial cells derived from umbilical veins. Identification by morphologic and immunologic criteria. *J Clin Invest* 52:2745-2750, 1973
 34. CONFORTI G, DOMINGUEZ-JIMENEZ C, ZANETTI A, GIMBRONE JR MA, CREMONA O, MARCHISIO PC, DEJANA E: Human endothelial cells express integrin receptors on the luminal aspects of their membrane. *Blood* 80:437-446, 1992
 35. PETERMANN A, FEES H, GRENZ H, GOODMAN SL, STERZEL RB: Polymerase chain reaction and focal contact indicate integrin expression in mesangial cells. *Kidney Int* 44:997-1005, 1993
 36. CHENG YF, CLYMAN RI, ENENSTEIN J, WALEH N, PYTELA R, KRAMER RH: The integrin complex alpha v beta 3 participates in the adhesion of microvascular endothelial cells to fibronectin. *Exp Cell Res* 184:69-77, 1991
 37. ZHANG Z, MORLA AO, VUORI K, BAUER JS, JULIANO RL, RUOSLAHTI E: The alpha v beta 1 integrin functions as a fibronectin receptor but does not support fibronectin matrix assembly and cell migration on fibronectin. *J Cell Biol* 122:235-242, 1993
 38. COSIO FG: Cell-matrix adhesion receptors: Relevance to glomerular pathology. *Am J Kidney Dis* 20:294-305, 1992
 39. OKADA M, YOSHIOKA K, TAKEMURA T, AKANO N, AYA N, MURAKAMI K, MAKI S: Immunohistochemical localization of C3 days fragment of complement and S-protein (vitronectin) in normal and diseased human kidneys: Association with the C5b-9 complex and vitronectin receptor. *Virchows Arch A Pathol Anat Histopathol* 422:367-373, 1993
 40. KANAHARA K, YORIOKA N, ARITA M, OHIRA N, YAMAKIDO M: Immunohistochemical studies of extracellular matrix components and integrins in IgA nephropathy. *Nephron* 66:29-37, 1994
 41. PATEY N, LESAVRE P, HALBWACHS-MECARELLI L, NOËL L H: Adhesion molecules in human crescentic glomerulonephritis. *J Pathol* 179:414-420, 1996
 42. KRIZ W, ELGER M, MUNDEL P, LEMLEY KV: Structure-stabilizing forces in the glomerular tuft. *J Am Soc Nephrol* 5:1731-1739, 1995
 43. TARABOLLETTI G, MORIGI M, FIGLIUZZI M, GIAVAZZI R, ZOJA C, REMUZZI G: Thrombospondin induces glomerular mesangial cell adhesion and migration. *Lab Invest* 67:566-571, 1992
 44. TSUKAMOTO M, TADA H: Thrombospondin modulates cell attachment, spreading and proliferation in cultured rat mesangial cells. *Nippon Junzo Gakkai Shi* 7:791-795, 1993
 45. LOPEZ CA, HOYER JR, WILSON PD, WATERHOUSE P, DENHARDT DT: Heterogeneity of osteopontin expression among nephrons in mouse kidneys and enhanced expression in sclerotic glomeruli. *Lab Invest* 69:355-363, 1993
 46. FALK RJ, PODACK E, DALMASSO AP, JENNETTE JC: Localization of S protein and its relationship to membrane attack complex of complement in renal tissue. *Am J Pathol* 127:182-190, 1987
 47. REILLY JT, NASH JRG: Vitronectin (serum spreading factor): Its localisation in normal and fibrotic tissue. *J Clin Pathol* 41:1269-1272, 1988
 48. PREISSNER KT, POTZSCH B: Vessel wall-dependent metabolic pathways of the adhesive proteins, von-Willebrand-factor and vitronectin. *Histol Histopathol* 10:239-251, 1995
 49. SAVILL J, SMITH J, SARRAF C, REN Y, ABBOTT F, REES A: Glomerular mesangial cells and inflammatory macrophages ingest neutrophils undergoing apoptosis. *Kidney Int* 42:924-936, 1992
 50. LESAVRE P, CHOUKROUN G, FRANÇOIS A, NOËL LH, HALBWACHS-MECARELLI L, DROZ D: CD44, a hyaluronan-binding molecule: Expression in normal kidneys and by cultured mesangial cells, in *Advances in Nephrology*, Chicago, Year Book Medical Publishers, 1966, pp 317-339