



Editorial: Extracellular Vesicles in CNS Diseases

Raffaella Soleti, Maria Antonietta Panaro, Tarek Benameur, Giovanni Messina, Chiara Porro

► To cite this version:

Raffaella Soleti, Maria Antonietta Panaro, Tarek Benameur, Giovanni Messina, Chiara Porro. Editorial: Extracellular Vesicles in CNS Diseases. *Frontiers in Molecular Biosciences*, 2022, 9, pp.943369. 10.3389/fmolb.2022.943369 . inserm-03823990

HAL Id: inserm-03823990

<https://inserm.hal.science/inserm-03823990>

Submitted on 21 Oct 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Editorial: Extracellular Vesicles in CNS Diseases

Raffaella Soleti^{1*}, Maria Antonietta Panaro^{2*}, Tarek Benameur^{3*}, Giovanni Messina^{4*} and Chiara Porro^{4*}

¹Univ Angers, Nantes Université, Inserm, CNRS, CRCI2NA, SFR ICAT, Angers, France, ²Department of Biosciences, Biotechnologies and Biopharmaceutics, University of Bari, Bari, Italy, ³Department of Biomedical Sciences, College of Medicine, King Faisal University, Al-Ahsa, Saudi Arabia, ⁴Department of Clinical and Experimental Medicine, University of Foggia, Foggia, Italy

Keywords: extracellular vesicles, CNS diseases, amyotrophic lateral sclerosis, miRNA - microRNA, exosomes

Editorial on the Research Topic

Extracellular Vesicles in CNS Diseases

OPEN ACCESS

Edited and reviewed by:

Cecilia Giulivi,
University of California, Davis,
United States

*Correspondence:

Raffaella Soleti
raffaella.soleti@univ-angers.fr
orcid.org/00000002-0271-0490
Maria Antonietta Panaro
mariaantonietta.panaro@uniba.it
orcid.org/0000-0001-5457-1069
Tarek Benameur
tbenameur@kfu.edu.sa
orcid.org/0000-0001-8683-2526
Giovanni Messina
giovanni.messina@unifg.it
orcid.org/0000-0002-0011867X
Chiara Porro
chiara.porro@unifg.it
orcid.org/0000-0002-7526-6968

Specialty section:

This article was submitted to
Cellular Biochemistry,
a section of the journal
Frontiers in Molecular Biosciences

Received: 13 May 2022

Accepted: 16 May 2022

Published: 09 June 2022

Citation:

Soleti R, Panaro MA, Benameur T,
Messina G and Porro C (2022)
Editorial: Extracellular Vesicles in
CNS Diseases.
Front. Mol. Biosci. 9:943369.
doi: 10.3389/fmolb.2022.943369

Extracellular vesicles (EVs) are considered as a new tool for cell to-cell communication in the brain, in both brain health and diseases. Under physiological and pathological conditions, all cells release EVs. Based on their biogenesis and size, EVs can be divided into three categories: exosomes, micro-vesicles, and apoptotic bodies. Many body fluids contain EVs, which enclose membrane, cytosolic membrane, and nuclear components such as proteins, lipids, and a variety of RNAs, including microRNAs, other non-coding RNA species with regulatory properties, as well as DNA fragments. EVs carry a variety of bioactive molecules and act as inter- and intra-cellular messenger, representing the fingerprints of their originating cells. EVs have the potential to influence target cell fate, inducing significant changes that may lead to their contribution to different pathophysiological processes. In the central nervous system (CNS) diseases, EVs and their content have emerged as potential biomarkers for diagnosis and prognosis (Panaro et al., 2020; Benameur et al., 2022).

The purpose of the present Research Topic (RT) is to bring together recent findings in the field of EVs in the brain. EVs have been studied for their role in cellular communication, but many questions remain unanswered, including those about EVs biogenesis, cargo characterization, and the role of EVs in the brain, with particular focus on their contribution to the progression of neurodegenerative diseases.

A better understanding of the pleiotropic role of EVs in brain homeostasis could help researchers to gain the knowledge they need to develop new therapeutic strategies to treat and reverse neurological diseases.

A total of 26 high-ranking international researchers and scientists have contributed to a compilation of four articles in the current Research Topic. Among the four accepted articles, readers will find one original research article (Yamaguchi et al.), two reviews (Chen et al.; Wang et al.) and one mini review (Nakamya et al.).

Modifications of protein cargo and nucleic acid from exosomes isolated from human biological fluids have been linked to several pathologies, including amyotrophic lateral sclerosis (ALS). ALS is a serious neurodegenerative disease characterized by a progressive motor neuron degeneration (MN) that results in muscle control loss. The majority of patients die of respiratory complications within 3–5 years of diagnosis with ALS (Cleveland and Rothstein, 2001; Chiò et al., 2009; Brown and Al-Chalabi, 2017; Hosaka et al., 2019; Roy et al., 2019). Regrettably, the molecular pathways involved in motor neuron degeneration remain unknown. Indeed, the research for useful diagnostic biomarkers and treatment strategies is difficult, despite the fact that genetic and non-genetic causes have been proposed. Exosomes have been identified as promising candidates for studying ALS pathogenesis, as they can act as biomarkers for disease diagnostic and progression. In their review, Chen et al. revisited a summary of existing knowledge on exosome the biogenesis and integration, as well as their function and heterogeneity in the brain. They demonstrated how ALS-related proteins are secreted and propagate through exosomes, probably contributing to progressive disease dissemination throughout the body. They also illustrated several extracellular miRNAs associated with the onset and progression of ALS, as well as the therapeutic potentials of exosomes, despite the lack of

standardized methodology for exosomes detection and isolation. Wang et al. focused their review on the use of EVs as therapeutic strategies for ALS. Exosomes can cross the Blood Brain Barrier (Ediriweera et al., 2021), this characteristic making them a potential candidate for non-cell therapy in neurodegenerative diseases, such as ALS (Ciregia et al., 2017). EVs can be isolated from a variety of cells of different origins. Mesenchymal stem cells have the capacity to release a higher amount of EVs that have beneficial effects on their target cells. Indeed, the authors demonstrated promising therapeutic strategies against ALS using of EVs derived from adipose-derived stem cells and human bone marrow-derived endothelial progenitor cells.

The authors then discussed the use of EVs as drug delivery tool for ALS. Chemical, biological, or physical methods can be used to enrich EVs with therapeutic drugs or bioeffectors (Bonafede et al., 2016). Finally, authors explored other interesting tools as potential therapeutics: plant-derived EVs and exosome-mimics, either unmodified or engineered. The innovative research on EVs represents an encouraging strategy for the treatment of ALS.

Yamaguchi et al. elegantly investigate the role of EVs in Hereditary Transthyretin Amyloidosis (ATTRy), a disease in which insoluble proteins, known as amyloids, are deposited in various organs, causing multiple damage. They showed that cell-derived EVs were involved in the formation of transthyretin (TTR) amyloid deposits on the membrane of small EVs and cells. Similarly, human serum EVs altered the intensity of TTR aggregation and deposition, resulting in lower TTR in serum-derived EVs from ATTR patient. The increased amount of EVs in ATTR patients suggested that they actively contribute to enhance ATTR amyloidosis and metabolism. As a result, serum-derived small EVs offer an opportunity to develop diagnostic and therapeutic tools for patients with ATTR amyloidosis.

To complete the analysis of the role of EVs in the CNS, Nakamya et al. present a synthesis of current knowledge about extracellular

vesicles carrying mitochondrial components, with a focus on several neurological disorders of the CNS such as Parkinson's Disease, Down Syndrome, Alzheimer's Diseases and aging. In their minireview the authors also discuss the current findings that highlight cross talk between mitophagy and the EVs that carry mitochondrial cargo. The release of mitochondria-derived vesicles (MDVs), a subtype of EVs containing mitochondrial DNA and proteins, is triggered by mitochondrial dysfunction due to dysregulated mitophagy and the induction of oxidative stress.

The MDV content could be disseminated. Consequently, MDVs contribute to disease pathogenesis and progression. MDVs, on the other hand, aid in the preservation of cell functions by promoting the elimination of their content in lysosomal compartments. MDVs, according to the authors, could be used as biomarkers or as tools in the development of novel therapeutic strategies for CNS disorders. Indeed, it is critical to develop research projects to broaden and standardize methodology for EVs studies, as well as to further investigate the functions and mechanisms involved in CNS disorders. Altogether, this collection of reviews and original researches for this Research Topic provides new perspectives and insights into the diverse links between EVs and their content, as well as their potential value as biomarkers of diagnosis and prognosis in CNS diseases.

AUTHOR CONTRIBUTIONS

RS, MAP, TB, GM, and CP wrote the editorial.

ACKNOWLEDGMENTS

We would like to thank to the contributors to this Research Topic as well as to all the referees and the editorial support of the Frontiers Team members.

REFERENCES

- Benaneur, T., Panaro, M. A., and Porro, C. (2022). Exosomes and Their Cargo as a New Avenue for Brain and Treatment of CNS-Related Diseases. *Open Neurology Journal* 16, 1190. doi:10.2174/1874205x-v16-e2201190
- Bonafede, R., Scambi, I., Peroni, D., Potrich, V., Boschi, F., Benati, D., et al. (2016). Exosome Derived from Murine Adipose-Derived Stromal Cells: Neuroprotective Effect on *In Vitro* Model of Amyotrophic Lateral Sclerosis. *Exp. Cell Res.* 340, 150–158. doi:10.1016/j.yexcr.2015.12.009
- Brown, R. H., and Al-Chalabi, A. (2017). Amyotrophic Lateral Sclerosis. *N. Engl. J. Med.* 377, 162–172. doi:10.1056/nejmra1603471
- Chiò, A., Logroscino, G., Hardiman, O., Swigler, R., Mitchell, D., Beghi, E., et al. (2009). Prognostic Factors in ALS: A Critical Review. *Amyotroph. Lateral Scler.* 10, 310–323. doi:10.3109/17482960802566824
- Ciregia, F., Urbani, A., and Palmisano, G. (2017). Extracellular Vesicles in Brain Tumors and Neurodegenerative Diseases. *Front. Mol. Neurosci.* 10, 276. doi:10.3389/fnmol.2017.00276
- Cleveland, D. W., and Rothstein, J. D. (2001). From Charcot to Lou Gehrig: Deciphering Selective Motor Neuron Death in ALS. *Nat. Rev. Neurosci.* 2, 806–819. doi:10.1038/35097565
- Ediriweera, G. R., Chen, L., Yerbury, J. J., Thurecht, K. J., and Vine, K. L. (2021). Non-Viral Vector-Mediated Gene Therapy for ALS: Challenges and Future Perspectives. *Mol. Pharm.* 18, 2142–2160. doi:10.1021/acs.molpharmaceut.1c00297
- Hosaka, T., Yamashita, T., Tamaoka, A., and Kwak, S. (2019). Extracellular RNAs as Biomarkers of Sporadic Amyotrophic Lateral Sclerosis and Other

Neurodegenerative Diseases. *Int. J. Mol. Sci.* 20 (13), 3148. doi:10.3390/ijms20133148

Panaro, M. A., Benaneur, T., and Porro, C. (2020). Extracellular Vesicles miRNA Cargo for Microglia Polarization in Traumatic Brain Injury. *Biomolecules* 10 (6), 901. doi:10.3390/biom10060901

Roy, J., Saucier, D., O'Connell, C., and Morin, P. J. (2019). Extracellular Vesicles and Their Diagnostic Potential in Amyotrophic Lateral Sclerosis. *Clin. Chim. Acta* 497, 27–34. doi:10.1016/j.cca.2019.07.012

Conflict of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Publisher's Note: All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Copyright © 2022 Soleti, Panaro, Benaneur, Messina and Porro. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.