

Imaging biomarker discovery in major depressive disorder with diffusion MRI multi-compartment models

Renaud Hedouin, Olivier Commowick, Gabriel Robert, Julie Coloigner

▶ To cite this version:

Renaud Hedouin, Olivier Commowick, Gabriel Robert, Julie Coloigner. Imaging biomarker discovery in major depressive disorder with diffusion MRI multi-compartment models. OHBM 2022 - 28th Annual Meeting of the Organization of Human Brain Mapping, Jun 2022, Glasgow, United Kingdom. pp.1-1. inserm-03934921

HAL Id: inserm-03934921 https://inserm.hal.science/inserm-03934921v1

Submitted on 11 Jan 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Imaging biomarker discovery in major depressive disorder with diffusion MRI multi-compartment models



R. Hedouin¹, O. Commowick¹, G. Robert^{1,2}, J. Coloigner¹

¹ INRIA Rennes - Bretagne Atlantique, VISAGES Research Team

² Adult University Psychiatry Department, Guillaume Régnier Hospital, Rennes, France

Introduction

- Major depressive disorder (MDD) is a disease widespread all over the world associated with a large and increasing economic, societal, and personal burden [Ferrari 2013]
- Despite a large number of studies, the pathophysiology underlying MDD is not well understood
- We propose a tract-based pipeline analysis using multicompartment models (MCM) that allows to model water diffusion under the voxel resolution

Data

- Actidep: Major depressive disorder in the elderly population 40 patients and 28 controls
- Clinical interest: Depressive symptoms in late life are associated with an increased risk of developing dementia [Barnes 2012]
- MRI acquisitions
 - Cusp: DWI with 1.5 mm isotropic resolution, 60 gradient orientations and 7 b0 [Scherrer 2012]
 - MPRAGE: T1-weighted images 1 mm isotropic
 - Resting state fMRI

Methods

Input:

- DWI with several b-values to estimate complex models
- T1-weighted image for registration purpose

DWI preprocessing

Eddy current correction


Block-matching distortion correction

NL-means denoising

Brain masking

Multi-compartments models

Estimate MCM [Stamm 2016] that decomposes the water diffusion as a weighted sum of anisotropic and isotropic compartments



Extract scalar maps (FA, MD, FW ...) from MCMs



MCM FA is the average of each anisotropic compartment FA



DTI FA

 Rigid registration of preprocessed DWIs and scalar maps on MNI152 Atlas


Tractography

For each subject:

- TractSeg performs a tractography for all registered DWIs based on ODF resulting in 72 identified fiber bundles [Wasserthal 2018]
- For each bundle a centroid is estimated and values (FA, MD, ...) are provided [Chandio 2020]

Statistics


- Performs group to group comparison along tracts
- Covariable (age, sexe ...)
 can be removed





Results


Statistic along fiber tracts

- Pipeline tested on the Actidep database
- Group controls (26 subjects) vs group patients (39 subjects) for different MCM metrics
- Corrected for age, sex and duration of depression









• Expected highlight within the corpus callosum

Conclusion & perspectives

- We developed a pipeline to perform tractography analysis using multi-compartments models
- We tested this pipeline on the Actidep database
- In the future we will extend this study to explore different database with additional MRI sequences, such as relaxometry, to derive new metrics and develop a multi-modal approach

Ferrari, Alize J., et al. "Burden of depressive disorders by country, sex, age, and year: findings from the global burden of disease study 2010."

Stamm, Aymeric, et al. "A Bayes Hilbert Space for Compartment Model Computing in Diffusion MRI, 2018.. Wasserthal, Jakob, et al. "TractSeg-Fast and accurate white matter tract segmentation «

Chandio, Bramsh Qamar, et al. "Bundle analytics, a computational framework for investigating the shapes and profiles of brain pathways across populations."

Scherrer, Benoit, and Simon K. Warfield. "Parametric representation of multiple white matter fascicles from cube and sphere diffusion MRI."

Barnes, Deborah E., et al. "Midlife vs late-life depressive symptoms and risk of dementia: differential effects for Alzheimer disease and vascular dementia."