



**HAL**  
open science

## AA Amyloidosis Secondary to Primary Immune Deficiency: About 40 Cases Including 2 New French Cases and a Systematic Literature Review

Marion Delplanque, Lionel Galicier, Eric Oziol, Stéphanie Ducharme-Bénard, Eric Oksenhendler, David Buob, Gilles Grateau, David Boutboul, Sophie Georgin-Lavialle

### ► To cite this version:

Marion Delplanque, Lionel Galicier, Eric Oziol, Stéphanie Ducharme-Bénard, Eric Oksenhendler, et al.. AA Amyloidosis Secondary to Primary Immune Deficiency: About 40 Cases Including 2 New French Cases and a Systematic Literature Review. *Journal of Allergy and Clinical Immunology: In Practice*, 2021, 9 (2), pp.745-752.e1. 10.1016/j.jaip.2020.09.023 . inserm-04050271

**HAL Id: inserm-04050271**

**<https://inserm.hal.science/inserm-04050271>**

Submitted on 22 Jul 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Copyright

1 **AA amyloidosis secondary to primary immune deficiency: about 40 cases including 2 new**  
2 **French cases and a systematic literature review.**

3

4 **Authors**

5 Marion Delplanque<sup>1</sup> MD, Lionel Galicier<sup>2</sup> MD, Eric Ozio<sup>3</sup> MD, Stéphanie Ducharme-Bénard<sup>4</sup>  
6 MD, Eric Oksenhendler<sup>2</sup> MD, PhD, David Buob<sup>5</sup> MD, PhD, Gilles Grateau<sup>1</sup> MD, PhD, David  
7 Boutboul<sup>2\*</sup> MD, Sophie Georgin-Lavialle<sup>1\*</sup> MD, PhD

8 **Affiliations:**

9 1 Sorbonne Université, AP-HP, Hôpital Tenon, Service de médecine interne, Centre de référence  
10 des maladies auto-inflammatoires et des amyloses d'origine inflammatoire (CEREMAIA), GRC-  
11 28 (Groupe de recherche clinique amylose AA Sorbonne université), F-75020, Paris

12 2 Service d'Immunopathologie Clinique, Hôpital Saint Louis, AP-HP, Paris

13 3 Service de Médecine Interne, Centre Hospitalier de Béziers, Béziers

14 4 Service de Médecine Interne, Hôpital du Sacré-Cœur de Montréal, Montréal, Québec, Canada

15 5 Sorbonne Université, AP-HP, Hôpital Tenon, Service d'Anatomie et Cytologie pathologiques,  
16 F-75020, Paris

17 \*Both authors co directed the study

18

19 **Corresponding author:**

20 Sophie Georgin-Lavialle, MD, PhD

21 Sorbonne Université, AP-HP, Hôpital Tenon, Service de médecine interne, Centre de référence  
22 des maladies auto-inflammatoires et des amyloses d'origine inflammatoire (CEREMAIA), 75020,  
23 Paris.

24 Tel: 00 331 56 01 60 77 Fax: 00 33 1 56 01 71 46 E-Mail: [sophie.georgin-lavialle@aphp.fr](mailto:sophie.georgin-lavialle@aphp.fr)

25

26 **Authors disclose any financial relationship with a biotechnology and/or pharmaceutical**  
27 **manufacturer that has an interest in the subject matter or materials discussed in the**  
28 **submitted manuscript.** Each author has no conflict of interest to declare concerning this manuscript.

29 **Word count: Manuscript: 3075 words; Abstract: 243 words**

30

31 **Highlights box:**

32 1. What is already known about this topic?

33 **AAA is a long term complication of PID with a poor prognosis**

34 2. What does this article add to our knowledge?

35 **If AAA patient displays bronchiectasis, PID should be sought, especially humoral**  
36 **deficiencies.**

37 3. How does this study impact current management guidelines?

38 **Renal function and proteinuria monitoring should be monitored among PID patients**

39

40

41 **Key words: Primary Immune deficiency, Common variable immunodeficiency, Humoral**  
42 **deficiency, Long term complication, AA Amyloidosis, nephrotic syndrome, prognosis,**  
43 **chronic infections, bronchiectasis, systematic review**

44

45

46

47 **Abbreviations**

48 AAA: AA Amyloidosis

49 BK: tuberculosis

50 CEREDIH: Reference Center for Hereditary Immune Deficiencies

51 CEREMAIA: Reference Center for Auto-Inflammatory Diseases and AA Amyloidosis

52 CGD: Chronic Granulomatous Disease

53 CRP: C-reactive protein

54 CT scan: Computerized tomography scan

55 CTC: Corticosteroids

56 CVID: Common variable immunodeficiency

57 DMARD: Disease-Modifying Antirheumatic Drugs

58 ESR: Erythrocyte sedimentation rate

59 F: Female

60 FMF: Familial Mediterranean Fever

61 GI: Gastrointestinal

62 HIES: Hyper IgE Syndrome

63 HIV: Human Immunodeficiency Virus

64 HPV: Human Papillomavirus

65 HSCT: Hematopoietic Stem Cell Transplantation

66 HSV: Herpes Simplex Virus

67 IM: Intramuscular

68 IRT: Immunoglobulin Replacement Therapy

69 IV: intravenous

70 LRBA: LPS Responsive Beige-like Anchor protein

- 71 M: Male
- 72 PAD: Predominantly Antibody Deficiency
- 73 PID: Primary Immune Deficiency
- 74 PRISMA: Preferred Reporting Items for Systematic Reviews and Meta-Analyses
- 75 RA: Rheumatoid arthritis
- 76 SAA: Serum Amyloid A
- 77 SC: Subcutaneous
- 78 TLR: Toll-Like Receptor
- 79 TTE: TransThoracic Echocardiography
- 80 VZV: Varicella-Zoster Virus
- 81 XLA: X-linked agammaglobulinemia
- 82 Yrs: Years
- 83

84 **Abstract (243 words)**

85 **Background:** Primary immune deficiencies (PID) are a heterogeneous group of disorders  
86 resulting from defects in immune system. They lead to increased susceptibility to infections and  
87 immune dysregulation. The resulting chronic inflammation can induce long-term complications,  
88 including AA amyloidosis (AAA).

89  
90 **Objectives:** To present the French cases of PID-related AAA and perform a systematic literature  
91 review to determine its main features and predisposing factors.

92  
93 **Methods:** A systematic literature review was performed by searching MEDLINE up until 2019.  
94 New French cases were identified with the help of the Reference Center for Auto-Inflammatory  
95 Diseases and AA Amyloidosis (CEREMAIA) and the Reference Center for Hereditary Immune  
96 Deficiencies (CEREDIH).

97  
98 **Results:** Forty patients were identified including 2 new French cases. PID were varied:  
99 immunoglobulin deficits (n=30), chronic granulomatous disease (n=3), Hyper IgM syndrome  
100 (n=3), hereditary complete C4 deficiency (n=1), Leucocyte Adhesion Deficiency Type-1 (n=1),  
101 Hyper IgE syndrome (n=1) and Chediak-Higashi syndrome (n=1). Mean age at PID diagnosis  
102 was  $22.2 \pm 16.02$  years. Renal involvement was the most common manifestation of AAA (80%).  
103 Infections were extremely heterogenous; bacterial infection with pulmonary involvement was the  
104 most frequent. Bronchiectasis was particularly common (52,5%). The delay between the first  
105 symptoms of PID and AAA diagnosis was  $16.18 \pm 7$  years. Thirteen concomitant diagnoses were  
106 made. Twenty patients died during follow-up.

107  
108 **Conclusion:** AAA is a rare life-threatening complication of PID, especially in cases of long  
109 diagnostic and therapeutic delays. Bronchiectasis should be considered as a warning sign of  
110 chronic inflammation and increased risk of AAA.

111

112

113 **Introduction**

114 Primary immune deficiencies (PID) are a heterogeneous group of diseases. While several  
115 classification systems exist, the one of the International Union of Immunological Societies (IUIS)  
116 expert committee for Primary Immunodeficiency is the most frequently used (1,2). PID are  
117 defined by an immune system failure with no underlying infectious, neoplastic or iatrogenic  
118 cause (3), leading to increased susceptibility to infection and immune dysregulation. The  
119 prevalence of PID in the general population is not well established, and probably underestimated  
120 because of the heterogeneity of these diseases. While severe and potentially life-threatening PID  
121 usually begin in the first weeks of life, others declare themselves later, including in adulthood.

122 More than 300 genes are associated with PID, causing mono or multigenic genetic abnormalities  
123 (4). Common variable immunodeficiency (CVID) is a type of PID generally affecting adults and  
124 characterized by impaired B cell differentiation resulting in hypogammaglobulinemia, normal or  
125 low numbers of B cells, and poor antibody response (5).

126 AA amyloidosis (AAA) is a multisystemic disease related to the deposition of serum amyloid A  
127 (SAA) protein secondary to chronic inflammation. Causes are multiple, the most frequent ones  
128 being chronic rheumatic and inflammatory bowel diseases, monogenic autoinflammatory  
129 diseases and chronic infections. However, immune deficiencies can also lead to AAA by  
130 predisposing to infection (6,7). Chronic and recurrent infections as well as immune dysregulation  
131 in patients with PID may lead to chronic inflammation, favoring extracellular deposition of SAA.  
132 Numerous assumptions exist about predisposing conditions of AAA complicating PID: severity,  
133 bronchiectasis, tuberculosis, delay in the diagnosis, delay of replacement therapy onset (8,9) but  
134 to the best of our knowledge, there is no previous study on such association.

135 Our objective was to perform a systematic review of the literature and report of new French cases  
136 to describe the characteristics of PID complicated by AAA, as well as risk factors and clinical  
137 presentation of PID-associated AAA.

138

139 **Materials and Methods**

140 We conducted our systematic review using Preferred Reporting Items for Systematic Reviews  
141 and Meta-Analyses (PRISMA) (10). No previously published research protocol was used. We  
142 searched MEDLINE for all articles pertaining to AAA in PID. We selected published articles  
143 indexed in PubMed involving humans and written in French, English or Spanish between 1946  
144 and 2019. The keywords used were the following ones: (“Common Variable  
145 Immunodeficiency”[Mesh] or “Severe Combined Immunodeficiency”[Mesh] or “Immune  
146 Deficiency Disease” [Supplementary Concept] or “Immune Deficiency, Familial Variable”  
147 [Supplementary Concept]) AND “Amyloidosis”[Mesh], “immune deficiency and amyloidosis”.  
148 We chose to study only primary immunodeficiency, thus excluding immune deficiency secondary  
149 to treatments, Human Immunodeficiency Virus (HIV), cancer or other diseases such as diabetes.  
150 An independent investigator (MD) first screened titles and abstract to exclude duplicate and those  
151 off-topic. Articles were then assessed for eligibility. Only full-text articles with detailed original  
152 cases were selected. These cases could be embedded in any type of clinical study. Two hundred  
153 sixty-two references were identified through database searching, and 44 articles were assessed for  
154 eligibility.

155  
156 French cases were identified using the CEREMAIA (Centre de Référence des Maladies Auto  
157 Inflammatoires rares et des Amyloses/National reference center for autoinflammatory diseases  
158 and inflammatory amyloidosis) and the CEREDIH (Centre de Référence des Déficits  
159 Immunitaires Héritaires). Their medical records were retrieved for analysis.

160  
161 “Non-specific agammaglobulinemia and hypogammaglobulinemia” terms designed  
162 predominantly antibody deficiency (PAD) patients who did not fulfill criteria for CVID nor X-  
163 linked and are defined by a decrease in IgG and IgA/IgM below 2 SD. CVID corresponds to a  
164 more stringent definition (ESID definition) including poor antibody response to vaccines (and/or  
165 absent isohemagglutinins) and/or defects in switched memory B cells (11), the definition of  
166 CVID was used only when comprehensive data were available. Hence, some patients with  
167 hypogammaglobulinemia/ non-specific agammaglobulinemia may correspond to CVID patients.  
168 We separated the cases of PID complicated by AAA in 3 different groups: all cases of PID



169 combined; cases affected by the most frequent PID, CVID, and cases affected by the most  
170 frequent type of PID, humoral deficiency (12).

171

172 Finally, we analyzed the 2 new cases from CEREMAIA and 35 articles from the systematic  
173 literature review (8,9,13–46). Descriptive statistics were performed using the excel software.  
174 (Figure 1)

175

176

177 **Results**

178 ***Description of the two new French cases***

179 *Case 1:* A 26-year-old patient from consanguineous parents was addressed in 2016 for weight  
180 loss, fever, diarrhea, ascites, dyspnea, and cough evolving for a few months. Two of his brothers  
181 had died before 30 years old years. His medical history revealed CVID with B-cell lymphopenia  
182 with absent switched memory B cells, which had not been treated with immunoglobulin  
183 replacement therapy (IRT). His serum C-reactive protein (CRP) was elevated during infections,  
184 and normalized in-between. The clinical examination revealed lymphadenopathy, clubbing  
185 (Figure 2 A), edema, and signs of respiratory failure. CT scan showed cystic bronchiectasis  
186 (Figure 2 B) and an irregular, excavated opacity with an air-fluid level associated with pleuritis  
187 and ascites. During a hospitalization for pulmonary infection, a nephrotic syndrome was detected  
188 with normal renal function. Laboratory exams showed normal total cells count except for  
189 lymphocytosis at 7.22 g/l, CRP at 23 mg/l, and total immunoglobulin (Ig) level of 0.6 g/L (IgG  
190 3.09 g/L, IgA <0.063 g/l and IgM 0.26 g/l). Lymphocyte immunophenotyping revealed B-cell  
191 lymphopenia without autoantibodies. A salivary gland biopsy showed abundant amyloid deposits  
192 staining with AA-antibody (Figure 2 C) confirming the diagnosis of AAA. No *MEFV* mutation  
193 was detected by Sanger sequencing. The patient was discharged with long-term roxithromycin  
194 treatment to prevent pulmonary infection, and IRT was started. Unfortunately, he died at 29 years  
195 from sepsis and acute respiratory failure.

196 *Case 2:* A woman was diagnosed with CVID at the age of 20 years because of recurrent severe  
197 respiratory tract infections, bronchiectasis, and severe hypogammaglobulinemia. Immunological  
198 studies demonstrated absence of IgG and IgA with very low IgM level (0.13 g/l), lymphopenia  
199 with marked CD4 lymphopenia (144/mm<sup>3</sup>), Between the age of 35 years and 58 years, she  
200 received intermittent intramuscular IRT, then continuous intravenous IRT, and finally  
201 subcutaneous IRT. Antibiotic prophylaxis with azithromycin and trimethoprim/sulfamethoxazole  
202 was added because of persistent infections including pneumonia and other respiratory tract  
203 infections, but also HPV, VZV, and HSV infections and Norovirus infection. At the age of 68  
204 years, she was admitted for diarrhea, weight loss, and severe portal hypertension, renal  
205 dysfunction, and microalbuminuria. A kidney biopsy and revealed tubular and vascular AA  
206 protein deposits. Genetic studies identified two compound heterozygous *LRBA* variants (*Exon 57*  
207 *c.8426 C>T* and *c.8464 G>A*). The first variant (*c.8426 C>T*; p.S2809L) is a very frequent SNP

208 found at 2.10-1 in GNOMAD public database. She died at 73 years old from complications of  
209 portal hypertension.

## 210 *Literature review*

### 211 *Main features of PID associated with AAA*

212 While various immune deficiencies and mechanisms of PID were found, antibody deficiencies  
213 were the most frequent type of PID associated with AAA. CVID (n=14) and non-specific  
214 agamma- and hypogammaglobulinemia (n=12) were predominant, followed by X-linked  
215 agammaglobulinemia (XLA) (n=4), chronic granulomatous disease (CGD) (n=3), Hyper IgM  
216 syndrome (n=3), hereditary complete C4 deficiency (n=1), Leucocyte Adhesion Deficiency  
217 Type-1 (n =1), Hyper IgE syndrome (n=1), and Chediak-Higashi syndrome (n=1) (Figure 3).  
218 While 8 patients had “infections only” CVID, 6 had CVID associated with other complications  
219 such as autoimmune disease, granulomatous disease and lymphocytic infiltration.

220 We studied AAA-associated PID in three different groups: first in all PID combined and then in  
221 two specific groups according to the IUIS classification (1):CVID group and predominantly  
222 antibody deficiency (PAD) group. In our review, CVID, non-specific agamma- and  
223 hypogammaglobulinemia, and X-linked agammaglobulinemia were classified as PAD according  
224 to the IUIS. Global sex ratio was 1.7. The mean age at first signs of immune deficiency was  
225 17.25 years in all PID combined group, 18.5 years in the CVID group, and 20.75 years in the  
226 PAD group. A diagnostic delay of 10.39 +/- 7.04 years was noted in all PID combined group,  
227 with 4 additional years in the CVID group. (Table 1)

228 Consanguinity was identified in 9 patients. Family history was compatible with PID in 12  
229 patients. Genetic analysis was performed in 6 cases (9,35,38,40 and case 2). One homozygous  
230 mutation was identified in exon 12 of *ITGB2* (C590R/C590R) (36), leading to a diagnosis of  
231 Leukocyte Adhesion Deficiency Type-1 (47). Two heterozygous non-synonymous amino acid  
232 substitutions in the *NLRP12* gene, encoding the NALP12 protein (p.H304Y and p.A629D) (41);  
233 and our case 2 with two heterozygous compound missense variants of *LRBA*.

234 PID was associated with lymphoproliferation (lymphadenopathy or splenomegaly n=8), hepatic  
235 complications (regenerative nodular hyperplasia and lymphoid infiltration n=1, hepatomegaly  
236 n=1), dermatological manifestations (oral and perianal ulcerations n=1, psoriasis n=1), clubbing  
237 (n=6), growth retardation (n=5), non-infectious arthritis (n=3), adrenal insufficiency (n=2),

238 anemia (n=2), cardiac features (n=1), partial albinism/ataxia/mental deficiency (n=1), and seizure  
239 (n=1). “Infections only” CVID represented 57% of the CVID group. The sites and types of  
240 infection were extremely varied and were mainly bacterial with pulmonary involvement, but  
241 viral, parasitic, and fungal infections were also reported. Twenty-one patients had bronchiectasis  
242 (52,5%). Eight patients had tuberculosis.

243 Treatments were mostly intended to prevent infections: IRT (n=26) given intravenously or  
244 intramuscularly, prophylactic antimicrobial therapy (trimethoprim n=2, co-trimoxazole n=1,  
245 levofloxacin/roxithromycine n=1, azithromycin n=1, penicillamine and itraconazole n=1, and  
246 undefined n=1), intravenous white blood cell transfusion (n=1), and interferon- $\gamma$  (n=1). A  
247 minority of treatment were directed against the PID itself: abatacept (n=1), omalizumab (n=1),  
248 and corticosteroids (n=1). Delay between first symptoms of PID and onset of treatment was 12.41  
249 +/-7.58 years. Three patients were not compliant with treatment. (Table E1)

250

#### 251 *Main features of PID-associated AAA (Figure 4)*

252 The mean age at diagnosis of AAA was 33.41 +/- 18.15 years in all PID combined group, ranging  
253 from 5 to 75 years. On average, CVID and PAD were diagnosed 4 years later. The average delay  
254 between first symptoms of PID and AAA diagnosis was 16.17 +/- 7.08 years in all PID combined  
255 group, like the CVID and PAD groups. Concomitant diagnosis of PID and AAA was made in 13  
256 patients.

257 The most common presentation of AAA was renal involvement, with nephrotic syndrome in the  
258 majority (n=21), non-nephrotic proteinuria (n=10), and renal failure (n=6). Other clinical  
259 manifestations involved gastro-intestinal signs and symptoms (mostly diarrhea, hepatic  
260 involvement, and splenomegaly), deterioration of general state, adrenal insufficiency, heart  
261 failure, and goiter. Histological diagnosis was made in 10 autopsies, mostly in the oldest cases  
262 reported. Renal tissue was most commonly analyzed (n=20), followed by the intestinal tract  
263 (n=13), node and spleen (n=10), liver (n=4), lungs (n=3), abdominal fat (n=1), thyroid (n=3),  
264 adrenal glands (n=3), skin (n=1), heart (n=1), and salivary glands (n = 1).

265 In patients who received therapy directed against AAA, 6 were given colchicine, 2 steroids, 1  
266 tocilizumab, 1 infliximab, 1 omalizumab, and 1 abatacept. Two patients underwent hemodialysis,  
267 and none were transplanted. (Table E2).

268 **Discussion**

269 AAA is a rare complication of PID (48). Available data is scarce, relying on case reports and,  
270 until more recently, on autopsies. Morbidity reviews and studies on PID usually do not mention  
271 AAA (49,50). With only 40 cases reported in more than 60 years and none reported in large case  
272 series, this work confirms the existence and rarity of AAA occurrence in the PID course (51–55).  
273 The typical clinical picture is a patient displaying recurrent infections, mostly bacterial  
274 pneumonia, since adolescence or beginning of adulthood, with a delayed diagnosis of PID and  
275 onset of treatment after 10 years of evolution. The course is marked by chronic pulmonary  
276 infections as illustrated by bronchiectasis for more than half of them, eventually leading to  
277 development of a nephrotic syndrome revealing AAA 16 years later (figure 4). AAA is  
278 associated with a poor prognosis in PID: 22 out of 40 patients, died during follow-up, at an  
279 average age of 36.1 years old.

280  
281 Antibody disorders, especially CVID and non-specified agamma- and hypogammaglobulinemia,  
282 are the predominant PID complicated by AAA, which parallels their prevalence in national  
283 registers (53,56–59). Thus, for AAA to complicate PID, the latter must be severe enough to  
284 develop chronic infections, but mild enough to allow a prolonged course of chronic  
285 inflammation. A possible additional explanation for overrepresentation of PAD in our study is the  
286 fact that patients with primary antibody defects are rarely undergoing hematopoietic stem cell  
287 transplantation (HSCT), as Ig supplementation is usually sufficient to avoid life-threatening  
288 complications. As AAA is linked to a persistent activation of innate immune cells resulting in a  
289 chronic biological inflammatory syndrome, HSCT might be involved in direct protection against  
290 amyloidosis. AAA risk might therefore be underevaluated in patients with early-onset PIDs  
291 hampering innate or non B cell adaptative immunity and leading to an early consideration for  
292 HSCT. However it is a complication that can develop rapidly in certain contexts, as illustrated by  
293 the case of a pediatric diagnostic LAD-1 that developed amyloidosis at the age of 11 years old.  
294 Moreover, 32,5% of PID diagnosis were made when AAA was diagnosed. From the clinical point  
295 of view, it suggests the value of looking for PID in the etiologic search for AAA.

296 The late PID diagnoses (51–55,60) and length of diagnostic delays (11,60–62) are partly due to  
297 older publications (12,61–63) as well as the numerous cases of CVID which are known to have

298 longer diagnostic delays. but it also may have promoted chronic inflammation for AAA. In CVID  
299 complicated by AAA, the first symptoms appeared rather early in life compared to non-AAA  
300 CVID (5,51,62,64–66), thus suggesting an underlying genetic defect. Moreover, despite having a  
301 generally better prognosis (49,67) “infections only” CVID was more often complicated by AAA  
302 than CVID presenting other non-infectious complications (49,67) highlighting a potential  
303 contributing role of infections in AAA secondary to CVID.

304  
305 Bronchiectasis was overrepresented (52,5%) (52,56,62,68). The specific association between  
306 antibody deficiencies and bronchiectasis as already been reported (69). Bronchiectasis was  
307 present in 90% of our CVID group, compared to 23% in Gathman’s series (51), 37% in  
308 Oksenhendler’s series, and 51% in Ramirez-Vargas’s series (68). Thus, bronchiectasis in PID  
309 could be used as a warning sign of chronic inflammation and increased risk of developing AAA.  
310 In the absence of prior PID diagnosis, diagnosis of bronchiectasis among patients with AAA  
311 diagnosis must lead to PID screening, particularly antibody deficiencies.

312  
313 PID patients with AAA share some similarities with other patients developing AAA: the renal  
314 involvement is the main symptomatic sign of AAA (48), the duration between the underlying  
315 disease onset and AAA diagnosis is similar to the mean delay all underlying disorders combined  
316 for AAA occurrence (6) and to the one reported in other immune dysregulation disease such as  
317 inflammatory bowel disease (70) and rheumatoid arthritis (71). However, some chronic infections  
318 such as tuberculosis (72) have a much shorter delay whereas it is slightly longer for the hereditary  
319 periodic fever syndromes (73). AAA seems to develop earlier in life in PID (6,74) than in other  
320 inflammatory diseases, which may be partly explained by the early onset of PID. Even if data for  
321 a prevalence evaluation of AAA in PID were not available because AAA is a very rare  
322 complication of PID as explained above, this prevalence correlates with the underlying disease .  
323 AAA is more common in monogenic periodic fever syndromes such as Familial Mediterranean  
324 Fever (75).

325  
326 This study raises the question of therapeutic issues in the management of PID complicated by  
327 AAA. Once the underlying PID is diagnosed and, if possible, treated, the first step should be to

328 optimize control of chronic inflammation caused by recurrent infections, with the possibility of  
329 inducing AAA regression (76). Although new protocol for Primary Immune Regulatory  
330 Disorders have been developed, to examine HSCT and other treatment for this group of disorders  
331 (77), in such population, we believe that immunomodulatory agents such as anti-IL1 or anti-IL6  
332 immunotherapies should be preferred as in other patients with AA amyloidosis without  
333 underlying immune defects. These agents increase the risk of severe bacterial events but this risk  
334 seems lesser than the treatment related toxicities of HSCT. Moreover, this risk can be  
335 circumscribed by Ig supplementation and antibioprohylaxis. In other PID situations,  
336 amyloidosis discovery could prompt to discuss early HSCT.

337 The occurrence of AAA in PID is low, suggesting that infections may not be the only  
338 predisposing factor for this complication. IBD and enteropathies are common in CVID, CGD,  
339 Leucocyte Adhesion Deficiency and could increase the risk of AAA in PID patients and one case  
340 of enteropathy (case 2) was reported but unfortunately, the available data was insufficient to  
341 provide a prevalence of IBD for patients with AAA secondary to PID. Many cases were reported  
342 before advances in genetics; given the high percentage of PID patients with consanguinity and  
343 culprit mutations identified since 2011, all PID patients developing AAA could get deep genetic  
344 analysis in order to identify underlying genetic defects and corresponding pathophysiological  
345 mechanisms that could have promotes chronic inflammation and trigger AAA. Going forward,  
346 the next step will be using next generation sequencing to determine if certain genotypes are  
347 particularly associated with AAA occurrence in PID patients.

348  
349 Finally, as a national reference center for AAA in an industrialized country, following the  
350 occurrence of 2 such cases within 1 year,, it appeared important for us to point this complication  
351 out to other specialists given the simplicity of non-invasive screening (proteinuria and renal  
352 function monitoring) versus the life threatening consequence of AAA. Because of the high  
353 prevalence of bronchiectasis in AAA associated to PID we propose to systematically look for  
354 hypogammaglobulinemia in AAA with no other cause than bronchiectasis and the realization of a  
355 thoracic imaging in the AAA check-up to detect bronchiectasis. In order to prevent and detect  
356 earlier the occurrence of AAA: we recommend for patients with PID the realization of an annual  
357 proteinuria/creatininuria ratio and a biannual plasma creatinine and CRP assays in the absence of  
358 acute infectious complications. This assessment seems to us to be sufficient, the digestive signs

359 of AAA being rather non-specific in particular diarrhea and the cardiac involvement being very  
360 rare in AAA unlike other amyloidosis such as AL amyloidosis and ATTR amyloidosis. In the  
361 future, a prospective study for proteinuria/creatininuria on a large cohort of PID and, depending  
362 on the results, a biopsy of the accessory salivary glands, which would be minimally invasive,  
363 could be interesting in order to evaluate this association and its incidence.

364



365 **Conclusion**

366 AAA is a long-term life-threatening complication of PID portending a poor prognosis. Despite  
367 improved understanding of PID, earlier diagnosis, and the availability of therapies for infection  
368 prevention and control, PID remain severe and potentially lethal diseases. This work emphasizes  
369 the importance of prompt identification of PID symptoms by general practitioners, pediatricians,  
370 and other specialists. Once a correct diagnosis is established, institution of measures to prevent  
371 infection (vaccination, IRT, and prophylactic antimicrobial therapy if needed) can play a key role  
372 in avoiding such complications.

373

374 **References**

- 375 1. Bousfiha A, Jeddane L, Picard C, Ailal F, Bobby Gaspar H, Al-Herz W, et al. The 2017 IUIS Phenotypic  
376 Classification for Primary Immunodeficiencies. *J Clin Immunol*. 2018;38(1):129-43.
- 377 2. Picard C, Bobby Gaspar H, Al-Herz W, Bousfiha A, Casanova J-L, Chatila T, et al. International Union  
378 of Immunological Societies: 2017 Primary Immunodeficiency Diseases Committee Report on Inborn  
379 Errors of Immunity. *J Clin Immunol*. 2018;38(1):96-128.
- 380 3. Mouthon L, Berezne A, Guillevin L, Lassoued K. [Diagnosis of primary immunodeficiency in adult  
381 patients]. *Presse Medicale Paris Fr* 1983. mai 2006;35(5 Pt 2):903-11.
- 382 4. Guffroy A, Gies V, Martin M, Korganow A-S. [Primary immunodeficiency and autoimmunity]. *Rev*  
383 *Med Interne*. juin 2017;38(6):383-92.
- 384 5. Cunningham-Rundles C, Bodian C. Common variable immunodeficiency: clinical and immunological  
385 features of 248 patients. *Clin Immunol Orlando Fla*. juill 1999;92(1):34-48.
- 386 6. Lachmann HJ, Goodman HJB, Gilbertson JA, Gallimore JR, Sabin CA, Gillmore JD, et al. Natural  
387 history and outcome in systemic AA amyloidosis. *N Engl J Med*. 7 juin 2007;356(23):2361-71.
- 388 7. Stojanovic KS, Georgin-Lavialle S, Grateau G. [AA amyloidosis]. *Nephrol Ther*. juin  
389 2017;13(4):258-64.
- 390 8. Soysal D, TÜRKKAN E, KARAKUŞ V, TATAR E, KABAYEĞİT ÖY, AVCI A. A case of common variable  
391 immunodeficiency disease and thyroid amyloidosis. 2009;39(3):467-73.
- 392 9. Esenboga S, Çağdas Ayvaz D, Sağlam Ayhan A, Peynircioglu B, Sanal O, Tezcan I. CVID Associated  
393 with Systemic Amyloidosis. *Case Rep Immunol*. 2015;2015:879179.
- 394 10. Moher D, Liberati A, Tetzlaff J, Altman DG, PRISMA Group. Preferred reporting items for systematic  
395 reviews and meta-analyses: the PRISMA statement. *PLoS Med*. 21 juill 2009;6(7):e1000097.
- 396 11. Seidel MG, Kindle G, Gathmann B, Quinti I, Buckland M, van Montfrans J, et al. The European  
397 Society for Immunodeficiencies (ESID) Registry Working Definitions for the Clinical Diagnosis of  
398 Inborn Errors of Immunity. *J Allergy Clin Immunol Pract*. août 2019;7(6):1763-70.
- 399 12. Gathmann B, Goldacker S, Klima M, Belohradsky BH, Notheis G, Ehl S, et al. The German national  
400 registry for primary immunodeficiencies (PID). *Clin Exp Immunol*. août 2013;173(2):372-80.
- 401 13. Teilum G. AMYLOIDOSIS SECONDARY TO AGAMMAGLOBULINAEMIA. *J Pathol Bacteriol*. juill  
402 1964;88:317-20.
- 403 14. Forssman O, Herner B. ACQUIRED AGAMMAGLOBULINAEMIA AND MALABSORPTION. *Acta Med*  
404 *Scand*. déc 1964;176:779-86.
- 405 15. Strachan RW, Gillanders LA. A CASE OF PRIMARY ACQUIRED HYPOGAMMAGLOBULINAEMIA WITH  
406 PERNICIOUS ANAEMIA AND AMYLOIDOSIS: DEMONSTRATED AT THE POSTGRADUATE MEDICAL  
407 SCHOOL OF LONDON. *Br Med J*. 2 janv 1965;1(5426):35-8.

- 408 16. Conn HO, Quintiliani R. Severe diarrhea controlled by gamma globulin in a patient with  
409 agammaglobulinemia, amyloidosis, and thymoma. *Ann Intern Med.* sept 1966;65(3):528-41.
- 410 17. Murray WD, Cook IA. Primary « acquired » hypogammaglobulinaemia and amyloidosis. *J Clin*  
411 *Pathol.* janv 1968;21(1):32-4.
- 412 18. Mawas C, Sors C, Bernier JJ. Amyloidosis associated with primary agammaglobulinemia, severe  
413 diarrhea and familial hypogammaglobulinemia. *Am J Med.* avr 1969;46(4):624-34.
- 414 19. De Seigneux R, Kanfer A, Terrioux P, Sraer JD, Whitworth JA. Letter: Renal amyloidosis in chronic  
415 granulomatous disease. *Br Med J.* 26 oct 1974;4(5938):230.
- 416 20. Ziegler JB, Penny R. Fatal echo 30 virus infection and amyloidosis in X-linked  
417 hypogammaglobulinemia. *Clin Immunol Immunopathol.* janv 1975;3(3):347-52.
- 418 21. Pick AI, Versano I, Schreibman S, Ben-Bassat M, Shoenfeld Y. Agammaglobulinemia, plasma cell  
419 dyscrasia, and amyloidosis in a 12-year-old child. *Am J Dis Child* 1960. juin 1977;131(6):682-6.
- 420 22. Gaffney EF, Lee JC. Systemic amyloidosis and hypogammaglobulinemia. *Arch Pathol Lab Med.* nov  
421 1978;102(11):558-9.
- 422 23. Podjarny E, Mekori YA, Bernhaim J, Klajman A. Sex-linked hypogammaglobulinemia and  
423 amyloidosis. *Arch Pathol Lab Med.* sept 1982;106(9):447-8.
- 424 24. Neild GH, Scott GW, Rowe IF, Pepys MB. Amyloid (type AA) in a patient with  
425 hypogammaglobulinemia. *N Engl J Med.* 14 févr 1985;312(7):446.
- 426 25. Maeda K, Sueishi K, Lida M. A case report of Chediak-Higashi syndrome complicated with systemic  
427 amyloidosis and olivo-cerebellar degeneration. *Pathol Res Pract.* août 1989;185(2):231-7.
- 428 26. Meysman M, Debeuckelaer S, Reynaert H, Schoors DF, Dehou MF, Van Camp B. Systemic  
429 amyloidosis-induced diarrhea in sex-linked agammaglobulinemia. *Am J Gastroenterol.* août  
430 1993;88(8):1275-7.
- 431 27. Kotilainen P, Vuori K, Kainulainen L, Aho H, Saario R, Asola M, et al. Systemic amyloidosis in a  
432 patient with hypogammaglobulinaemia. *J Intern Med.* août 1996;240(2):103-6.
- 433 28. Tezcan I, Ersoy F, Sanal O, Gönc EN, Arici M, Berkel I. A case of X linked agammaglobulinaemia  
434 complicated with systemic amyloidosis. *Arch Dis Child.* juill 1998;79(1):94.
- 435 29. Oner A, Demircin G, Erdoğan O, Bülbül M, Memiş L, Arda N, et al. A family with  
436 hyperimmunoglobulin M syndrome and systemic amyloidosis. *Nephrol Dial Transplant Off Publ Eur*  
437 *Dial Transpl Assoc - Eur Ren Assoc.* sept 2000;15(9):1480-2.
- 438 30. Celik AF, Altiparmak MR, Pamuk GE, Pamuk ON, Tabak F. Association of secondary amyloidosis with  
439 common variable immune deficiency and tuberculosis. *Yonsei Med J.* 31 déc 2005;46(6):847-50.
- 440 31. Kaltenis P, Mudeniené V, Maknavicius S, Seinins D. Renal amyloidosis in a child with chronic  
441 granulomatous disease and invasive aspergillosis. *Pediatr Nephrol Berl Ger.* mai 2008;23(5):831-4.

- 442 32. Aghamohammadi A, Shafiei A, Abolhassani H, Sherkat R, Mahjoub F, Rezaei N. Renal amyloidosis in  
443 common variable immunodeficiency. *Nefrol Publicacion Of Soc Espanola Nefrol.* 2010;30(4):474-6.
- 444 33. Turkmen K, Anil M, Solak Y, Atalay H, Esen H, Tonbul HZ. A hepatitis C-positive patient with new  
445 onset of nephrotic syndrome and systemic amyloidosis secondary to common variable  
446 immunodeficiency. *Ann Saudi Med.* oct 2010;30(5):401-3.
- 447 34. Aydin Z, Gursu M, Ozturk S, Kilicaslan I, Kazancioglu R. A case of primary immune deficiency  
448 presenting with nephrotic syndrome. *NDT Plus.* oct 2010;3(5):456-8.
- 449 35. Helal I, Goucha R, Hamida FB, Elyounsi F, Maiz HB, Kheder A. Renal AA amyloidosis in a patient with  
450 hereditary complete complement C4 deficiency. *Saudi J Kidney Dis Transplant Off Publ Saudi Cent  
451 Organ Transplant Saudi Arab.* sept 2011;22(5):1008-11.
- 452 36. Rawat A, Singh S, Sharma D, Suri D, Rajwanshi A, Etzioni A. Amyloidosis in a child with leucocyte  
453 adhesion deficiency type-1: an unusual association. *Indian J Pediatr.* déc 2011;78(12):1546-8.
- 454 37. Firinu D, Serusi L, Lorrain MM, Grande M, Murgia G, Barca MP, et al. Systemic reactive (AA)  
455 amyloidosis in the course of common variable immunodeficiency. *Amyloid Int J Exp Clin Investig Off  
456 J Int Soc Amyloidosis.* juin 2011;18 Suppl 1:214-6.
- 457 38. Kadiroğlu AK, Yıldırım Y, Yılmaz Z, Kayabaşı H, Avcı Y, Yıldırım MS, et al. A rare cause of secondary  
458 amyloidosis: common variable immunodeficiency disease. *Case Rep Nephrol.* 2012;2012:860208.
- 459 39. Gonzalo-Garijo MA, Sánchez-Vega S, Pérez-Calderón R, Pérez-Rangel I, Corrales-Vargas S,  
460 Fernández de Mera JJ, et al. Renal amyloidosis in a patient with X-linked agammaglobulinemia  
461 (Bruton's disease) and bronchiectasis. *J Clin Immunol.* janv 2014;34(1):119-22.
- 462 40. Meira T, Sousa R, Cordeiro A, Ilgenfritz R, Borralho P. Intestinal Amyloidosis in Common Variable  
463 Immunodeficiency and Rheumatoid Arthritis. *Case Rep Gastrointest Med.* 2015;2015:405695.
- 464 41. Borte S, Celiksoy MH, Menzel V, Ozkaya O, Ozen FZ, Hammarström L, et al. Novel NLRP12  
465 mutations associated with intestinal amyloidosis in a patient diagnosed with common variable  
466 immunodeficiency. *Clin Immunol Orlando Fla.* oct 2014;154(2):105-11.
- 467 42. Arslan S, Ucar R, Yavsan DM, Esen H, Maden E, Reislı I, et al. Common variable immunodeficiency  
468 and pulmonary amyloidosis: a case report. *J Clin Immunol.* mai 2015;35(4):344-7.
- 469 43. Balwani MR, Kute VB, Shah PR, Wakhare P, Trivedi HL. Secondary renal amyloidosis in a patient of  
470 pulmonary tuberculosis and common variable immunodeficiency. *J Nephro pharmacology.*  
471 2015;4(2):69-71.
- 472 44. Darougar S, Rashid Farokhi F, Tajik S, Baghaie N, Amirmoini M, Bashardoust B, et al. Amyloidosis as  
473 a Renal Complication of Chronic Granulomatous Disease. *Iran J Kidney Dis.* juill 2016;10(4):228-32.
- 474 45. Gupta K, Rawat A, Agrawal P, Jindal A, Nada R, Saikia B, et al. Infectious and non-infectious  
475 complications in primary immunodeficiency disorders: an autopsy study from North India. *J Clin  
476 Pathol.* mai 2018;71(5):425-35.

- 477 46. González Sanchidrián S, Davin Carrero E, Gallego Domínguez S. AA amyloidosis associated with  
478 recurrent infections in the hyperimmunoglobulin E syndrome. *Med Clin (Barc)*. 20  
479 2017;149(6):274-5.
- 480 47. van de Vijver E, Maddalena A, Sanal Ö, Holland SM, Uzel G, Madkaikar M, et al. Hematologically  
481 important mutations: leukocyte adhesion deficiency (first update). *Blood Cells Mol Dis*. 15 janv  
482 2012;48(1):53-61.
- 483 48. Papa R, Lachmann HJ. Secondary, AA, Amyloidosis. *Rheum Dis Clin North Am*. 2018;44(4):585-603.
- 484 49. Cunningham-Rundles C. Common variable immune deficiency: Dissection of the variable. *Immunol*  
485 *Rev*. 2019;287(1):145-61.
- 486 50. Kobrynski L, Powell RW, Bowen S. Prevalence and morbidity of primary immunodeficiency diseases,  
487 United States 2001-2007. *J Clin Immunol*. nov 2014;34(8):954-61.
- 488 51. Gathmann B, Mahlaoui N, CEREDIH, Gérard L, Oksenhendler E, Warnatz K, et al. Clinical picture and  
489 treatment of 2212 patients with common variable immunodeficiency. *J Allergy Clin Immunol*. juill  
490 2014;134(1):116-26.
- 491 52. Edgar JDM, Buckland M, Guzman D, Conlon NP, Knerr V, Bangs C, et al. The United Kingdom  
492 Primary Immune Deficiency (UKPID) Registry: report of the first 4 years' activity 2008-2012. *Clin Exp*  
493 *Immunol*. janv 2014;175(1):68-78.
- 494 53. Marschall K, Hoernes M, Bitzenhofer-Grüber M, Jandus P, Duppenenthaler A, Wuillemin WA, et al.  
495 The Swiss National Registry for Primary Immunodeficiencies: report on the first 6 years' activity  
496 from 2008 to 2014. *Clin Exp Immunol*. oct 2015;182(1):45-50.
- 497 54. Joshi AY, Iyer VN, Hagan JB, St Sauver JL, Boyce TG. Incidence and temporal trends of primary  
498 immunodeficiency: a population-based cohort study. *Mayo Clin Proc*. 2009;84(1):16-22.
- 499 55. Veramendi-Espinoza LE, Zafra-Tanaka JH, Pérez-Casquino GA, Córdova-Calderón WO. Diagnostic  
500 Delay of Primary Immunodeficiencies at a Tertiary Care Hospital in Peru- Brief Report. *J Clin*  
501 *Immunol*. mai 2017;37(4):383-7.
- 502 56. Kirkpatrick P, Riminton S. Primary immunodeficiency diseases in Australia and New Zealand. *J Clin*  
503 *Immunol*. sept 2007;27(5):517-24.
- 504 57. Matamoros Florí N, Mila Llambi J, Español Boren T, Raga Borja S, Fontan Casariego G. Primary  
505 immunodeficiency syndrome in Spain: first report of the National Registry in Children and Adults. *J*  
506 *Clin Immunol*. juill 1997;17(4):333-9.
- 507 58. Abuzakouk M, Feighery C. Primary immunodeficiency disorders in the Republic of Ireland: first  
508 report of the national registry in children and adults. *J Clin Immunol*. janv 2005;25(1):73-7.
- 509 59. Stray-Pedersen A, Abrahamsen TG, Frøland SS. Primary immunodeficiency diseases in Norway. *J*  
510 *Clin Immunol*. nov 2000;20(6):477-85.

- 511 60. Al-Saud B, Al-Mousa H, Al Gazlan S, Al-Ghoniaum A, Arnaout R, Al-Seraihy A, et al. Primary  
512 Immunodeficiency Diseases in Saudi Arabia: a Tertiary Care Hospital Experience over a Period of  
513 Three Years (2010-2013). *J Clin Immunol.* oct 2015;35(7):651-60.
- 514 61. Gathmann B, Binder N, Ehl S, Kindle G, ESID Registry Working Party. The European internet-based  
515 patient and research database for primary immunodeficiencies: update 2011. *Clin Exp Immunol.*  
516 mars 2012;167(3):479-91.
- 517 62. Oksenhendler E, Gérard L, Fieschi C, Malphettes M, Mouillot G, Jaussaud R, et al. Infections in 252  
518 patients with common variable immunodeficiency. *Clin Infect Dis Off Publ Infect Dis Soc Am.* 15 mai  
519 2008;46(10):1547-54.
- 520 63. Seymour B, Miles J, Haeney M. Primary antibody deficiency and diagnostic delay. *J Clin Pathol.* mai  
521 2005;58(5):546-7.
- 522 64. Abbott JK, Gelfand EW. Common Variable Immunodeficiency: Diagnosis, Management, and  
523 Treatment. *Immunol Allergy Clin North Am.* nov 2015;35(4):637-58.
- 524 65. Lin L-J, Wang Y-C, Liu X-M. Clinical and immunological features of common variable  
525 immunodeficiency in China. *Chin Med J (Engl).* 5 févr 2015;128(3):310-5.
- 526 66. Quinti I, Soresina A, Spadaro G, Martino S, Donnanno S, Agostini C, et al. Long-term follow-up and  
527 outcome of a large cohort of patients with common variable immunodeficiency. *J Clin Immunol.*  
528 mai 2007;27(3):308-16.
- 529 67. Chapel H, Lucas M, Lee M, Bjorkander J, Webster D, Grimbacher B, et al. Common variable  
530 immunodeficiency disorders: division into distinct clinical phenotypes. *Blood.* 15 juill  
531 2008;112(2):277-86.
- 532 68. Ramírez-Vargas N, Arablin-Oropeza SE, Mojica-Martínez D, Yamazaki-Nakashimada MA, de la Luz  
533 García-Cruz M, Terán-Juárez LM, et al. Clinical and immunological features of common variable  
534 immunodeficiency in Mexican patients. *Allergol Immunopathol (Madr).* juin 2014;42(3):235-40.
- 535 69. Soler-Palacín P, de Gracia J, González-Granado LI, Martín C, Rodríguez-Gallego C, Sánchez-Ramón S,  
536 et al. Primary immunodeficiency diseases in lung disease: warning signs, diagnosis and  
537 management. *Respir Res.* 12 nov 2018;19(1):219.
- 538 70. Greenstein AJ, Sachar DB, Panday AK, Dikman SH, Meyers S, Heimann T, et al. Amyloidosis and  
539 inflammatory bowel disease. A 50-year experience with 25 patients. *Medicine (Baltimore).* sept  
540 1992;71(5):261-70.
- 541 71. Fiter Aresté J, Nolla Solé JM, Gómez Vaquero C, Valverde García J, Roig Escofet D. [Secondary  
542 amyloidosis in rheumatoid arthritis. A clinical study of 29 patients]. *An Med Interna Madr Spain*  
543 1984. déc 1999;16(12):615-9.
- 544 72. Dixit R, Gupta R, Dave L, Prasad N, Sharma S. Clinical profile of patients having pulmonary  
545 tuberculosis and renal amyloidosis. *Lung India Off Organ Indian Chest Soc.* avr 2009;26(2):41-5.

- 546 73. Lane T, Loeffler JM, Rowczenio DM, Gilbertson JA, Bybee A, Russell TL, et al. AA amyloidosis  
547 complicating the hereditary periodic fever syndromes. *Arthritis Rheum.* avr 2013;65(4):1116-21.
- 548 74. Real de Asúa D, Costa R, Galván JM, Filigheddu MT, Trujillo D, Cadiñanos J. Systemic AA  
549 amyloidosis: epidemiology, diagnosis, and management. *Clin Epidemiol.* 2014;6:369-77.
- 550 75. Grateau G, Hentgen V, Stojanovic KS, Jéru I, Amselem S, Steichen O. How should we approach  
551 classification of autoinflammatory diseases? *Nat Rev Rheumatol.* oct 2013;9(10):624-9.
- 552 76. Castellano I, Gómez-Martino JR, Hernández MT, Novillo R, Covarsí A. [Remission of nephrotic  
553 syndrome caused by renal amyloidosis secondary to pulmonary tuberculosis after tuberculostatic  
554 treatment]. *Nefrol Publicacion Of Soc Espanola Nefrol.* févr 2001;21(1):88-91.
- 555 77. Torgerson TR. Primary Immune Regulatory Disorders: Clinical Presentations, Treatments and  
556 Outcomes available from <https://grantome.com/grant/NIH/U54-AI082973-11-6475>. NIH; 2019.
- 557
- 558
- 559
- 560
- 561

562 **Table's legends**

563 **Table 1 Epidemiological data of PID complicated by AAA**

564 **Abbreviations;** AAA AA Amyloidosis; PID Primitive Immunodeficiency; Yrs Years; CVID  
565 Common variable immunodeficiency

566



567 **Figure's legends**

568 **Figure 1. Flow-chart diagram depicting the selection process**

569

570

571


572 **Figure 2. Case 1**

573 **A. Clubbing B. CT Chest: Diffuse bronchiectasis C. Microscopic feature of salivary biopsy**  
574 **showing amyloid deposition around the salivary ducts (C.1 Congo red stain, ×100 and**  
575 **C.2Immunohistochemical reactivity of deposited material with anti-amyloid A protein x400**

576 **Figure 3. Distribution of included patients into the 9 major diagnostic categories of PID (1)**


577

578 **Figure 4 Synthesis of the natural history of AAA in PID**







A.




B.



C.1




C.2





- Immunodeficiencies affecting cellular and humoral immunity
- Combined immunodeficiencies with associated or syndromic features
- Predominantly antibody deficiency
- Diseases of immune dysregulation
- Congenital defects of phagocyte number, function, or both
- Defects in Intrinsic and Innate Immunity
- Complement deficiencies
- Autoinflammatory disorders
- Phenocopies of PID





**Consanguinity**  
22,5%





	Mean age (yrs)			Minimum age (yrs)			Maximum age (yrs)			Standard deviation (yrs)		
	All PID	CVID	PAD	All PID	CVID	PAD	All PID	CVID	PAD	All PID	CVID	PAD
<b>First signs of PID</b>	17.25	18.50	20.75	0	7	0	64	64	64	17.98	18.86	17.91
<b>Diagnosis of PID</b>	22.20	28.86	25.64	0.17	9.00	0.17	64	64	64	16.02	14.75	15.99
<b>Initiation of treatment of PID</b>	27.06	33.00	31.19	0.17	24.00	0.17	74	48	74	20.33	7.71	17.67
<b>First signs of AAA</b>	32.41	38.14	38.26	7	20	12	72	72	72	17.32	16.83	16.23
<b>Diagnosis of AAA</b>	33.41	39.03	39.13	7	20.5	12	75	75	68	18.15	17.34	16.77
<b>Death</b>	36.13	41.00	41.88	8.5	21	15	75	73	75	20.27	20.49	18.41
<b>Delay between first signs of PID and AAA diagnosis</b>	16.17	16.14	16.82	2	2	2	29	22	29	7.08	6.94	7.41
<b>Delay between first signs of PID and PID diagnosis</b>	10.39	14.00	11.12	0	0	0	22	22	22	7.04	7.95	7.35
<b>Delay between first signs of PID and initiation of treatment</b>	12.41	20.67	13.90	0	0	0	22	22	22	7.58	1.54	7.43