



The missing link between familial Mediterranean fever and recurrent aseptic meningitis

J. Capron, G. Grateau, O. Steichen

► To cite this version:

J. Capron, G. Grateau, O. Steichen. The missing link between familial Mediterranean fever and recurrent aseptic meningitis: Letter to the Editor. 2019, pp.350. 10.1016/j.pedneo.2019.03.007 . inserm-04051002

HAL Id: inserm-04051002


<https://inserm.hal.science/inserm-04051002>

Submitted on 29 Mar 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

Copyright

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Letter to the Editor

The missing link between familial Mediterranean fever and recurrent aseptic meningitis



To the Editor,

Yusef et al. reported a case of familial Mediterranean fever (FMF) presenting with recurrent aseptic meningitis (RAM).¹ There is no consensus on a possible association between the two conditions, and we have proposed criteria to assess the strength of evidence of case reports.² In the present case, FMF was diagnosed based on a homozygous E148Q mutation on the MEFV gene. However, the pathogenicity of the E148Q mutation is uncertain, and the diagnosis of FMF requires clinical features that are missing in this case.³ Although colchicine seems to have prevented further episodes of aseptic meningitis, the lack of other systemic features or inflammation during the episodes questions the association of RAM with any systemic disease. Nevertheless, in this young Jordanian patient, one may advise further investigations, such as ophthalmologic examination, to rule out Behcet's disease, which may cause RAM.⁴ According to our criteria, the evidence supporting FMF as a cause of RAM is missing in most case reports, including this one.

Conflicts of interest

The authors have no conflicts of interest relevant to this article.

References

1. Yusef D, Khasawneh W. Familial Mediterranean fever presenting with recurrent aseptic meningitis: a case report. *Pediatr Neonatol* 2017;58:378–9.
2. Capron J, Grateau G, Steichen O. Is recurrent aseptic meningitis a manifestation of familial Mediterranean fever? A systematic review. *Clin Exp Rheumatol* 2013;31:127–32.
3. Giancane G, Ter Haar NM, Wulffraat N, Vastert SJ, Barron K, Hentgen V, et al. Evidence-based recommendations for genetic diagnosis of familial Mediterranean fever. *Ann Rheum Dis* 2015;74:635–41.
4. Kidd D, Steuer A, Denman A, Rudge P. Neurological complications in Behcet's syndrome. *Brain* 1999;122:2183–94.

J. Capron*

Assistance Publique - Hôpitaux de Paris, Hôpital Saint Antoine, Service de neurologie, F-75012, Paris, France

G. Grateau
O. Steichen

Assistance Publique - Hôpitaux de Paris, Hôpital Tenon, Service de médecine interne, F-75020, Paris, France

*Corresponding author. Service de neurologie, Hôpital Saint Antoine, 184 rue du Faubourg Saint Antoine, 75012, Paris, France.

E-mail address: jean.capron@aphp.fr (J. Capron)

Sep 6, 2016