



**HAL**  
open science

# Estimation of Donor Renal Function After Living Donor Nephrectomy: The Value of the Toulouse-Rangueil Predictive Model

Thomas Prudhomme, Mathieu Roumiguie, Marc Olivier Timsit

► **To cite this version:**

Thomas Prudhomme, Mathieu Roumiguie, Marc Olivier Timsit. Estimation of Donor Renal Function After Living Donor Nephrectomy: The Value of the Toulouse-Rangueil Predictive Model. *Transplant International*, 2023, 36, pp.11393. 10.3389/ti.2023.11393 . inserm-04200067

**HAL Id: inserm-04200067**

**<https://inserm.hal.science/inserm-04200067v1>**

Submitted on 8 Sep 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



# Estimation of Donor Renal Function After Living Donor Nephrectomy: The Value of the Toulouse-Rangueil Predictive Model

Thomas Prudhomme<sup>1,2\*</sup>, Mathieu Roumiguie<sup>1</sup> and Marc Olivier Timsit<sup>3</sup>

<sup>1</sup>Department of Urology and Kidney Transplantation, Toulouse-Rangueil University Hospital, Toulouse, France, <sup>2</sup>Center for Research in Transplantation and Translational Immunology, Nantes University, INSERM, UMR 1064, Nantes, France,

<sup>3</sup>Department of Urology and Transplant Surgery, AP-HP, Necker Hospital and European Hospital Georges Pompidou, Paris, France

**Keywords:** living donor nephrectomy, renal function, external validation, predictive model, concordance

**A Forum discussing:**

**External Validation of the Toulouse-Rangueil Predictive Model to Estimate Donor Renal Function After Living Donor Nephrectomy**

by Almeida M, Calheiros Cruz G, Sousa C, Figueiredo C, Ventura S, Silvano J, Pedroso S, Martins LS, Ramos M and Malheiro J (2023). *Transpl Int* 36:11151. doi: 10.3389/ti.2023.11151

Assessment of renal function after living donor nephrectomy represents a current hot topic in kidney transplantation. It is well known that living donor kidney transplantation (LDKT) is the best treatment for end-stage renal disease (ESRD) patients eligible for transplantations [1]. However, it has been demonstrated that living donors are at an increased risk of chronic kidney disease (CKD) and ESRD comparing to healthy non-donors [2, 3], exposing them to cardiovascular and global morbidity and mortality of CKD [4]. For these reasons, personalized instruments to evaluate post-operative renal function are necessary in order to offer the benefit of LDKT to recipients while controlling the immediate and long-term negative effects of reduced nephron mass for the donor.

Almeida et al. [5], have just published in *Transplant International*, an external validation of the predictive model to estimate the donor renal function after living donor nephrectomy, developed by our team at the Rangueil University Hospital in Toulouse. They reported a significant correlation (Pearson  $r = 0.67$ ;  $p < 0.001$ ) and concordance (Bland-Altman plot with 95% limits of agreement  $-21.41$  to  $26.47$  mL/min/1.73 m<sup>2</sup>;  $p < 0.001$ ) between predicted and observed 1-year estimated glomerular filtration rate (eGFR). The area under the receiver operating characteristic curve (AUROC) showed a good discriminative ability of the formula to predict CKD [AUC: 0.83 (CI 95%: 0.78–0.88;  $p < 0.001$ )].

It is very rewarding to see a confirmation of our previous report regarding this predictive model to estimate the 1-year post-donation eGFR and risk of CKD. First, Benoit et al. [6] retrospectively evaluated our living donor cohort and demonstrated that age and preoperative eGFR were strong independent predictors of postoperative eGFR. A formula, using multiple linear regression model, was designed: postoperative eGFR (CKD-EPI, mL/min/1.73 m<sup>2</sup>) =  $31.71 + (0.521 \times \text{preoperative eGFR}) - (0.314 \times \text{age})$ . The internal validation reported an optimal statistical performance with a significant correlation ( $r = 0.65$ ) and an AUROC of the model of 0.83 (CI 95%: 0.72–0.93;  $p < 0.001$ ). Then, this model was externally validated using the Necker Hospital living donor cohort [7]. A significant correlation (Pearson  $r = 0.66$ ;  $p < 0.001$ ) and concordance (Bradley-Blackwood  $F = 49.189$ ;  $p < 0.001$ ) were reported between predicted and observed 1-year eGFR. The AUROC in this external population confirmed discriminative ability



OPEN ACCESS

**\*Correspondence:**

Thomas Prudhomme  
prudhomme-thomas@hotmail.fr

**Received:** 23 March 2023

**Accepted:** 11 May 2023

**Published:** 19 May 2023

**Citation:**

Prudhomme T, Roumiguie M and Timsit MO (2023) Estimation of Donor Renal Function After Living Donor Nephrectomy: The Value of the Toulouse-Rangueil Predictive Model. *Transpl Int* 36:11393. doi: 10.3389/ti.2023.11393

of this model to predict CKD (AUC: 0.86 (CI 95%: 0.82–0.89;  $p < 0.01$ )). Kullik et al. [8] also externally validated the model and reported a good correlation to the observed 1-year post-donation eGFR.

External validation is necessary to determine predictive model reproducibility and generalizability to different patients [9]. Correlation and concordance are usually degraded by external validations [10], the outcomes reported by Almeida et al. [5] confirmed the effectiveness of the formula to estimate the 1-year post-donation eGFR and risk of CKD. The strength of their study is the evaluation of model performance according to the sex of the donor; they reported similar performance between females and males.

Along with an extension of marginal deceased donors (extended criteria donor, donation after circulatory death donors) [11, 12], extending the age limit for living donors to expand the pool has been considered [13, 14]. However, donor age was to be a strong predictor of CKD after living donor nephrectomy which may defer the wish to extend the age limit of donors [6, 15]. Preoperative eGFR was also an independent predictor of postoperative eGFR [6]. Thus, a particular attention should be paid to preoperative eGFR and the risk of postoperative CKD with potential an increasing pool of donor over 60 years. However, considering the performance of the predictive model, the use of this tool at the living donor evaluation consultation, combined with a global donor assessment (i.e., comorbidities assessment, donor age and

expected lifespan), could contraindicated some potential donors with predicted lower 1-year eGFR.

The prediction of the postoperative renal function is the key point of donor's candidate evaluation. This predictive model, represents a simple, low cost, non-invasive tool that can be easily joined to the living donor evaluation routine consultation, paving the way for an improved decision-making process.

## DATA AVAILABILITY STATEMENT

The original contributions presented in the study are included in the article/supplementary material, further inquiries can be directed to the corresponding author.

## AUTHOR CONTRIBUTIONS

TP: Manuscript writing, MR: Manuscript editing, MT: Manuscript editing.

## CONFLICT OF INTEREST

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

## REFERENCES

1. Gruessner RWG, Gruessner AC. Solid-organ Transplants from Living Donors: Cumulative United States Experience on 140,156 Living Donor Transplants over 28 Years. *Transpl Proc* (2018) 50:3025–35. doi:10.1016/j.transproceed.2018.07.024
2. Mjoen G, Hallan S, Hartmann A, Foss A, Midtvedt K, Øyen O, et al. Long-term Risks for Kidney Donors. *Kidney Int* (2014) 86:162–7. doi:10.1038/ki.2013.460
3. Muzaale AD, Massie AB, Wang MC, Montgomery RA, McBride MA, Wainright JL, et al. Risk of End-Stage Renal Disease Following Live Kidney Donation. *JAMA* (2014) 311:579–86. doi:10.1001/jama.2013.285141
4. Ross LF, Thistlethwaite JR, Jr. Long-term Consequences of Kidney Donation. *N Engl J Med* (2009) 360:2371–2.
5. Almeida M, Calheiros Cruz G, Sousa C, Figueiredo C, Ventura S, Silvano J, et al. External Validation of the Toulouse-Rangueil Predictive Model to Estimate Donor Renal Function after Living Donor Nephrectomy. *Transpl Int* (2023) 36:11151. doi:10.3389/ti.2023.11151
6. Benoit T, Game X, Roumiguie M, Sallusto F, Doumerc N, Beauval JB, et al. Predictive Model of 1-year Postoperative Renal Function after Living Donor Nephrectomy. *Int Urol Nephrol* (2017) 49:793–801. doi:10.1007/s11255-017-1559-1
7. Benoit T, Prudhomme T, Adypagavane A, Malavaud B, Soulié M, Gamé X, et al. External Validation of a Predictive Model to Estimate Renal Function after Living Donor Nephrectomy. *Transplantation* (2021) 105(11):2445–50. doi:10.1097/TP.0000000000003643
8. Kulik U, Gwiasda J, Oldhafer F, Kaltenborn A, Arelin V, Gueler F, et al. External Validation of a Proposed Prognostic Model for the Prediction of 1-year Postoperative eGFR after Living Donor Nephrectomy. *Int Urol Nephrol* (2017) 49:1937–40. doi:10.1007/s11255-017-1683-y
9. Ramspek CL, Jager KJ, Dekker FW, Zoccali C, van Diepen M. External Validation of Prognostic Models: What, Why, How, When and Where? *Clin Kidney J* (2021) 14:49–58. doi:10.1093/ckj/sfaa188
10. Sperrin M, Riley RD, Collins GS, Martin GP. Targeted Validation: Validating Clinical Prediction Models in Their Intended Population and Setting. *Diagn Progn Res* (2022) 6:24. doi:10.1186/s41512-022-00136-8
11. Neyrinck A, Van Raemdonck D, Monbaliu D. Donation after Circulatory Death: Current Status. *Curr Opin Anaesthesiol* (2013) 26:382–90. doi:10.1097/ACO.0b013e328360dc87
12. Israni AK, Zaun D, Rosendale JD, Snyder JJ, Kasiske BL. OPTN/SRTR 2012 Annual Data Report: Deceased Organ Donation. *Am J Transpl* (2014) 14(1):167–83. doi:10.1111/ajt.12585
13. Serrano OK, Yadav K, Bangdiwala A, Vock DM, Dunn TB, Finger EB, et al. Age Alone Is Not a Contraindication to Kidney Donation: Outcomes of Donor Nephrectomy in the Elderly. *Clin Transpl* (2018) 32:e13287. doi:10.1111/ctr.13287
14. Cantarelli C, Cravedi P. Criteria for Living Donation from Marginal Donors: One, No One, and One Hundred Thousand. *Nephron* (2019) 142:227–32. doi:10.1159/000500498
15. Ibrahim HN, Foley R, Tan L, Rogers T, Bailey RF, Guo H, et al. Long-term Consequences of Kidney Donation. *N Engl J Med* (2009) 360:459–69. doi:10.1056/NEJMoa0804883

Copyright © 2023 Prudhomme, Roumiguie and Timsit. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.