



**HAL**  
open science

# Contacting the gut: Mitochondria-associated Endoplasmic Reticulum Membranes in the Enteric Nervous System

Jenny Greig, Giada Delfino, Pascal Derkinderen, Sébastien Paillusson

► **To cite this version:**

Jenny Greig, Giada Delfino, Pascal Derkinderen, Sébastien Paillusson. Contacting the gut: Mitochondria-associated Endoplasmic Reticulum Membranes in the Enteric Nervous System. Contact, 2024, 7, 10.1177/25152564241244941 . inserm-04587199

**HAL Id: inserm-04587199**

**<https://inserm.hal.science/inserm-04587199v1>**

Submitted on 26 Oct 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

# Contacting the gut: Mitochondria-associated Endoplasmic Reticulum Membranes in the Enteric Nervous System

Jenny Greig<sup>1,\*</sup>, Giada Delfino<sup>2,\*</sup>, Pascal Derkinderen<sup>2</sup>, and Sébastien Paillusson<sup>2</sup> 

Contact  
Volume 7: 1–3  
© The Author(s) 2024  
Article reuse guidelines:  
sagepub.com/journals-permissions  
DOI: 10.1177/25152564241244941  
journals.sagepub.com/home/ctc



## Abstract

Changes in the connections between the endoplasmic reticulum (ER) and mitochondria, as well as alterations in mitochondria-associated ER membrane (MAM) signalling, have been documented in various neurodegenerative diseases affecting the brain. Despite the growing recognition of the significance of the gut-brain axis in neurodegenerative conditions, there has been no prior investigation into the biology of MAM within the enteric nervous system (ENS). Our recent research reveals, for the first time, the existence of connections between the ER and mitochondria within enteric neurons. Additionally, we observed alterations in the dynamics of these connections in the enteric neurons from a mouse model exhibiting age-related neurodegeneration. These findings provide the first detailed characterization of MAM in the ENS under physiological conditions and in a mouse model of age-associated neurodegeneration and shed new light on the potential role of enteric MAM in the context of neurodegenerative disorders.

## Keywords

enteric nervous system, mitochondria-associated ER membranes, Parkinson's disease, Alzheimer's disease, ageing

Over the past decade, changes in both mitochondrial-associated membrane (MAM) dynamics and signalling have consistently been linked to age-related neurodegenerative disorders such as Parkinson's disease (PD) and Alzheimer's disease (AD). Neuronal damage in these disorders often coincides with cellular functions regulated by contacts between the endoplasmic reticulum (ER) and mitochondria, as well as MAM signalling (Markovinovic et al., 2022).

A growing body of research indicates that the gut plays a significant role in neurodegeneration, as seen in PD, AD, and age-related neurodegeneration. The enteric nervous system (ENS), a key component of the gut-brain axis, governs major gastrointestinal functions (Markovinovic et al., 2022). Despite extensive research on MAM in the central nervous system (CNS), no data are available regarding MAM in the ENS, and this forms the focal point of the current study.

In our recently published study (Delfino et al., 2024), we have, for the first time, identified the presence of ER-mitochondria contacts in enteric neurons. Utilizing electron microscopy, we conducted a detailed characterization of these interactions. Our initial findings demonstrated that mitochondria naturally form MAM with ER membranes in enteric neurons of the myenteric plexus in young adult C57BL/6J mice. Approximately 14% of the total mitochondrial surface was found to be in contact with ER membranes. Notably, these connections exhibit similarities and are found

in the same order of magnitude to what is found in the neurons in the CNS, where 10 to 15% percent of the total mitochondrial surface is occupied by ER (Stoica et al., 2014, 2016). We further characterized the length, spatial distribution, and surface occupation of ER on mitochondria, revealing the dynamic nature of MAM in enteric neurons. Key MAM proteins were expressed in the gut, such as Sigma1R and mitofusin 2 and tethers such as VAPB and PTPIP51 were actively interacting in enteric neurons.

These observations prompted an exploration of ER-mitochondria associations in enteric neurons in a pathological context, using SAMP8 mice — a non-transgenic model of age-related neurodegeneration displaying both

<sup>1</sup>INSERM, Centre de Recherche en Transplantation et Immunologie, UMR 1064, ITUN, Nantes, France

<sup>2</sup>Nantes Université, INSERM, CHU Nantes, The Enteric Nervous System in Gut and Brain Disorders, Nantes, France

\*These authors have equally contributed to this work

Received February 1, 2024. Revised March 12, 2024. Accepted March 18, 2024

## Corresponding Author:

Sébastien Paillusson, Nantes Université, INSERM, CHU Nantes, The Enteric Nervous System in Gut and Brain Disorders, Nantes, France.  
Email: sebastien.paillusson@univ-nantes.fr



CNS and GI dysfunctions. Our investigation revealed constipation in 8-month-old SAMP8 mice compared to control animals. Concurrently, SAMP8 enteric neurons exhibited an increased number of ER-mitochondria contacts compared to control mice, accompanied by a reduction in PGP9.5 and in HuD positive cells by 75% and 53%, respectively, indicative of enteric neuronal loss. Interestingly, no difference in ER-mitochondria contacts was observed in longitudinal smooth muscle cells. We also found that calcium homeostasis we altered in SAMP8 mice enteric neurons. Because the non-transgenic SAMP8 model recapitulates some of the complexity found in patients, it is not easy to draw a direct link between MAM dysfunction and slow transit as symptoms cannot be solely attributed to MAM alteration. However, our findings are rather suggesting that the observed slow transit constipation in SAMP8 mice is the result of neuronal dysfunction leading to the loss of enteric neurons, rather than a functional defect in smooth muscle cells. This indicates that damage made to MAM in the enteric neurons could contribute to altered gut function.

There is now mounting evidence showing that slow transit is an enteric neuropathy rather than a functional disorder (Bassotti et al., 2013), with many studies showing that severely constipated patients exhibit abnormalities in the ENS, such as changes in the neurochemical coding (Wattchow et al., 2008). MAM plays a critical role in regulating key cellular functions in neurons, which include calcium homeostasis, mitochondria biogenesis and oxidative phosphorylation needed for ATP production, neurite growth, synaptic activity and autophagy amongst others (Markovinovic et al., 2022). Therefore, a defect in MAM signalling could trigger neuronal dysfunctions in the ENS leading to gastrointestinal dysfunction. Despite current knowledge, whether and how MAM regulate digestive functions (e.g. intestinal motility and barrier permeability) remains unexamined until our work. One way to understand the direct role that MAM could play in regulating digestive functions through enteric neurons is to modulate (up or down) the amount of tethering between ER and mitochondria in a neuronal-specific manner. In a recent publication, Lu et al. have linked a reduction in ER-mitochondria associations with cardiac dysfunctions during ageing suggesting that damage made to MAM during ageing could affect other key organs like the heart (Lu et al., 2022).

From the literature (Markovinovic et al., 2022), both VAPB-PTPIP51 and mitofusin 2 seem to be the most reliable and interesting tethering proteins to modulate, since they seem to be involved in short and large contact spacing respectively which regulate different MAM functions. Moreover, knock-out animal models for VAPB, PTPIP51 and mitofusin 2 are already available but haven't been studied for gastrointestinal features yet.

Gastrointestinal disorders, in particular constipation, is now considered as a characteristic feature in neurodegenerative disorders such as PD or AD (Rao and Gershon, 2016);

however, the mechanistic make up behind this is still poorly understood as no obvious neuronal loss seems to occur. In the case of PD, growing evidence show that the aggregation of alpha-synuclein, one of the major PD drivers, occurs early in enteric neurons and closely mirrors the synucleinopathy found in central neurons (de Guilhem de Lataillade et al., 2020). Our hypothesis is that pathological drivers like accumulation of alpha-synuclein or A $\beta$  triggers damage to MAM homeostasis in enteric neurons, which alters neuronal function even without any neuronal loss in a similar fashion to the CNS. In order to confirm this hypothesis, a first critical step would be to perform an in-depth characterization of MAM in the ENS of PD patients at different disease stages. To do so, submucosal biopsies from PD patients would be an ideal place to start as they offer fresh tissue without any post-mortem artefacts, and multiple samples could be taken from the same patient to perform a comprehensive study, as one biopsy could be used for electronic microscopy, one for proximity ligation assay, one for IHC and a last one for western blotting, giving a multiple range of assays to draw a reliable conclusion. Aiming in the same direction, more research is needed to characterize whether MAM biology is affected both in the CNS and the ENS during neurodegenerative disorders like PD, and if so in what time scale. Therefore restoring the MAM in neurodegenerative disorders seems to be a way forward in alleviating disease symptoms, and some clinical trials have even reached phase II for PD with this approach (Payne et al., 2020). However, because of the complexity of the damage to MAM in a given disease, it is often difficult to determine whether ER-mitochondria interactions would benefit from being tightened or loosened, making it important to understand the nature of the dysfunction in a particular patient. Using enteric biopsies to monitor MAM status and changes in tethering proteins in a patient could help to determine the individual requirements for a modulating treatment for MAM dysfunction.

### Acknowledgements


We acknowledge the graphical abstract was partly generated using BioRender software.

### Declaration of Conflicting Interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

### Funding

The author(s) disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: This work was supported by the Fédération pour la Recherche sur le Cerveau (FRC), Association AMADYS, Conseil Régional des Pays de la Loire, Agence Nationale de la Recherche, (grant number PO CCI 2014 FR16M20P008, ANR-16-IDEX-0007).

**ORCID iD**Sébastien Paillusson  <https://orcid.org/0000-0001-7013-5582>**References**

- Bassotti G, Villanacci V, Crețoiu D, Crețoiu SM, Becheanu G (2013). Cellular and molecular basis of chronic constipation: Taking the functional/idiopathic label out. *World J Gastroenterol* *19*, 4099–4105. doi: 10.3748/wjg.v19.i26.4099
- de Guilhem de Lataillade A, Lebouvier T, Noble W, Leclair-Visonneau L, Derkinderen P (2020). Enteric synucleinopathy: From trendy concept to real entity. *Free Neuropathol* *1*, 1–26. doi: 10.17879/freeneuropathology-2020-2920
- Delfino G, Briand JB, Oullier T, Nienkemper L, Greig J, Véziers J, Neunlist M, Derkinderen P, Paillusson S (2024). Characterisation of mitochondria-associated ER membranes in the enteric nervous system under physiological and pathological conditions. *Am J Physiol Gastrointest Liver Physiol* *326*, G330–G343. doi: 10.1152/ajpgi.00224.2023
- Lu X, Gong Y, Hu W, Mao Y, Wang T, Sun Z, Su X, Fu G, Wang Y, Lai D (2022). Ultrastructural and proteomic profiling of mitochondria-associated endoplasmic reticulum membranes reveal aging signatures in striated muscle. *Cell Death Dis* *13*, 296. doi: 10.1038/s41419-022-04746-4
- Markovinovic A, Greig J, Martín-Guerrero SM, Salam S, Paillusson S (2022). Endoplasmic reticulum-mitochondria signaling in neurons and neurodegenerative diseases. *J Cell Sci* *135*, jcs248534. doi: 10.1242/jcs.248534
- Payne T, Sassani M, Buckley E, Moll S, Anton A, Appleby M, Maru S, Taylor R, McNeill A, Hoggard N, et al. (2020). Ursodeoxycholic acid as a novel disease-modifying treatment for Parkinson's disease: Protocol for a two-centre, randomised, double-blind, placebo-controlled trial, the “UP” study. *BMJ Open* *10*, e038911. doi: 10.1136/bmjopen-2020-038911
- Rao M, Gershon MD (2016). The bowel and beyond: The enteric nervous system in neurological disorders. *Nat Rev Gastroenterol Hepatol* *13*, 517–528. doi: 10.1038/nrgastro.2016.107
- Stoica R, De Vos KJ, Paillusson S, Mueller S, Sancho RM, Lau KF, Vizcay-Barrena G, Lin WL, Xu YF, Lewis J, et al. (2014). ER-mitochondria associations are regulated by the VAPB-PTPIP51 interaction and are disrupted by ALS/FTD-associated TDP-43. *Nat Commun* *5*, 3996. doi: 10.1038/ncomms4996
- Stoica R, Paillusson S, Gomez-Suaga P, Mitchell JC, Lau DH, Gray EH, Sancho RM, Vizcay-Barrena G, De Vos KJ, Shaw CE, et al. (2016). ALS/FTD-associated FUS activates GSK-3 $\beta$  to disrupt the VAPB-PTPIP51 interaction and ER-mitochondria associations. *EMBO Rep* *17*, 1326–1342. doi: 10.15252/embr.201541726
- Wattchow D, Brookes S, Murphy E, Carbone S, de Fontgalland D, Costa M (2008). Regional variation in the neurochemical coding of the myenteric plexus of the human colon and changes in patients with slow transit constipation. *Neurogastroenterol Motil* *20*, 1298–1305. doi: 10.1111/j.1365-2982.2008.01165.x